



SLOVENSKI STANDARD

SIST EN ISO 21569:2005

01-oktober-2005

Živila - Analitske metode za odkrivanje gensko spremenjenih organizmov in njihovih produktov - Kvalitativne metode na osnovi nukleinske kisline (ISO 21569:2005)

Foodstuffs - Methods of analysis for the detection of genetically modified organisms and derived products - Qualitative nucleic acid based methods (ISO 21569:2005)

iTeh STANDARD PREVIEW

Lebensmittel - Verfahren zum (Nachweis von gentechnisch modifizierten Organismen und ihren Produkten - Qualitative auf Nukleinsäuren basierende Verfahren (ISO 21569:2005)

SIST EN ISO 21569:2005

Produits alimentaires - Méthodes d'analyse pour la détection des organismes génétiquement modifiés et des produits dérivés - Méthodes qualitatives basées sur l'utilisation des acides nucléiques (ISO 21569:2005)

Ta slovenski standard je istoveten z: EN ISO 21569:2005

ICS:

67.050	Splošne preskusne in analizne metode za živilske proizvode	General methods of tests and analysis for food products
--------	--	---

SIST EN ISO 21569:2005

en

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[SIST EN ISO 21569:2005](#)

<https://standards.iteh.ai/catalog/standards/sist/4b0aeb49-39da-4bae-8ed4-f09f2577a058/sist-en-iso-21569-2005>

EUROPEAN STANDARD
NORME EUROPÉENNE
EUROPÄISCHE NORM

EN ISO 21569

June 2005

ICS 67.050

English version

Foodstuffs - Methods of analysis for the detection of genetically modified organisms and derived products - Qualitative nucleic acid based methods (ISO 21569:2005)

Produits alimentaires - Méthodes d'analyse pour la détection des organismes génétiquement modifiés et des produits dérivés - Méthodes qualitatives basées sur l'utilisation des acides nucléiques (ISO 21569:2005)

Lebensmittel - Verfahren zum Nachweis von gentechnisch modifizierten Organismen und ihren Produkten - Qualitative auf Nucleinsäuren basierende Verfahren (ISO 21569:2005)

This European Standard was approved by CEN on 6 June 2005.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the Central Secretariat or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the Central Secretariat has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: rue de Stassart, 36 B-1050 Brussels

EN ISO 21569:2005 (E)

Foreword

This document (EN ISO 21569:2005) has been prepared by Technical Committee CEN/TC 275 "Food analysis - Horizontal methods", the secretariat of which is held by DIN, in collaboration with Technical Committee ISO/TC 34 "Agricultural food products".

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by December 2005, and conflicting national standards shall be withdrawn at the latest by December 2005.

According to the CEN/CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.

iTeh STANDARD PREVIEW (standards.iteh.ai)

[SIST EN ISO 21569:2005](https://standards.iteh.ai/catalog/standards/sist/4b0acb49-39da-4bae-8ed4-f09f2577a058/sist-en-iso-21569-2005)

<https://standards.iteh.ai/catalog/standards/sist/4b0acb49-39da-4bae-8ed4-f09f2577a058/sist-en-iso-21569-2005>

INTERNATIONAL
STANDARD

ISO
21569

First edition
2005-06-15

**Foodstuffs — Methods of analysis for the
detection of genetically modified
organisms and derived products —
Qualitative nucleic acid based methods**

*Produits alimentaires — Méthodes d'analyse pour la détection des
organismes génétiquement modifiés et des produits dérivés —
Méthodes qualitatives basées sur l'utilisation des acides nucléiques*

iTeh STANDARD PREVIEW
(standards.iteh.ai)

SIST EN ISO 21569:2005

<https://standards.iteh.ai/catalog/standards/sist/4b0acb49-39da-4bae-8ed4-f09f2577a058/sist-en-iso-21569-2005>



Reference number
ISO 21569:2005(E)

© ISO 2005

ISO 21569:2005(E)**PDF disclaimer**

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

iTeh STANDARD PREVIEW (standards.iteh.ai)

[SIST EN ISO 21569:2005](https://standards.iteh.ai/catalog/standards/sist/4b0acb49-39da-4bae-8ed4-f09f2577a058/sist-en-iso-21569-2005)

<https://standards.iteh.ai/catalog/standards/sist/4b0acb49-39da-4bae-8ed4-f09f2577a058/sist-en-iso-21569-2005>

© ISO 2005

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Contents

Page

Foreword.....	iv
Introduction	v
1 Scope	1
2 Normative references	1
3 Terms and definitions.....	1
4 Principle of the method	2
5 Reagents	2
6 Apparatus and equipment	3
7 Procedure	3
8 Interpretation.....	6
9 Expression of results and quality assurance	6
10 Test report	7
Annex A (informative) Target-taxon-specific methods.....	8
Annex B (informative) Screening methods.....	24
Annex C (informative) Construct-specific methods	40
Annex D (informative) Event-specific methods.....	62
Bibliography	67

ISO 21569:2005(E)**Foreword**

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

ISO 21569 was prepared by the European Committee for Standardization (CEN) Technical Committee CEN/TC 275, *Food analysis — Horizontal methods*, in collaboration with Technical Committee ISO/TC 34, *Food products*, in accordance with the Agreement on technical cooperation between ISO and CEN (Vienna Agreement).

iTeh STANDARD PREVIEW **(standards.iteh.ai)**

[SIST EN ISO 21569:2005](https://standards.iteh.ai/catalog/standards/sist/4b0aeb49-39da-4bae-8ed4-f09f2577a058/sist-en-iso-21569-2005)

<https://standards.iteh.ai/catalog/standards/sist/4b0aeb49-39da-4bae-8ed4-f09f2577a058/sist-en-iso-21569-2005>

Introduction

The search for a genetically modified origin of ingredients is performed by means of the following successive (or simultaneous) steps. After sample collection, nucleic acids are extracted from the test portion. Extracted nucleic acids can be further purified, simultaneously or after the extraction process. Afterwards, they are quantified (if necessary), diluted (if necessary) and subjected to analytical procedures (such as PCR). These steps are detailed in this International Standard and in the following series of International Standards with the general title *Foodstuffs — Methods of analysis for the detection of genetically modified organisms and derived products*:

- *Sampling* (ISO 21568);
- *Quantitative nucleic acid based methods* (ISO 21570);
- *Nucleic acid extraction* (ISO 21571).

Further information about general requirements and definitions involving the steps cited above are collected in ISO 24276.

The qualitative detection of DNA target sequences is performed in order to obtain a yes or no answer to the question whether a certain target sequence is detected or not relative to appropriate controls and within the detection limits of the analytical method used and test portion analysed.

The specificity of the methods, as described in Annexes A to D, ranges from screening methods to detect common DNA sequences characteristic of GMOs, to specific identification of a genetic construct or a specific transformation event.

The International Organization for Standardization (ISO) draws attention to the fact that it is claimed that compliance with this document may involve the use of a patent concerning the PCR technology.

ISO takes no position concerning the evidence, validity and scope of this patent right.

ISO has been informed that Applied Biosystems, Roche Molecular Systems, Inc. and F. Hoffman La Roche Ltd. hold patent rights concerning the PCR technology. The companies have assured the ISO that they are willing to negotiate licences under reasonable and non-discriminatory terms and conditions with applicants throughout the world. In this respect, the statements of the holders of these patent rights are registered with ISO. Information may be obtained from:

Licensing Department

Applied Biosystems
850 Lincoln Centre Drive
Foster City, CA 94404
USA

and

Roche Molecular Systems, Inc.
Licensing Department
1145 Atlantic Avenue
Alameda, CA 94501
USA

ISO 21569:2005(E)

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights other than those identified above. ISO shall not be held responsible for identifying any or all such patent rights.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

SIST EN ISO 21569:2005

<https://standards.iteh.ai/catalog/standards/sist/4b0acb49-39da-4bae-8ed4-f09f2577a058/sist-en-iso-21569-2005>

Foodstuffs — Methods of analysis for the detection of genetically modified organisms and derived products — Qualitative nucleic acid based methods

1 Scope

This International Standard describes the procedure to qualitatively detect genetically modified organisms (GMOs) and derived products by analysing the nucleic acids extracted from the sample under study. The main focus is on polymerase chain reaction (PCR) based amplification methods.

It gives general requirements for the specific detection and identification of target nucleic acid sequences (DNA) and for the confirmation of the identity of the amplified DNA sequence.

Guidelines, minimum requirements and performance criteria laid down in this International Standard are intended to ensure that comparable, accurate and reproducible results are obtained in different laboratories.

This International Standard has been established for food matrices, but could also be applied to other matrices (e.g. feed and plant samples from the environment).

Specific examples of methods are provided in Annexes A to D.

[SIST EN ISO 21569:2005](https://standards.iteh.ai/catalog/standards/sist/4b0aeb49-39da-4bae-8ed4-f09f2577a058/sist-en-iso-21569-2005)

[https://standards.iteh.ai/catalog/standards/sist/4b0aeb49-39da-4bae-8ed4-](https://standards.iteh.ai/catalog/standards/sist/4b0aeb49-39da-4bae-8ed4-f09f2577a058/sist-en-iso-21569-2005)

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 21571:2005, *Foodstuffs — Methods of analysis for the detection of genetically modified organisms and derived products — Nucleic acid extraction*

ISO 24276:—¹⁾, *Foodstuffs — Nucleic acid based methods of analysis for the detection of genetically modified organisms and derived products — General requirements and definitions*

3 Terms and definitions

For the purposes of this document, the terms and definitions given in ISO 24276 apply.

1) To be published.

ISO 21569:2005(E)

4 Principle of the method

4.1 General

Qualitative analysis consists of specific detection of target nucleic acid sequences in the test samples. Each method shall specify the target sequence.

A qualitative result shall clearly demonstrate the presence or absence of the genetic element under study, relative to appropriate controls and within the detection limits of the analytical method used and test portion analysed.

4.2 PCR amplification

Amplification of the target sequence occurs *in vitro* through a reaction catalysed by a DNA polymerase in the presence of oligonucleotide primers and deoxyribonucleoside triphosphates in a defined reaction buffer^{[1], [2]}. An important prerequisite for the amplification of the target sequence is that the reaction mixture contains no polymerase inhibitors. Amplification of the DNA is a cyclical process consisting of

- denaturation of the double-stranded DNA into single-stranded nucleic acid by means of heating,
- annealing of the primers to the target sequence at a suitable temperature, and
- extension of the primers, which are bound to both single strands, by a DNA polymerase suitable for PCR, at an appropriate temperature.

4.3 Detection and confirmation of PCR products

PCR products are detected by gel electrophoresis or an appropriate alternative, if necessary, after isolation by means of a suitable separation procedure.

The identity of any detected target sequence can be verified by an appropriate technique (e.g. by restriction enzyme analysis, by hybridization or by DNA sequence analysis).

In the case of real-time PCR analysis, amplification and detection occur simultaneously.

5 Reagents

It is generally advisable to store the reaction solutions required for the analytical method at approximately $-20\text{ }^{\circ}\text{C}$ if not specified otherwise.

It may also be appropriate to aliquot the reaction solutions required for the analytical method in order to avoid subjecting them to repeated freeze-thaw cycles, and/or to reduce chances of cross contamination.

5.1 Target DNA/control

5.2 Water

5.3 Deoxyribonucleoside triphosphate (dNTP) solution, containing dATP, dCTP, dGTP, and dTTP or dUTP.

NOTE The use of dUTP can interfere with restriction enzyme analyses of PCR products.

5.4 PCR buffer solution

The PCR buffer solution is usually delivered with the DNA polymerase, which may or may not include $MgCl_2$ in a concentration specified by the manufacturer. The final $MgCl_2$ concentrations are method specific and are therefore listed in each annex. Ready-to-use reagents may be commercially available. The manufacturer's instructions for use should be considered.

5.5 Thermostable DNA polymerase

5.6 Forward primer

5.7 Reverse primer

6 Apparatus and equipment

See ISO 24276 and Annexes A to D for details.

7 Procedure

7.1 Quality, integrity and amplifiability of nucleic acid extracts

The nucleic acid solution shall be pure enough for subsequent analysis^[3]. The quality and amount of nucleic acid extracted using a given method on a given matrix shall be both repeatable and reproducible.

NOTE The quality, integrity and amount of the DNA template influences the outcome of the PCR, and hence the analytical results obtained. The limit of detection of a specific method may therefore depend on whether the material to be analysed has been processed or refined, and on the degree of degradation of the DNA therein.

Nucleic acids for use in PCR should be substantially free of PCR inhibitors^[4]. PCR inhibition controls shall be included as described in ISO 24276.

7.2 Performance criteria

General performance criteria are described in ISO 24276.

The values for the performance characteristics are given for each method as outlined in Annexes A to D and should take into account the genome sizes; see Reference [5].

The reaction conditions, especially the $MgCl_2$ concentration and the thermocycling conditions should be optimized for every primer pair and/or system. When any primer system is used for the first time, it is necessary to demonstrate beforehand that the cycle conditions chosen for the particular matrix to be studied avoid undesirable competitive products that would otherwise reduce the sensitivity of the PCR detection.

In an optimal reaction, less than 40 cycles are required to amplify ≥ 10 target molecules to produce a product that is readily detectable by standard methods. As the cycle number increases, non-specific products could accumulate. The optimized PCR should be able to amplify in 40 PCR cycles from a pure reference sample of 100 copies of template DNA enough copies of the PCR product to be detectable. The characteristic temperature/time profile for each primer system and the reaction mixture appropriate for the apparatus used and the number of cycles shall be strictly adhered to.

In general, the specificity of the reaction should be enhanced as much as possible (e.g. by using hot-start PCR). Hot-start PCR is recommended as a means of reducing side reactions such as the amplification of non-target sequences in background DNA (mispriming) and primer-oligomerization (it thus increases specificity).

The values derived from the validation study may not be applicable to analyte concentration ranges and matrices other than given in the respective annexes.

ISO 21569:2005(E)**7.3 Aspects of PCR design****7.3.1 General**

Because the performance of each specific PCR should be comparable with other specific PCRs, the following aspects of PCR design shall be taken into account.

7.3.2 Size of PCR products

The size of the target sequence shall be selected to match the range of molecular mass available in the nucleic acid extract being analysed.

EXAMPLE For highly degraded DNA from processed foodstuffs, the size of the PCR product should ideally be in the range of 60 bp to 150 bp. For raw materials, a broader range of PCR products up to, for example, 250 bp is applicable.

However, if prior experimental studies are carried out to determine the validity of primer sets yielding different sized PCR products, these may be used on the matrix for which they have been validated.

7.3.3 Primers**7.3.3.1 General**

Primer sequence information is included in Annexes A to D.

7.3.3.2 Primer design

iTeh STANDARD PREVIEW
(standards.iteh.ai)

The primer sequences should preferably have the following characteristics wherever practicable:

- length of each primer: 18 to 30 nucleotides; [SIST EN ISO 21569:2005](https://standards.iteh.ai/catalog/standards/sist/4b0acb49-39da-4bae-8ed4-091818181818/sist-en-iso-21569-2005)
- optimal annealing temperature $\approx 60\text{ }^{\circ}\text{C}$ (should be established experimentally), i.e. estimated melting temperature $\leq 65\text{ }^{\circ}\text{C}$;
- GC:AT ratio = 50:50 if possible, or else as close to this ratio as possible;
- high internal stability (avoid concentration of Gs and Cs in short segments of primers);
- minimal 3' end complementarity to avoid primer-dimer formation;
- minimal secondary structure;
- minimal dimer formation with specific detection probe(s) designed for the PCR.

Software packages are available to help with primer design.

7.3.3.3 Validation of primers**7.3.3.3.1 General**

The ability of the primers to detect the target sequence shall be validated.

Primer validation should be carried out in two steps: a first theoretical evaluation, and a second experimental evaluation.

7.3.3.3.2 Theoretical evaluation of the specificity

Theoretical evaluation shall as a minimum be carried out by performing a sequence similarity search (e.g. FastA, Blast^{®2)}) against one of the major nucleic acid sequence databases (e.g. EMBL, GenBank^{®2)}). Homologous gene sequences may be retrieved from the sequence databases and aligned to obtain an estimate of the chance of finding similar sequences in the target taxon or other organisms.

7.3.3.3.3 Experimental evaluation of the specificity

Irrespective of the design criteria used, the specificity of primers shall always be experimentally evaluated to confirm the primers' ability to discriminate between the target and closely related non-target sequences.

Primers designed to detect taxon-specific target sequences should be shown to detect these sequences reliably in a representative number of different members of the taxon.

7.4 PCR target descriptions

For the qualitative detection and identification of GMOs, various PCR tests may be performed, depending on the type of matrices under study and/or the requirements of the analysis. These analyses may be directed at sequences specific for target taxa, genetic constructs and transformation events, as well as elements suitable for screening purposes.

7.5 Controls

Because of the risk of obtaining false positive and/or false negative results, appropriate controls shall be included in each diagnostic PCR assay (see ISO 24276).

If available and appropriate, certified reference materials should be used as positive and negative controls.

7.6 PCR set-up, detection and confirmation of PCR products

Annexes A to D give details on the specific PCR procedure steps.

NOTE In the case of detection of the PCR products by gel electrophoresis, the size of the PCR products can be estimated using a suitable DNA size marker of known length to run in parallel with the PCR products under test.

It may be desirable in some cases to confirm a positive or negative result for a particular genetic modification. This may be achieved by employing primers to an alternative target sequence; this is particularly suitable for confirmation of screening test results.

A positive identification of the specific target DNA sequence may be confirmed by an appropriate method other than size determination of the PCR product, for example

- by hybridization of the PCR product with specific probes, or
- by carrying out restriction analyses of the PCR product; the length of the resulting fragments has to correspond to the expected length of the target DNA sequence after restriction, or
- by sequencing of the PCR product, or
- other equivalent confirmation.

2) Blast and GenBank are examples of suitable products available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of these products. Equivalent products may be used if they can be shown to give the same results.