

---

**Kakovost vode - Ugotavljanje strupenosti snovi s sladkovodnimi ribami v podaljšanem testu - Metoda za vrednotenje vplivov snovi na hitrost rasti šarenke (Oncorhynchus mykiss Walbaum (Teleostei, Salmonidae))**

Water quality -- Determination of the prolonged toxicity of substances to freshwater fish -- Method for evaluating the effects of substances on the growth rate of rainbow trout (Oncorhynchus mykiss Walbaum (Teleostei, Salmonidae))

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

Qualité de l'eau -- Détermination de la toxicité à long terme de substances vis-à-vis d'un poisson d'eau douce -- Méthode d'évaluation des effets de substances sur le taux de croissance de la truite arc-en-ciel (Oncorhynchus mykiss Walbaum (Teleostei, Salmonidae))

**Ta slovenski standard je istoveten z: ISO 10229:1994**

---

**ICS:**

13.060.70	Preiskava bioloških lastnosti vode	Examination of biological properties of water
-----------	------------------------------------	---

**SIST ISO 10229:1997**

**en**

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

SIST ISO 10229:1997

<https://standards.iteh.ai/catalog/standards/sist/7db0f7f5-8499-42d7-98b4-c490c4b92d28/sist-iso-10229-1997>

INTERNATIONAL  
STANDARDISO  
10229First edition  
1994-12-01

---

---

**Water quality — Determination of the prolonged toxicity of substances to freshwater fish — Method for evaluating the effects of substances on the growth rate of rainbow trout [*Oncorhynchus mykiss* Walbaum (Teleostei, Salmonidae)]**

[SIST ISO 10229:1997](https://standards.iteh.ai/catalog/standards/sist/7db0f7f5-8499-42d7-98b4-)

<https://standards.iteh.ai/catalog/standards/sist/7db0f7f5-8499-42d7-98b4->

*Qualité de l'eau — Détermination de la toxicité prolongée de substances vis-à-vis d'un poisson d'eau douce — Méthode d'évaluation des effets de substances sur le taux de croissance de la truite arc-en-ciel [*Oncorhynchus mykiss* Walbaum (Teleostei, Salmonidae)]*



Reference number  
ISO 10229:1994(E)

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

International Standard ISO 10229 was prepared by Technical Committee ISO/TC 147, *Water quality*, Subcommittee SC 5, *Biological methods*.

Annexes A and B of this International Standard are for information only.

ITeH STANDARD PREVIEW  
(standards.iteh.ai)  
SIST ISO 10229:1997  
https://standards.iteh.ai/catalog/standards/sist/75-8499-42d7-98b4-c490c4b92d28/sist-iso-10229-1997

© ISO 1994

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from the publisher.

International Organization for Standardization  
Case Postale 56 • CH-1211 Genève 20 • Switzerland

Printed in Switzerland

## Introduction

This International Standard describes methods of determining the long-term toxicity of substances to freshwater fish. The methods include the determination of lethal and sublethal responses. In all cases, the specified use of one species does not preclude the use of other species. It is possible for the methods presented here to be used for other species of freshwater fish, provided that appropriate modifications to, for example, quality of water used for dilution, temperature conditions and feeding rates are made and the relationships between growth rates, feeding rates and conversion efficiencies are established.

Within this International Standard, a choice may be made between semi-static and flow-through methods. These have already been described for the determination of the acute lethal toxicity of substances to a freshwater fish in ISO 7346-2 and ISO 7346-3 respectively. The choice of the method of presentation of the test solution depends on two factors: the nature of the test substance and the form of the test method.

The flow-through method can be used for most types of substances, including those unstable in water, but the concentrations of the test substance are determined wherever possible. The semi-static method can be used for testing those substances whose concentrations can be maintained satisfactorily throughout the test by renewal of the solutions every 24 h.

To assist in the preparation and maintenance of concentrations of substances which may invoke the chosen responses in the fish at concentrations close to that of their aqueous solubility, a small volume of solvent is sometimes used, as specified in the methods.

## iTeh STANDARD PREVIEW

This page intentionally left blank.  
(standards.iteh.ai)

[SIST ISO 10229:1997](https://standards.iteh.ai/catalog/standards/sist/7db0f7f5-8499-42d7-98b4-c490c4b92d28/sist-iso-10229-1997)

<https://standards.iteh.ai/catalog/standards/sist/7db0f7f5-8499-42d7-98b4-c490c4b92d28/sist-iso-10229-1997>

# Water quality — Determination of the prolonged toxicity of substances to freshwater fish — Method for evaluating the effects of substances on the growth rate of rainbow trout [*Oncorhynchus mykiss* Walbaum (Teleostei, Salmonidae)]

## 1 Scope

This International Standard specifies a method for the determination of the long-term sublethal toxicity of substances soluble in water under specified conditions to a species of anadromous fish [*Oncorhynchus mykiss* Walbaum (Teleostei, Salmonidae) — common name, rainbow trout] in fresh water.

NOTE 1 “Substances” is considered to include pure chemicals tested singly or in mixtures, waste waters, process waters or other waters whose sublethal toxicity to fish it is necessary to determine. A range of concentrations of waste and process waters should be tested (see 8.2).

The response measured is the change in rate of growth of fish exposed to the test substance over periods of 14 d and 28 d (see [1] in annex B).

NOTE 2 The mass of each fish at the start of the test, the period of exposure, the feeding regime and temperature range have been chosen to ensure that the control fish will be growing exponentially throughout the test.

The method is applicable for assigning, for each test substance, a category of prolonged sublethal toxicity to *Oncorhynchus mykiss* under the test conditions.

The results are insufficient by themselves to define water quality limits for environmental protection.

It is possible for the method to be adapted for use with other freshwater fish and marine and brackish water fish with appropriate modification of the test conditions, particularly with respect to the tempera-

ture, the quantity and quality of the dilution water, the food and the fish-marking technique.

## 2 Normative references

The following standards contain provisions which, through reference in this text, constitute provisions of this International Standard. At the time of publication, the editions indicated were valid. All standards are subject to revision, and parties to agreements based on this International Standard are encouraged to investigate the possibility of applying the most recent editions of the standards indicated below. Members of IEC and ISO maintain registers of currently valid International Standards.

ISO 7346-2:1984, *Water quality — Determination of the acute lethal toxicity of substances to a freshwater fish (Brachydanio rerio Hamilton-Buchanan (Teleostei, Cyprinidae)) — Part 2: Semi-static method.*

ISO 7346-3:1984, *Water quality — Determination of the acute lethal toxicity of substances to a freshwater fish (Brachydanio rerio Hamilton-Buchanan (Teleostei, Cyprinidae)) — Part 3: Flow-through method.*

## 3 Principle

Determination, under specified conditions, of the concentrations at which a substance statistically and significantly reduces the rate of growth of a test population of *Oncorhynchus mykiss* after exposure periods of 14 d and 28 d to that substance in the ambient water.

The test is carried out in two stages:

- a) a preliminary test, carried out using rainbow trout as the test species but in accordance with ISO 7346-2 or ISO 7346-3 (although the rainbow trout is not specifically named in these parts of ISO 7346), which determines the concentrations at which the substance is lethal to 50 % of a test population after exposure to the substance in the ambient water for 96 h: this median lethal concentration is designated as 96 h LC50 and serves to determine the range of concentrations for the final test (see note 11 to 8.1).
- b) a final test, using 16 fish per concentration, in which the results are recorded as both the lowest observed effect concentration (LOEC) and the no observed effect concentration (NOEC). They are prefixed by an indication of the exposure period concerned, for example "28 d LOEC". The data may also be fitted to a regression model in order to calculate the IC10 (inhibition concentration for a 10 % effect), or a similar statistic.

Where analytical data show that median test concentrations remain relatively constant (i.e. within 20 % of the nominal values) throughout the test, the method allows the use of either measured or nominal concentrations in the statements on LOEC and NOEC. Where such analyses show that the concentrations present remain relatively constant (i.e.  $\pm 20\%$  of the median) but are less than 80 % or greater than 120 % of the nominal values, then the analytical values derived from this method are used in presenting the LOEC and NOEC. Where no evidence is available to show that the test concentrations remained at an acceptable level throughout the test period or where it is known that the concentrations of the test chemical have changed significantly (i.e. by greater than 20 % of the median) at any stage during the test, then irrespective of whether or not chemical analytical data are available the LOEC and NOEC cannot be defined using this test method. In these cases, the test is not necessarily invalidated but it can only be stated that the LOEC of the substance is  $\leq x$  mg/l, and that the NOEC of the substance is  $\leq y$  mg/l, the values  $x$  and  $y$  given being estimated from the nominal concentrations used.

NOTE 3 Wherever possible, it is very important to determine the concentrations of the test substance to which the fish are exposed.

Fish can be individually recognized throughout the test if the freeze-branding technique or an equivalent method (e.g. tagging) is used for marking, but this is not mandatory.

## 4 Test organism

The recommended test species is *Oncorhynchus mykiss* Walbaum (Teleostei, Salmonidae), commonly known as the rainbow trout. Test fish shall be selected from a population of a single stock. This stock shall have been acclimatized and, in any case, maintained for at least 2 weeks prior to the test in dilution water, continuously aerated (using bubbled air), under conditions of water quality, feeding regime and illumination similar to those used in the test.

Except where otherwise stated (see clause 8), the fish shall be fed at a minimum rate of 1 % of their wet body mass per day during any pre-acclimation and acclimation period and 4 % of their mass per day during the final test (see 8.2).

NOTE 4 This latitude in feeding rate during the pretest period enables the fish to be grown to, or held at, a suitable size (see below) for the start of the test.

The food shall be a dry proprietary salmonid fry food and shall be divided into two equal portions and given to the fish in two feeds per day separated by at least 5 h. After exposure of 14 d to the test substance, when the fish are weighed again, the rations shall be recalculated for each test vessel.

At the start of the test, each fish shall have a mass of not less than 3 g and not more than 5 g (but see 8.2). For the whole batch of fish used in the test, the range in individual masses at the start of the test shall lie within  $\pm 10\%$  of the arithmetic mean of the masses.

Test fish shall be free of overt disease or visible malformation. They shall not receive treatment for disease during the test or in the 2 weeks preceding the test.

Maintain the fish in environmental conditions suitable for rainbow trout.

NOTE 5 Fish cultivation manuals should be consulted, for example [2] in annex B.

## 5 Reagents

Use only reagents of recognized analytical grade. The water used for the preparation of stock solutions shall be glass-distilled or deionized water of at least equivalent purity.

### 5.1 Dilution water

The dilution water shall be suitable for the long-term survival and growth of the test fish. The average pH



of the dilution water shall be within the range 6,7 to 8,5, but during a given test the pH of the dilution water shall not vary by more than  $\pm 0,2$  pH units from the mean value.

NOTE 6 Such a pH range can be expected to be associated with a bicarbonate alkalinity of 1 mg/l to 100 mg/l  $\text{CaCO}_3$  and total hardness values of 20 mg/l to 300 mg/l  $\text{CaCO}_3$ , but these ranges will depend on the precise constitution of the dilution water and the time available for equilibration of the water. A suitable artificial water can be made with distilled water, as follows:

294,0 mg/l  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$

123,3 mg/l  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$

63,0 mg/l  $\text{NaHCO}_3$

5,5 mg/l KCl

## 5.2 Stock solutions of test substances

Prepare a stock solution of the test substance by dissolving a known amount of test substance in a defined volume of dilution water, deionized water or glass-distilled water.

NOTE 7 In the case where the test "substance" is a waste water, process water or other water (see clause 1, note 1) and not one or more pure chemicals, these waters are to be taken as the stock solution.

Prepare or obtain the stock solution as frequently as necessary to maintain stable solutions. Although the use of solvents is not desirable, to enable stock solutions to be prepared and to assist in their transfer to the test vessels, it shall be permissible to dissolve or disperse substances of low aqueous solubility by suitable means, for example ultrasonic devices or the use of organic solvents of low toxicity to fish. If any such organic solvent is used, its concentration in the test solution shall not exceed 0,1 ml/l (and preferably be below 0,01 ml/l), and two sets of controls, one containing solvent at the maximum concentration used in any test vessel and one without solvent or test substance, shall be included.

NOTE 8 It is preferable that the same concentration of solvent be used in all test vessels.

## 5.3 Test solutions

Prepare test solutions by adding appropriate amounts of the stock solution of the test substance (5.2) to the dilution water to give the required concentrations (8.2). It is recommended that when a stock solution

is prepared in distilled or deionized water, no more than 100 ml of stock solution should be added per 10 litres of dilution water. Wherever possible, it is very important to determine the concentrations of the test substance in each test vessel, with the exception of waste waters where this may not be possible.

## 6 Apparatus

### 6.1 General

All materials which may come into contact with any liquid into which the fish are to be placed, or with which they may come into contact, shall be inert and shall not absorb the test substance significantly.

Usual laboratory equipment [including a dip-net, made of nylon or of another chemically inert material, for the control vessels and another for all the test vessels (6.2)] shall be used.

6.2 Test vessels, of capacity at least 45 litres. Their sides shall be covered with an opaque material to minimize disturbance of the fish.

Before use, carefully wash new test vessels and then rinse successively with water and the dilution water (5.1). At the end of the test, empty the vessels, clean them, for example with a non-ionic detergent (followed by acid and solvent washes for substances expected to adsorb strongly to the vessel), rinse with water to remove all traces of the test substance and cleaning aid, and dry.

Rinse test vessels with dilution water (5.1) just before use.

6.3 Temperature control, to maintain the temperature of the test solutions and the water in the fish stock tanks in the range 12,5 °C to 17,5 °C and to regulate it within  $\pm 1$  °C.

6.4 Solution-replacement equipment, comprising a dosing and mixing apparatus capable of maintaining the required concentrations of the stock solution in the test vessels to within 10 % of the mean. The apparatus shall be set to renew the test solutions in the vessels at a rate which is sufficient to prevent the concentration of dissolved oxygen in the vessels from falling below 70 % air saturation value (ASV).

### 6.5 Equipment for marking fish

If individual fish are to be marked, one of several techniques is permissible, for example freeze branding.