



**SLOVENSKI STANDARD**  
**SIST ISO 10706:2010**

**01-september-2010**

---

**Kakovost vode - Določevanje dolgotrajne strupenosti snovi z Daphnia magna Straus (Cladocera, Crustacea)**

Water quality - Determination of long term toxicity of substances to Daphnia magna Straus (Cladocera, Crustacea)

**iTeh STANDARD PREVIEW**

Qualité de l'eau - Détermination de la toxicité à long terme de substances vis-à-vis de Daphnia magna Straus (Cladocera, Crustacea)

[SIST ISO 10706:2010](https://standards.iteh.ai/catalog/standards/sist/bdd3e916-1486-438b-94f6-14297bd48a5d/sist-iso-10706-2010)

**Ta slovenski standard je istoveten z: ISO 10706:2000**

---

**ICS:**

13.060.70	Preiskava bioloških lastnosti vode	Examination of biological properties of water
-----------	------------------------------------	---

**SIST ISO 10706:2010**

**en,fr**

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

SIST ISO 10706:2010

<https://standards.iteh.ai/catalog/standards/sist/bdd3e916-1486-438b-94f6-f4297bd48a5d/sist-iso-10706-2010>

INTERNATIONAL  
STANDARD

ISO  
10706

First edition  
2000-04-01

---

---

**Water quality — Determination of long term  
toxicity of substances to *Daphnia magna*  
Straus (*Cladocera*, *Crustacea*)**

*Qualité de l'eau — Détermination de la toxicité à long terme de substances  
vis-à-vis de Daphnia magna Straus (Cladocera, Crustacea)*

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

SIST ISO 10706:2010

<https://standards.iteh.ai/catalog/standards/sist/bdd3e916-1486-438b-94f6-f4297bd48a5d/sist-iso-10706-2010>



Reference number  
ISO 10706:2000(E)

© ISO 2000

**ISO 10706:2000(E)****PDF disclaimer**

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

## iTeh STANDARD PREVIEW (standards.iteh.ai)

SIST ISO 10706:2010

<https://standards.iteh.ai/catalog/standards/sist/bdd3e916-1486-438b-94f6-f4297bd48a5d/sist-iso-10706-2010>

© ISO 2000

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Case postale 56 • CH-1211 Geneva 20  
Tel. + 41 22 749 01 11  
Fax + 41 22 734 10 79  
E-mail [copyright@iso.ch](mailto:copyright@iso.ch)  
Web [www.iso.ch](http://www.iso.ch)

Printed in Switzerland

## Contents

Page

Foreword.....	iv
Introduction.....	v
1 Scope .....	1
2 Normative references .....	1
3 Principle.....	1
4 Test environment.....	1
5 Reagents and materials .....	2
6 Apparatus .....	3
7 Treatment and preparation of samples .....	3
7.1 Special precautions for sampling and transportation of water or effluent samples .....	3
7.2 Preparation of solutions of substances to be tested.....	3
8 Procedure .....	4
8.1 Controls .....	4
8.2 Selection of exposure concentrations.....	4
8.3 Renewal of exposure solutions.....	5
8.4 Introduction of animals into the test system.....	5
8.5 Feeding of organisms .....	5
8.6 Observations and measurements .....	5
8.7 Validity of the test.....	6
9 Data analysis and expression of results.....	6
10 Test report .....	7
Annex A (informative) OECD M4 and M7 media — Preparation and acclimation of Elendt M4 and M7 media.....	9
Annex B (informative) ASTM reconstituted hard fresh water.....	12
Annex C (informative) Data collection sheet .....	13
Annex D (informative) Calculation of time-weighted mean concentration.....	15
Bibliography.....	17

**ISO 10706:2000(E)****Foreword**

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 3.

Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this International Standard may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

International Standard ISO 10706 was prepared by Technical Committee ISO/TC 147, *Water quality*, Subcommittee SC 5, *Biological methods*.

Annexes A, B, C and D of this International Standard are for information only.

**ITC147 STANDARD PREVIEW**  
**(standards.iteh.ai)**

[SIST ISO 10706:2010](https://standards.iteh.ai/catalog/standards/sist/bdd3e916-1486-438b-94f6-f4297bd48a5d/sist-iso-10706-2010)

<https://standards.iteh.ai/catalog/standards/sist/bdd3e916-1486-438b-94f6-f4297bd48a5d/sist-iso-10706-2010>

## Introduction

This International Standard defines a procedure for the determination of the long term sublethal toxicity of chemicals, waters and waste waters to the water-flea *Daphnia magna* Straus. The methodology is adapted from a guideline produced by the Organisation for Economic Co-operation and Development (OECD Guideline 211, see reference [1] in the Bibliography).

## iTeh STANDARD PREVIEW (standards.iteh.ai)

[SIST ISO 10706:2010](https://standards.iteh.ai/catalog/standards/sist/bdd3e916-1486-438b-94f6-f4297bd48a5d/sist-iso-10706-2010)

<https://standards.iteh.ai/catalog/standards/sist/bdd3e916-1486-438b-94f6-f4297bd48a5d/sist-iso-10706-2010>

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

SIST ISO 10706:2010

<https://standards.iteh.ai/catalog/standards/sist/bdd3e916-1486-438b-94f6-f4297bd48a5d/sist-iso-10706-2010>



# Water quality — Determination of long term toxicity of substances to *Daphnia magna* Straus (Cladocera, Crustacea)

**WARNING** — Activated sludge and sewage contain potentially pathogenic organisms. Therefore, appropriate precautions should be taken when handling them. Toxic test compounds and those whose properties are unknown should be handled with care.

## 1 Scope

This International Standard describes a method for the determination of the long term sublethal toxicity to *Daphnia magna* Straus (Cladocera, Crustacea) of:

- a) chemical substances, which are soluble under the conditions of the test, or can be maintained as stable suspensions or dispersions under the conditions of the test,
- b) industrial or sewage effluents, treated or untreated, after decanting, filtration, or centrifugation,
- c) surface or ground waters.

NOTE This International Standard is adapted from OECD Guideline 211<sup>[1]</sup>.

## 2 Normative references

The following normative documents contain provisions which, through reference in this text, constitute provisions of this International Standard. For dated references, subsequent amendments to, or revisions of, any of these publications do not apply. However, parties to agreements based on this International Standard are encouraged to investigate the possibility of applying the most recent editions of the normative documents indicated below. For undated references, the latest edition of the normative document referred to applies. Members of ISO and IEC maintain registers of currently valid International Standards.

ISO 5667-2, *Water quality — Sampling — Part 2: Guidance on sampling techniques*.

ISO 5667-16, *Water quality — Sampling — Part 16: Guidance on biotesting of samples*.

## 3 Principle

*Daphnia magna* females, less than 24 h old, are exposed for a period of 21 days to a test substance, industrial or sewage effluent or surface/ground water added to dilution water in a range of concentrations in a semi-static or flow-through system. The survival of parents is recorded together with the number of live offspring produced per live parent at the end of the test.

The survival is reported and the reproductive output of exposed parents living at the end of the test (expressed in a number of terms) is compared to those of the control parents.

## 4 Test environment

The test atmosphere shall be free from vapours or dusts which may be toxic to *Daphnia magna* Straus.

## ISO 10706:2000(E)

The exposure solutions shall not be aerated.

The dissolved oxygen concentration in the exposure solutions shall be above 3 mg/l and the pH shall be within the range of pH 6 to pH 9 and not change by more than 1,5 pH units for the duration of the test. Hardness shall be above 140 mg/l (as CaCO<sub>3</sub>), which has been demonstrated to be necessary to promote reproductive performance that shall meet the validity criteria (see 8.7).

The photoperiod of the test shall be 16 h of light and 8 h of darkness. The intensity shall be in the range 600 lux to 800 lux but not exceeding 1 200 lux.

The temperature of the test exposures shall be maintained within the range of 18 °C to 22 °C and the temperature for the test shall not vary by more than 2 °C (for instance 18 °C to 20 °C, 19 °C to 21 °C or 20 °C to 22 °C) throughout the test.

## 5 Reagents and materials

**5.1 Test organism:** *Daphnia magna* Straus (*Cladocera, Crustacea*), hereafter referred to as *D. magna*, obtained by ayclical parthenogenesis for at least three generations under specified culture conditions.

The age and the source (including clone, if possible) of the *D. magna* culture shall be indicated in the test report, since the sensitivity of *D. magna* to toxicants can be affected by the source of the culture.

The animals used for the test shall be less than 24 h old and be from the second to fifth brood. The *D. magna* shall be from a healthy stock showing no signs of stress such as mortality > 20 %, presence of males, ehippia, or discoloured animals and there shall be no delay in the production of the first brood.

The stock animals shall be maintained in culture conditions (light, temperature, medium, feeding and animals per unit volume) similar to those in the test. If the culture conditions differ from test conditions, it is recommended that one generation be acclimated under the test conditions for about three weeks so as to avoid stressing the parent animals.

<https://standards.iteh.ai/catalog/standards/sist/bdd3e916-1486-438b-94f6-f4297bd48a5d/sist-iso-10706-2010>

**5.2 Dilution water,** synthetic as described in 5.3 or uncontaminated natural waters with comparable pH and hardness characteristics to these dilution waters for culturing and testing.

**5.3 Synthetic dilution water,** prepared using one of the following methods:

- a) OECD M4 and M7 media (see informative annex A);
- b) ASTM reconstituted hard freshwater (see informative annex B).

The dilution water shall be aerated until the dissolved oxygen concentration has reached 95 % saturation and the pH has stabilized. If necessary, adjust the pH to  $8,0 \pm 0,5$  by adding sodium hydroxide (NaOH) solution or hydrochloric acid (HCl) solution. The dilution water prepared in this way shall not be further aerated before use.

It is recommended that the TOC levels be < 5 mg/l before the addition of algae. The use of dilution water containing EDTA (for instance M4 and M7) is not recommended when testing compounds containing metals because chelation may reduce the toxicity of the compound.

If the test has to be performed for purposes necessitating the use of a dilution water with characteristics differing from those described above, mention is necessary in the test report of the main characteristics (for instance pH, hardness, TOC, COD) of the synthetic dilution water used.

## 6 Apparatus

Ordinary laboratory apparatus and in particular the following.

**6.1 Environmental controls**, for controlling temperature, photoperiod and light intensity.

**6.2 Measuring apparatus and/or instruments**, for measuring dissolved oxygen, pH, hardness, total organic carbon, chemical oxygen demand, light intensity and temperature.

**6.3 Test containers**, of chemically inert material and of sufficient capacity for the tests (for example 50 ml or 100 ml glass test tubes or beakers) and which are clean and uncontaminated.

## 7 Treatment and preparation of samples

### 7.1 Special precautions for sampling and transportation of water or effluent samples

Sampling of water or effluent shall be carried out in accordance with the general procedure specified in ISO 5667-2. Bottles shall be completely filled to exclude air.

The preservation and storage of water or effluent samples shall be carried out in accordance with ISO 5667-16; the following is only a summary. Carry out the toxicity test as soon as possible, ideally within 12 h of collection. If this time interval cannot be observed, cool the sample (0 °C to 4 °C) and test the sample within 24 h. If testing cannot be carried out within 48 h, the sample may be frozen (below –18 °C) for testing within 2 months of collection.

All portions (subsamples) shall be pretreated identically (i.e. if freezing of unstable water is necessary all portions, including those needed for the first day, shall be frozen prior to testing see ISO 5667-16).

Because of the duration of this test (21 days) and the periodic renewal of solutions, a sufficient number of portions of the sample must be frozen to renew test solutions and to repeat the test, if needed (reserve samples). The minimum volume of the frozen portions is dependent upon the toxicity of the sample to be tested. The volume of sample needed for a semi-static test with ten daphnids exposed individually in 100 ml of undiluted sample is 9 l; additional sample will be required for diluted exposures.

If the test is conducted on site or close to the sampling site, fresh samples may be used to replace the test solutions. In this case the variations at the sampling site are incorporated into the test design.

### 7.2 Preparation of solutions of substances to be tested

#### 7.2.1 Preparation of stock solutions

A stock solution of the substance(s) to be tested shall be prepared by dissolving a known quantity of the substance(s) in a specified volume of dilution water, deionized water or distilled water in a glass container. The stock solution shall be prepared immediately before preparing the exposure solutions unless the substance(s) is/are known to be stable in defined storage conditions, in which case the stock solution may be prepared in advance of testing and stored in these conditions.

Stock solutions or suspensions of substance(s) which are poorly soluble in water can be solubilized or dispersed directly in the medium with ultrasonic dispersion, and/or stirring or with solvents or dispersants of low toxicity to *D. magna* as discussed in ISO 5667-16. If a solvent is used, the concentration of the solvent in the stock solution shall be such that the concentration in the highest exposure solution does not exceed 0,1 ml/l.

The use of organic solvents should be avoided. If they are required organic solvents such as acetone, ethanol, methanol, dimethylformamide, triethylene glycol or dispersants such as Cremophor RH40, methylcellulose 0,1 %, and HCO-40 may be used to produce a suitably concentrated stock solution. They are not toxic to *D. magna* at 0,1 ml/l concentrations. No single procedure for the preparation of stock solutions of poorly soluble substances can be recommended due to the differing nature of chemicals.