

---

---

**Meat and meat products — Enumeration of  
*Escherichia coli* — Colony-count technique  
at 44 °C using membranes**

*Viandes et produits à base de viande — Dénombrement des Escherichia coli — Méthode par comptage des colonies obtenues sur membranes à 44 °C*

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

[ISO 6391:1997](#)

<https://standards.iteh.ai/catalog/standards/sist/73a6f77b-4d7d-4bf8-b772-4973f8178074/iso-6391-1997>



## Contents

Page

1	Scope.....	1
2	Normative references .....	1
3	Definitions .....	1
4	Principle .....	2
5	Diluent, culture media and reagents .....	2
6	Apparatus and glassware.....	4
7	Sampling .....	5
8	Preparation of test sample.....	5
9	Procedure.....	5
10	Expression of results .....	7
11	Test report.....	8
Annex A	Bibliography .....	9

<https://standards.iteh.ai/catalog/standards/sis/75a0177b-4d7d-4bf8-b772-4973f8178074/iso-6391-1997>

© ISO 1997

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from the publisher.

International Organization for Standardization  
Case postale 56 • CH-1211 Genève 20 • Switzerland  
Internet central@iso.ch  
X.400 c=ch; a=400net; p=iso; o=isocs; s=central

Printed in Switzerland

## Foreword

ISO (the International Organization for Standardization) is a world-wide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

International Standard ISO 6391 was prepared by Technical Committee ISO/TC 34, *Agricultural food products*, Subcommittee SC 6, *Meat and meat products*.

This second edition cancels and replaces the first edition (ISO 6391:1988), which has been technically revised.

Annex A of this International Standard is for information only.

## Introduction

Although, for well-accepted statistical reasons, the limit for the lowest number of colonies counted per plate of selective medium is set at 15, for practical purposes it is often desirable to establish an estimated count of lower numbers of *Escherichia coli*. The confidence limits of such determinations are to be found in ISO 7218.

## iTeh STANDARD PREVIEW (standards.iteh.ai)

[ISO 6391:1997](#)

<https://standards.iteh.ai/catalog/standards/sist/73a6f77b-4d7d-4bf8-b772-4973f8178074/iso-6391-1997>

# Meat and meat products — Enumeration of *Escherichia coli* — Colony-count technique at 44 °C using membranes

## 1 Scope

This International Standard specifies a method for the enumeration of viable *Escherichia coli* present in all kinds of meat and meat products, including poultry.

The method will detect *Escherichia coli* (biotype 1) and lactose non-fermenting or anaerogenic variants (see reference [1]).

NOTE 1 This method may not be appropriate for the examination of processed meat having a high content of aerobic spore-forming bacteria.

NOTE 2 Some pathogenic strains of *Escherichia coli* may not be detected by this method.

[ISO 6391:1997](https://standards.iteh.ai/catalog/standards/sist/73a6f77b-4d7d-4bf8-b772-4973f8178074/iso-6391-1997)

## 2 Normative references

<https://standards.iteh.ai/catalog/standards/sist/73a6f77b-4d7d-4bf8-b772-4973f8178074/iso-6391-1997>

The following standards contain provisions which, through reference in this text, constitute provisions of this International Standard. At the time of the publication, the editions indicated were valid. All standards are subject to revision, and parties to agreements based on this International Standard are encouraged to investigate the possibility of applying the most recent editions of the standards indicated below. Members of IEC and ISO maintain registers of currently valid International Standards.

ISO 6887:1983, *Microbiology — General guidance for the preparation of dilutions for microbiological examination*

ISO 7218:1996, *Microbiology of food and animal feeding stuffs — General rules for microbiological examinations*

## 3 Definitions

For the purposes of this International Standard, the following definition applies.

### 3.1

#### ***Escherichia coli***

Bacteria which at 44 °C form indole-positive (pink) colonies on cellulose acetate membranes overlaid on tryptone bile agar, under the conditions specified in this International Standard.

## 4 Principle

In general, the detection of *Escherichia coli* requires three successive stages.

### 4.1 Resuscitation

Inoculation of a specified quantity of the initial suspension onto cellulose acetate membranes overlaid on mineral-modified glutamate agar, and incubation at 37 °C for 4 h.

NOTE — This procedure enables *Escherichia coli* damaged by storage under frozen, dried or chill conditions or damaged by heat or chemical processes to be resuscitated. It also permits the diffusion of high concentrations of any fermentable carbohydrate, if present in the test sample, which would otherwise interfere with indole production during the subsequent isolation stage.

### 4.2 Isolation

Transfer of membranes from the resuscitation stage on mineral-modified glutamate agar to tryptone bile agar. Then incubation at 44 °C for 18 h to 20 h.

### 4.3 Detection

Demonstration of the presence of *Escherichia coli* on the membranes by the production of indole by each colony.

### 4.4 Calculation

Calculation of the number of *Escherichia coli* per millilitre or per gram of sample from the number of indole-positive colonies obtained on membranes at dilution levels chosen to give a significant result.

**iTeh STANDARD PREVIEW**  
(standards.iteh.ai)  
<https://standards.iteh.ai/catalog/standards/sist/73a6f77b-4d7d-4bf8-b772-4973f8178074/iso-6391-1997>

## 5 Diluent, culture media and reagents

### 5.1 General

For current laboratory practice, see ISO 7218.

### 5.2 Diluent

See ISO 6887 for the preparation of dilutions.

### 5.3 Resuscitation media: Mineral-modified glutamate agar

#### 5.3.1 Composition

Sodium glutamate	6,35 g
Lactose	10,0 g
Sodium formate	0,25 g
L(-)-cystine	0,02 g
L(-)-aspartic acid	0,024 g
L(+)-arginine	0,02 g
Thiamine	0,001 g
Nicotinic acid	0,001 g
Pantothenic acid	0,001 g
Magnesium sulfate (MgSO <sub>4</sub> .7H <sub>2</sub> O)	0,100 g
Ammonium iron(II) citrate [at least 15 % Fe (m/m)]	0,010 g
Calcium chloride (CaCl <sub>2</sub> .2H <sub>2</sub> O)	0,010 g
Dipotassium hydrogen phosphate	0,90 g
Ammonium chloride	2,5 g
Agar	12 g to 18 g <sup>1)</sup>
Water	1 000 ml

1) Depending on the gel strength of the agar.

#### 5.3.2 Preparation

Dissolve the ammonium chloride in the water. Add the other components and heat to boiling. Adjust the pH so that after sterilization it is 6,7 at 25 °C. Transfer 100 ml volumes of the medium to suitable containers (6.5) and sterilize for 10 min in the autoclave (6.1) set at 121 °C.

<https://standards.iteh.ai/catalog/standards/sist/73a6f77b-4d7d-4bf8-b772-4973f8178074/iso-6391-1997>

#### 5.3.3 Preparation of agar plates

Pour 12 ml to 15 ml of the medium, cooled to 47 °C in the water bath (6.10), into sterile Petri dishes (6.6) and allow to solidify. The plates may be stored at between 0 °C and +5 °C for up to one week.

Immediately before use, dry the agar plates carefully (e.g. in the cabinet or oven (6.4) set at 50 °C), preferably with the lids off and agar surface downwards, until any droplets have disappeared from the surface of the medium. Do not dry them any further.

NOTE — The agar should be dry enough so that 1 ml of inoculum is totally absorbed into the membrane/agar within 15 min (see 9.2.3).

## 5.4 Selective medium: Tryptone bile agar (see reference [3])

### 5.4.1 Composition

Tryptone (peptic digest of casein)	20,0 g <sup>1)</sup>
Bile salts (refined)	1,5 g <sup>2)</sup>
Agar	12 g to 18 g <sup>3)</sup>
Water	1 000 ml

1) Commercial brands which favour indole formation should be used.

2) Oxoid Bile salts No. 3 is an example of a suitable product available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this product.

3) Depending on the gel strength of the agar.

### 5.4.2 Preparation

Dissolve the components in the water and heat to boiling. Adjust the pH so that after sterilization it is 7,2 at 25 °C. Transfer 100 ml volumes of the medium to suitable containers, and sterilize for 15 min in the autoclave (6.1) set at 121 °C.

### 5.4.3 Preparation of agar plates

Pour 12 ml to 15 ml of the medium, cooled to 47 °C in the water bath (6.10), into sterile Petri dishes (6.6) and allow to solidify. These plates may be stored at between 0 °C and +5 °C for up to 4 days.

Immediately before use, dry the agar plates carefully (e.g. in the cabinet or oven (6.4) set at 50 °C), preferably with the lids off and agar surface downwards, until any droplets have disappeared from the surface of the medium. Do not dry them any further.

## 5.5 Indole detection reagent (Vracko and Sherris reagent)

### 5.5.1 Composition

<i>p</i> -Dimethylaminobenzaldehyde	5 g
Hydrochloric acid, <i>c</i> (HCl) = 1 mol/l	100 ml

### 5.5.2 Preparation

Dissolve the reagent in the hydrochloric acid and store at room temperature for not longer than one month.

## 6 Apparatus and glassware

Usual microbiological laboratory apparatus and, in particular, the following.

### 6.1 Apparatus for dry sterilization (oven) or wet sterilization (autoclave)

See ISO 7218.



- 6.2 Incubator**, capable of being maintained at  $35\text{ °C} \pm 1\text{ °C}$  or  $37\text{ °C} \pm 1\text{ °C}$ .
- 6.3 Incubator**, capable of being maintained at  $44\text{ °C} \pm 0,5\text{ °C}$ .
- 6.4 Drying cabinet** or **oven**, ventilated by convection, capable of being maintained at  $50\text{ °C} \pm 1\text{ °C}$ .
- 6.5 Test tubes**, 18 mm x 180 mm, and **flasks** or **bottles**, 125 ml to 300 ml, for sterilization and storage of culture media.
- 6.6 Petri dishes**, made of glass or plastic, of 90 mm to 100 mm diameter.
- 6.7 Cellulose acetate membranes**, of 0,45  $\mu\text{m}$  to 1,2  $\mu\text{m}$  pore size and of 85 mm diameter.
- 6.8 Pipettes**, calibrated for bacteriological use, of 1 ml nominal capacity, graduated in divisions of 0,1 ml, and with an outflow opening of 2 mm to 3 mm in diameter.
- 6.9 Spreaders**, made of plastic or glass, for example hockey sticks made from a glass rod of approximately 3,5 mm diameter, and 20 cm length, bent at right angles about 3 cm from one end and with the cut ends made smooth by heating.
- 6.10 Water bath**, or similar apparatus, capable of being maintained at  $47\text{ °C} \pm 0,2\text{ °C}$ .
- 6.11 Longwave (365 nm) ultraviolet lamp**, fitted with a suitable filter to remove UV radiation below 310 nm.
- 6.12 pH-meter**, accurate to  $\pm 0,1$  pH units at  $25\text{ °C}$ .

ISO 6391:1997

- 7 Sampling** <https://standards.iteh.ai/catalog/standards/sist/73a6f77b-4d7d-4bf8-b772-4973f8178074/iso-6391-1997>

Sampling is not part of the method specified in this International Standard. A recommended sampling method is given in ISO 3100-1 [4].

It is important that the laboratory receive a sample which is truly representative and has not been damaged or changed during transport or storage.

## 8 Preparation of test sample

Prepare the sample in accordance with the specified International Standard appropriate to the product concerned. If there is no specific International Standard, it is recommended that the parties concerned come to an agreement on this subject.

A recommended method is given in ISO 3100-2 [5].

## 9 Procedure

### 9.1 Test portion, initial suspension and dilutions

See ISO 6887 and the specific International Standard appropriate to the product concerned.