



SLOVENSKI STANDARD
SIST-TS CEN/TS 15921:2010
01-februar-2010

Gnojila - Ekstrakcija topnega fosforja po Petermannu pri 65 °C

Fertilizers - Extraction of soluble phosphorus according to Petermann at 65 °C

Düngemittel - Extraktion des löslichen Phosphors nach Petermann bei 65 °C

Engrais - Extraction du phosphore soluble selon Petermann à 65 °C

Ta slovenski standard je istoveten z: CEN/TS 15921:2009

[SIST-TS CEN/TS 15921:2010](https://standards.iteh.ai/catalog/standards/sist/b3711bd6-67db-4c0d-9bbc-915773b5d5a5/sist-ts-cen-ts-15921-2010)

<https://standards.iteh.ai/catalog/standards/sist/b3711bd6-67db-4c0d-9bbc-915773b5d5a5/sist-ts-cen-ts-15921-2010>

ICS:

65.080 Gnojila Fertilizers

SIST-TS CEN/TS 15921:2010 **en,fr,de**

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[SIST-TS CEN/TS 15921:2010](#)

<https://standards.iteh.ai/catalog/standards/sist/3711bd6-67db-4c0d-9bbc-915773b5d5a5/sist-ts-cen-ts-15921-2010>

TECHNICAL SPECIFICATION
SPÉCIFICATION TECHNIQUE
TECHNISCHE SPEZIFIKATION

CEN/TS 15921

September 2009

ICS 65.080

English Version

**Fertilizers - Extraction of soluble phosphorus according to
Petermann at 65 °C**

Engrais - Extraction du phosphore soluble selon Petermann
à 65 °C

Düngemittel - Extraktion des löslichen Phosphors nach
Petermann bei 65 °C

This Technical Specification (CEN/TS) was approved by CEN on 3 August 2009 for provisional application.

The period of validity of this CEN/TS is limited initially to three years. After two years the members of CEN will be requested to submit their comments, particularly on the question whether the CEN/TS can be converted into a European Standard.

CEN members are required to announce the existence of this CEN/TS in the same way as for an EN and to make the CEN/TS available promptly at national level in an appropriate form. It is permissible to keep conflicting national standards in force (in parallel to the CEN/TS) until the final decision about the possible conversion of the CEN/TS into an EN is reached.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.

[SIST-TS CEN/TS 15921:2010](https://standards.iteh.ai/catalog/standards/sist/b3711bd6-67db-4c0d-9bbc-915773b5d5a5/sist-ts-cen-ts-15921-2010)

<https://standards.iteh.ai/catalog/standards/sist/b3711bd6-67db-4c0d-9bbc-915773b5d5a5/sist-ts-cen-ts-15921-2010>



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: Avenue Marnix 17, B-1000 Brussels

Contents		Page
Foreword.....		3
1	Scope	4
2	Normative references	4
3	Terms and definitions	4
4	Principle.....	4
5	Sampling.....	4
6	Reagents.....	5
7	Apparatus	6
8	Procedure	6
Bibliography.....		7

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[SIST-TS CEN/TS 15921:2010](https://standards.iteh.ai/catalog/standards/sist/3711bd6-67db-4c0d-9bbc-915773b5d5a5/sist-ts-cen-ts-15921-2010)
<https://standards.iteh.ai/catalog/standards/sist/3711bd6-67db-4c0d-9bbc-915773b5d5a5/sist-ts-cen-ts-15921-2010>

Foreword

This document (CEN/TS 15921:2009) has been prepared by Technical Committee CEN/TC 260 "Fertilizers and liming materials", the secretariat of which is held by DIN.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

This document has been prepared under a mandate given to CEN by the European Commission and the European Free Trade Association.

According to the CEN/CENELEC Internal Regulations, the national standards organizations of the following countries are bound to announce this Technical Specification: Austria, Belgium, Bulgaria, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and the United Kingdom.

iTeh STANDARD PREVIEW (standards.iteh.ai)

[SIST-TS CEN/TS 15921:2010](https://standards.iteh.ai/catalog/standards/sist/3711bd6-67db-4c0d-9bbc-915773b5d5a5/sist-ts-cen-ts-15921-2010)

<https://standards.iteh.ai/catalog/standards/sist/3711bd6-67db-4c0d-9bbc-915773b5d5a5/sist-ts-cen-ts-15921-2010>

CEN/TS 15921:2009 (E)**1 Scope**

This document specifies the procedure for the determination of soluble phosphorus in alkaline ammonium citrate.

The method is applicable exclusively to precipitated dehydrated dicalcium phosphate ($\text{CaHPO}_4 \cdot 2\text{H}_2\text{O}$).

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN 1482-2, *Fertilizers and liming materials — Sampling and sample preparation — Part 2: Sample preparation*

EN 12944-1:1999, *Fertilizers and liming materials and soil improvers — Vocabulary — Part 1: General terms*

EN 12944-2:1999, *Fertilizers and liming materials and soil improvers — Vocabulary — Part 2: Terms relating to fertilizers*

EN 15475, *Fertilizers — Determination of ammoniacal nitrogen*

ITeH STANDARD PREVIEW
(standards.iteh.ai)

3 Terms and definitions

[SIST-TS CEN/TS 15921:2010](https://standards.iteh.ai/catalog/standards/sist/3711bd6-67db-4c0d-9bbc-915775b5d5a5/sist-ts-cen-ts-15921-2010)

For the purposes of this document, the terms and definitions given in EN 12944-1:1999 and EN 12944-2:1999 apply.

4 Principle

Extraction of phosphorus from the test portion at a temperature of 65 °C with an alkaline solution of ammonium citrate (Petermann's solution) under the specified conditions.

5 Sampling

Sampling is not part of the method specified in this document. A recommended sampling method is given in EN 1482-1.

Sample preparation shall be carried out in accordance with EN 1482-2. Grinding of the laboratory sample is recommended for homogeneity reasons.

6 Reagents

6.1 Water, distilled or demineralized having the same characteristics as distilled water.

6.2 Petermann's solution

6.2.1 Characteristics of the Petermann's solution

- citric acid ($C_6H_8O_7 \cdot H_2O$): 173 g/l;
- ammonia: 42 g/l of ammoniacal nitrogen;
- pH between 9,4 and 9,7.

6.2.2 Preparation from diammonium citrate

Dissolve 931 g of diammonium citrate (molecular mass 226,19) in about 3 500 ml of water (6.1), in a 5 l standard flask. Stand in a bath of running water, mix and cool and add in small amounts ammonia. For example, for $d_{20} = 906$ g/ml corresponding to a level of 20,81 % by mass of ammoniacal nitrogen, it is necessary to use 502 ml of ammonia solution. Adjust the temperature to 20 °C, make up to volume with water (6.1) and mix.

6.2.3 Preparation from citric acid and ammonia

Dissolve 865 g of citric acid monohydrate in about 2 500 ml of water (6.1) in a container of about 5 l capacity. Place the container in an ice bath, and add in small amounts, shaking constantly, ammonia solution using a funnel, the stem of which is immersed in the citric acid solution. For example, for $d_{20} = 906$ g/ml corresponding to a level of 20,81 % by mass of ammoniacal nitrogen, it is necessary to add 1 114 ml of ammonia solution. Adjust the temperature to 20 °C, transfer to a 5 l standard flask, make up to the mark with water (6.1) and mix.

6.2.4 Checking of the ammoniacal nitrogen content

Transfer 25 ml of the solution into a 250 ml standard flask and make up to volume with water (6.1) and mix. Determine the ammoniacal content on 25 ml of this solution according to EN 15475. If the solution is correct, an amount of 15 ml of $c = 0,5$ mol/l H_2SO_4 shall be used.

If the strength of ammoniacal nitrogen is greater than 42 g/l, NH_3 can be expelled by a stream of inert gas or by moderate heating to bring back the pH to 9,7. Carry out a second determination.

If the strength of ammoniacal nitrogen is less than 42 g/l, it will be necessary to add a mass M in grams of ammonia solution:

$$M = (42 - n \times 2,8) \times \frac{500}{20,81} \quad (1)$$

or a volume, V , at 20 °C:

$$V = \frac{M}{0,906} \quad (2)$$

If V is less than 25 ml, add it directly to the 5 l flask with a mass of $V \times 0,173$ g powdered citric acid.

If V is greater than 25 ml, it will be convenient to prepare a new litre of reagent in the following way.

CEN/TS 15921:2009 (E)

Weigh 173 g of citric acid. Dissolve it in 500 ml of water. And, taking the precautions indicated, add not more than $225 + V \times 1.206$ ml of ammonia solution which was used to prepare the 5 l of reagent. Make up to volume with water and mix.

Mix this volume of 1 l with the 4 975 ml previously prepared.

7 Apparatus

- 7.1 **Common laboratory equipment and glassware**, in particular equipment according to 7.2 to 7.4.
- 7.2 **Water bath**, which can be maintained at a temperature of (65 ± 1) °C.
- 7.3 **500 ml graduated flask**, e.g. Stohmann.
- 7.4 **Dry fluted filter paper**, free from phosphates.

8 Procedure**8.1 Test portion**

Weigh, to the nearest 0,001 g, 1 g of the prepared laboratory sample and transfer it to the graduated flask (7.3).

iTeh STANDARD PREVIEW
(standards.iteh.ai)

8.2 Extraction

Add 200 ml of alkaline ammonium citrate solution (6.2). Stopper the flask and shake vigorously by hand to avoid the formation of lumps and to prevent any adherence of the substance to the sides.

Place the flask in the water bath set at 65 °C (7.2) and shake every 5 min during the first 0,5 h. After each shaking, raise the stopper to equilibrate the pressure. The level of water in the water bath ought to be above the level of solution in the flask. Allow the flask to remain in the water bath a further 1 h at 65 °C and shake every 10 min. Remove the flask, cool to a temperature of about 20 °C, make up to a volume of 500 ml with water. Mix and filter through a dry fluted filter paper (7.4), rejecting the first portion of filtrate. Continue the filtering until a sufficient quantity of filtrate is obtained to carry out the phosphorus determination.