

SLOVENSKI STANDARD SIST EN 12932:2008

01-november-2008

Nadomešča:

SIST EN 12932:2000

Kemikalije, ki se uporabljajo za pripravo pitne vode - Kemikalije za uporabo v sili - Natrijev dikloroizociaourat, dehidriran

Chemicals used for treatment of water intended for human consumption - Chemicals for emergency use - Sodium dichloroisocyanurate, dihydrate

iTeh STANDARD PREVIEW

Produkte zur Aufbereitung von Wasser für den menschlichen Gebrauch - Produkte für den Notfall - Natriumdichlorisocyanurat-dihydrat

SIST EN 12932:2008

Produits chimiques pour le traitement de l'eau destinée à la consommation humaine -Produits chimiques utilisés en cas d'urgence - Dichloroisocyanurate de sodium, dihydraté

Ta slovenski standard je istoveten z: EN 12932:2008

ICS:

13.060.20 Pitna voda Drinking water

71.100.80 Kemikalije za čiščenje vode Chemicals for purification of

water

SIST EN 12932:2008 en,fr,de

SIST EN 12932:2008

iTeh STANDARD PREVIEW (standards.iteh.ai)

SIST EN 12932:2008

https://standards.iteh.ai/catalog/standards/sist/38afb2e1-1396-41a3-8bc2-07b28fb41e2e/sist-en-12932-2008

EUROPEAN STANDARD

EN 12932

NORME EUROPÉENNE

EUROPÄISCHE NORM

July 2008

ICS 71.100.80

Supersedes EN 12932:2000

English Version

Chemicals used for treatment of water intended for human consumption - Chemicals for emergency use - Sodium dichloroisocyanurate, dihydrate

Produits chimiques pour le traitement de l'eau destinée à la consommation humaine - Produits chimiques utilisés en cas d'urgence - Dichloroisocyanurate de sodium, dihydraté

Produkte zur Aufbereitung von Wasser für den menschlichen Gebrauch - Produkte für den Notfall -Natriumdichlorisocyanurat-dihydrat

This European Standard was approved by CEN on 28 June 2008.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latyia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.

07b28fb41e2e/sist-en-12932-2008



EUROPEAN COMMITTEE FOR STANDARDIZATION COMITÉ EUROPÉEN DE NORMALISATION EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: rue de Stassart, 36 B-1050 Brussels

Contents

		Page
Fore	eword	3
Intro	oduction	5
1	Scope	6
2	Normative references	6
3	Description	6
4	Purity criteria	8
5	Test methods	9
6	Labelling - Transportation - Storage	16
Ann	ex A (informative) General information on sodium dichloroisocyanurate, dihydrate	19
Ann	ex B (normative) General rules relating to safety	21
Ann	ex C (normative) Determination of arsenic, antimony and selenium (atomic absorption spectrometry hydride technique)	23
	iography(Standards.iteh.ai)	

SIST EN 12932:2008

https://standards.iteh.ai/catalog/standards/sist/38afb2e1-1396-41a3-8bc2-07b28fb41e2e/sist-en-12932-2008

Foreword

This document (EN 12932:2008) has been prepared by Technical Committee CEN/TC 164 "Water supply", the secretariat of which is held by AFNOR.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by January 2009, and conflicting national standards shall be withdrawn at the latest by January 2009.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

This document supersedes EN 12932:2000.

Significant technical difference between this edition and EN 12932:2000 is as follows:

— Deletion of reference to EU Directive 80/778/EEC of July 15, 1980 in order to take into account the latest Directive in force (see [1]).

Other amendments:

- iTeh STANDARD PREVIEW
- a) section 3.1.2 addition of "Troclosene sodium, dihydrate" as a synonym; (standards, iteh.ai)
- b) section 4.3 inclusion of the maximum admissible water content {14%};
- c) section 5.2: SIST EN 12932:2008 https://standards.iteh.ai/catalog/standards/sist/38afb2e1-1396-41a3-8bc2-
 - 1) inclusion of section 5.2.2.1 Water;
 - 2) change numbering of following section to 5.2.2.2 Sodium chloride;
- d) section 6.1 inclusion of big bags as permitted means of delivery;
- e) section 6.2 updating of risk and safety labelling according to EU directives (Annex I of the Directive 67/548EEC and its latest Adaptations to Technical Progress);
 - 1) correction of hazard symbols {"N: Dangerous for the environment" instead of "O: Oxidizing"}
 - 2) addition of R phrase:
 - R 50/53: Very toxic to aquatic organisms, may cause long-term adverse effects in the aquatic environment.
 - 3) addition of S phrases:
 - S2: Keep out of the reach of children.
 - S 60: This material and its container must be disposed of as hazardous waste.
 - S 61: Avoid release to the environment. Refer to special instructions/safety data sheets.
- f) section 6.3:

- 1) precision of the transport classification {"is not subject to the UN Model Regulations on the Transport of Dangerous goods" instead of "is not listed under a UN Number"};
- 2) change to note N° 5) {"United Nations Model Regulations on the Transport of dangerous goods, chapter 3.3.1, Special provision N° 135" instead of "United Nations Number"};
- g) section 6.4 addition of "Troclosene sodium, dihydrate" as one possible chemical name.

According to the CEN/CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and the United Kingdom.

iTeh STANDARD PREVIEW (standards.iteh.ai)

<u>SIST EN 12932:2008</u> https://standards.iteh.ai/catalog/standards/sist/38afb2e1-1396-41a3-8bc2-07b28fb41e2e/sist-en-12932-2008

Introduction

In respect of potential adverse effects on the quality of water intended for human consumption, caused by the product covered by this European Standard:

- a) this European Standard provides no information as to whether the product may be used without restriction in any of the Member States of the EU or EFTA;
- b) it should be noted that, while awaiting the adoption of verifiable European criteria, existing national regulations concerning the use and/or the characteristics of this product remain in force.

NOTE Conformity with this European Standard does not confer or imply acceptance or approval of the product in any of the Member States of the EU or EFTA. The use of the product covered by this European Standard is subject to regulation or control by National Authorities.

iTeh STANDARD PREVIEW (standards.iteh.ai)

<u>SIST EN 12932:2008</u> https://standards.iteh.ai/catalog/standards/sist/38afb2e1-1396-41a3-8bc2-07b28fb41e2e/sist-en-12932-2008

1 Scope

This European Standard is applicable to sodium dichloroisocyanurate dihydrate used for emergency treatment of water intended for human consumption. It describes the characteristics of sodium dichloroisocyanurate dihydrate and specifies the requirements and the corresponding test methods for sodium dichloroisocyanurate dihydrate. It gives information on its use in water treatment. It also determines the rules relating to safe handling and use of sodium dichloroisocyanurate dihydrate (see Annex B).

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN 1233, Water quality - Determination of chromium - Atomic absorption spectrometric methods

EN 1483, Water quality - Determination of mercury - Method using atomic absorption spectrometry

EN ISO 3696, Water for analytical laboratory use - Specification and test methods (ISO 3696:1987)

ISO 3165, Sampling of chemical products for industrial use - Safety in sampling

ISO 6206, Chemical products for industrial use. Sampling - Vocabulary

ISO 8213, Chemical products for industrial use. Sampling techniques - Solid chemical products in the form of particles varying from powders to coarse lumps

ISO 8288:1986, Water quality - Determination of cobalt, nickel, copper, zinc, cadmium and lead - Flame atomic absorption spectrometric methods iteh alcatalog/standards/sist/38afb2e1-1396-41a3-8bc2-07b28fb41e2e/sist-en-12932-2008

3 Description

3.1 Identification

3.1.1 Chemical name

1-sodium, 3, 5-dichloro-1,3,5-triazine-2, 4, 6-trione dihydrate.

3.1.2 Synonym or common name

Sodium dichloroisocyanurate, dihydrate.

Troclosene sodium, dihydrate

3.1.3 Relative molecular mass

256,02

3.1.4 Empirical formula

C₃N₃O₃Cl₂Na.2H₂O

3.1.5 Chemical formula

$$Na$$
 $| Na$
 $|$

3.1.6 CAS Registry Number 1)

51580-86-0

3.1.7 EINECS reference 2)

Not listed.

3.2 Commercial formeh STANDARD PREVIEW

The product is available in various granular forms ds.iteh.ai)

3.3 Physical properties

SIST EN 12932:2008

https://standards.iteh.ai/catalog/standards/sist/38afb2e1-1396-41a3-8bc2-

3.3.1 Appearance and odour 07b28fb41e2e/sist-en-12932-2008

The product is a white granular solid with chlorinous odour.

3.3.2 Density

The bulk density of the product is approximately 0,9 g/cm³.

3.3.3 Solubility in water

The solubility of the product in water is 291 g/l at 25 °C.

3.3.4 Vapour pressure

Not applicable.

3.3.5 Boiling point at 100 kPa 3)

Not applicable, the product decomposes before fusion.

- 1) Chemical Abstracts Service Registry Number.
- 2) European Inventory of Existing Commercial Chemical Substances.
- 3) 100 kPa = 1 bar

3.3.6 Melting point

Not applicable.

3.3.7 Specific heat

 $(1,090 \pm 0,040)$ kJ/kg.K at 20 °C.

3.3.8 Viscosity

Not applicable.

3.3.9 Critical temperature

Not applicable.

3.3.10 Critical pressure

Not applicable.

3.3.11 Physical hardness

Not applicable.

iTeh STANDARD PREVIEW

3.4 Chemical properties

(standards.iteh.ai)

The product is a strong oxidant. It liberates chlorine by hydrolysis.

SIST EN 12932:2008

4 Purity criteria

https://standards.iteh.ai/catalog/standards/sist/38afb2e1-1396-41a3-8bc2-07b28fb41e2e/sist-en-12932-2008

4.1 General

This European Standard specifies the minimum purity requirements for dihydrate sodium dichloroisocyanurate used for the treatment of water intended for human consumption. Limits are given for impurities commonly present in the product. Depending on the raw material and the manufacturing process other impurities may be present and, if so, this shall be notified to the user and when necessary to the relevant authorities.

NOTE Users of this product should check the national regulations in order to clarify whether it is of appropriate purity for treatment of water intended for human consumption, taking into account raw water quality, required dosage, contents of other impurities and additives used in the product not stated in this product standard.

Limits have been given for impurities and chemical parameters where these are likely to be present in significant quantities from the current production process and raw materials. If the production process or raw materials lead to significant quantities of impurities, by-products or additives being present, this shall be notified to the user.

4.2 Composition of commercial product

The product shall contain at least a mass fraction of 55 % of available chlorine as calculated in accordance with the corresponding method given in 5.2.1.

4.3 Impurities and main by-products

The water content shall be less than a mass fraction of 14 % of the product.

The sodium chloride content shall be less than a mass fraction of 0,05 % of the product.

4.4 Chemical parameters

The product shall conform to the requirements specified in Table 1.

Table 1 - Chemical parameters

Element		Limit mg/kg of product	
		Type 1	Type 2
Arsenic (As)	max.	10	10
Cadmium (Cd)	max.	1	1
Chromium (Cr)	max.	6	10
Mercury (Hg)	max.	0,02	0,02
Nickel (Ni)	max.	3	3
Lead (Pb)	max.	4	4
Antimony (Sb)	max.	5	5
Selenium (Se)	max.	1	1

NOTE. Cyanide, which does not exist in a strong oxidizing medium such as sodium dichloroisocyanurate, dihydrate, is not a relevant chemical parameter. Pesticides and polycyclic aromatic hydrocarbons are not by-products of the manufacturing process. For parametric values of sodium dichloroisocyanurate on trace metal content in drinking water, see [1].

SIST EN 12932:2008

5 Test methods*tps://standards.iteh.ai/catalog/standards/sist/38afb2e1-1396-41a3-8bc2-07b28fb41e2e/sist-en-12932-2008

5.1 Sampling

Observe the general recommendations of ISO 3165 and take account of ISO 6206. Prepare the laboratory sample(s) required by the relevant procedure described in ISO 8213.

5.2 Analysis

5.2.1 Determination of available chlorine (main product)

5.2.1.1 Principle

The available chlorine is determined by measuring active chlorine in the sample. The oxidizing chlorine reacts with potassium iodide releasing iodine which is then titrated with sodium thiosulfate standard volumetric solution in the presence of starch indicator solution.

5.2.1.2 Reagents

All reagents shall be of a recognized analytical grade and the water used shall conform to grade 3 in accordance with EN ISO 3696.

5.2.1.2.1 Potassium iodide crystals (KI).

5.2.1.2.2 Glacial acetic acid.

5.2.1.2.3 Sodium thiosulfate standard volumetric solution, $c(Na_2S_2O_3.5H_2O) = 0.1 \text{ mol/l.}$

Standard volumetric solutions of sodium thiosulfate are commercially available.

Alternatively a standard volumetric solution can be prepared by the following procedure:

Dissolve 24,8 g Na₂S₂O₃ . 5 H₂O in a 1 000 ml one-mark volumetric flask in about 0,75 l water. After the temperature has equalized make up to the mark with water and mix thoroughly.

To standardize:

Weigh, to the nearest 0,1 mg, 3,600 g (m)of dry potassium iodate. Dissolve in water in a 1 000 ml one-mark volumetric flask, make up to the mark with water and mix (standard reference solution $c(1/6 \text{ KIO}_3) = 0,1 \text{ mol/l})$. Place 200 ml of water in a 500 ml stoppered conical flask, add ($2 \pm 0,5$) g of potassium iodide and stir to dissolve. Then introduce,by means of a pipette, 10,0 ml of sodium thiosulfate solution for standardization, add (15 ± 1) ml of hydrochloric acid solution (diluted 1 + 1 by volume) and (5 ± 1) ml of starch solution (5.2.1.2.4). Titrate immediately with the potassium iodate standard reference solution until the appearance of a blue coloration persisting for at least 30 s occurs . Record the volume (V_1) of iodate used.

The actual concentration, c, of the sodium thiosulfate standard volumetric solution (Na₂S₂O_{3.5}H₂O), expressed in moles per litre is given by the following equation:

$$c = \frac{V_1 \times c_1}{V} \tag{1}$$

where

- is the concentration, expressed in moles per litre, of the potassium jodate standard reference solution [c(1/6 KIO₃) = 0,1 mol/l]; (standards.iteh.ai)
- V is the volume, in millilitres, of the sodium thiosulfate standard volumetric solution used for the standardization (V= 10 ml); SIST EN 12932:2008
- https://standards.iteh.ai/catalog/standards/sist/38afb2e1-1396-41a3-8bc2- V_1 is the volume, in millilitres, of potassium iodate standard reference solution used in the titration.

5.2.1.2.4 Starch solution, mass fraction 1 %.

Make a slurry with $(1 \pm 0,1)$ g starch and (5 ± 1) ml water. Add (90 ± 5) ml boiling water to the slurry. Stir to dissolve it and cool the solution. This solution needs refrigeration to avoid the decomposition of the starch which results in a vague end point. Keep the solution cool and use it within one week.

NOTE Commercial indicators for iodine titration exist and can be used in place of the specified starch solution provided that their efficiency has been previously tested.

5.2.1.3 Apparatus

5.2.1.3.1 Ordinary laboratory apparatus and glassware.

5.2.1.4 Procedure

5.2.1.4.1 Test portion

Weigh, to the nearest 0,1 mg, 0,25 g of the laboratory sample (m_0) into a tared stoppered weighing bottle.

5.2.1.4.2 Determination

Transfer the test portion to a 200 ml volumetric flask.