

---

---

**Animal feeding stuffs — Enzymatic  
determination of total starch content**

*Aliments des animaux — Détermination enzymatique de la teneur totale  
en amidon*

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

[ISO 15914:2004](https://standards.iteh.ai/catalog/standards/sist/cd5c8cc9-ac42-4da1-ae81-667c8b249b4e/iso-15914-2004)

[https://standards.iteh.ai/catalog/standards/sist/cd5c8cc9-ac42-4da1-ae81-  
667c8b249b4e/iso-15914-2004](https://standards.iteh.ai/catalog/standards/sist/cd5c8cc9-ac42-4da1-ae81-667c8b249b4e/iso-15914-2004)



**PDF disclaimer**

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

ISO 15914:2004

<https://standards.iteh.ai/catalog/standards/sist/cd5c8cc9-ac42-4da1-ae81-667c8b249b4e/iso-15914-2004>

© ISO 2004

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Case postale 56 • CH-1211 Geneva 20  
Tel. + 41 22 749 01 11  
Fax + 41 22 749 09 47  
E-mail [copyright@iso.org](mailto:copyright@iso.org)  
Web [www.iso.org](http://www.iso.org)

Published in Switzerland

## Contents

Page

Foreword.....	iv
1 Scope.....	1
2 Normative references .....	1
3 Terms and definitions .....	1
4 Principle .....	2
5 Reagents .....	2
6 Apparatus.....	4
7 Preparation of test sample .....	5
8 Procedure.....	5
9 Calculation and expression of results .....	7
10 Precision .....	8
11 Test report.....	9
Annex A (informative) Notes on procedure .....	10
Annex B (informative) Results of interlaboratory test .....	11
Bibliography .....	12

<https://standards.iteh.ai/catalog/standards/sist/cd5c8cc9-ac42-4da1-ac81-667c8b249b4e/iso-15914-2004>  
 ISO 15914:2004

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 15914 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 10, *Animal feeding stuffs*.

## iTeh STANDARD PREVIEW (standards.iteh.ai)

[ISO 15914:2004](https://standards.iteh.ai/catalog/standards/sist/cd5c8cc9-ac42-4da1-ae81-667c8b249b4e/iso-15914-2004)

<https://standards.iteh.ai/catalog/standards/sist/cd5c8cc9-ac42-4da1-ae81-667c8b249b4e/iso-15914-2004>

# Animal feeding stuffs — Enzymatic determination of total starch content

## 1 Scope

This international Standard specifies a method for the enzymatic determination of the total starch content of animal feeding stuffs and raw materials for animal feeding stuffs.

The method is also applicable to the determination of the purity of starch.

It is important that in the sample matrix no components are present which contribute to the measured extinction at 340 nm.

The analytical range of the method is 40 g/kg to 1 000 g/kg starch. The standard procedure is applicable to the range 200 g/kg to 1000 g/kg. For the lower range, 40 g/kg to 200 g/kg, another dilution procedure for the standard glucose solution and samples can be used.

iTeh STANDARD PREVIEW

## 2 Normative references (standards.iteh.ai)

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 3696:1987, *Water for analytical laboratory use — Specification and test methods*

ISO 6498:1998, *Animal feeding stuffs — Preparation of test samples*

## 3 Terms and definitions

For the purpose of this document, the following terms and definitions apply.

### 3.1

#### starch

natural vegetable polymer consisting of long linear unbranched chains of 1,4- $\alpha$ -D-glucose units (amylose) and/or long  $\alpha$ -1,6-branched chains of  $\alpha$ -1,4-linked glucose units (amylopectin)

### 3.2

#### starch content

mass fraction of starch and its high molecular mass breakdown products, insoluble in 40 % ethanol, and determined in accordance with the method of this International Standard

NOTE The starch content is expressed in grams per kilogram.

## 4 Principle

The milled test sample is extracted with 40 % ethanol to remove the soluble sugars. Sample disintegration and starch solubilization is achieved by dispersing the extracted test portion in aqueous DMSO (90 % volume fraction) at 100 °C, followed by addition of concentrated hydrochloric acid at 60 °C. The solubilized and dissolved starch is quantitatively converted into glucose by the enzyme amyloglucosidase. The glucose quantification is carried out by the well-known hexokinase method (see [1], [2]).

## 5 Reagents

Use only reagents of recognized analytical grade.

**5.1 Water**, complying with at least grade 3 in accordance with ISO 3696:1987.

**5.2 Ethanol** (C<sub>2</sub>H<sub>5</sub>OH), 40 % (volume fraction)

Take 417 ml of ethanol (96 % volume fraction) and dilute with water to 1 000 ml

**5.3 Hydrochloric acid**,  $c(\text{HCl}) = 12 \text{ mol/l}$ .

**5.4 Aqueous sodium hydroxide**,  $c(\text{NaOH}) = 4 \text{ mol/l}$ .

Dissolve in a beaker 40 g of NaOH in about 50 ml water. After cooling, transfer quantitatively to a 250 ml volumetric flask and dilute to the mark with water.

**WARNING** — Heat develops. Wear safety glasses.

**5.5 Acetic acid solution**,  $c(\text{CH}_3\text{COOH}) = 2 \text{ mol/l}$ .

Add to a 500 ml volumetric flask about 200 ml water, followed by 59 ml of glacial acetic acid. Dilute to the mark with water.

**5.6 Sodium acetate solution**,  $c(\text{CH}_3\text{COONa}) = 2 \text{ mol/l}$ .

Dissolve, in a 500 ml volumetric flask, 82,0 g of sodium acetate in water and dilute to the mark with water.

**5.7 Sodium acetate buffer**,  $c(\text{CH}_3\text{COONa}/\text{H}) = 2 \text{ mol/l}$ , pH = 4,8.

Mix 41 ml of acetic acid solution (5.5) with 59 ml of sodium acetate solution (5.6). Check the pH with a pH-meter and, if necessary, adjust to obtain the correct pH with the acetic acid or the sodium acetate solution.

Prepare a fresh buffer solution daily.

**5.8 Aqueous dimethylsulfoxide**, ( $\sigma_{\text{DMSO}}$ ) = 90 % (volume fraction).

Mix pure DMSO and water in a volume ratio of 9:1.

**5.9 Clarifying solutions**, according to Carrez, prepared as follows.

**5.9.1 Potassium hexacyanoferrate(II) solution**,  $c[\text{K}_4\text{Fe}(\text{CN})_6] = 0,25 \text{ mol/l}$ .

In a 1 litre volumetric flask, dissolve 106 g of potassium hexacyanoferrate(II) trihydrate  $[\text{K}_4\text{Fe}(\text{CN})_6 \cdot 3\text{H}_2\text{O}]$  in water. Dilute to the mark with water.

**5.9.2 Zinc acetate**, solution in 0,5 mol/l acetic acid,  $c[\text{Zn}(\text{CH}_3\text{CO}_2)_2] = 1 \text{ mol/l}$ .

In a 1 litre volumetric flask, dissolve 219,5 g of zinc acetate dihydrate  $[\text{Zn}(\text{CH}_3\text{CO}_2)_2 \cdot 2\text{H}_2\text{O}]$  and 30 g of glacial acetic acid in water. Dilute to the mark with water.

### 5.10 Iodine solution in potassium iodide

In a 1 litre volumetric flask, dissolve 12,7 g of iodine ( $\text{I}_2$ ) and 24,0 g of potassium iodide (KI) in water. Dilute to the mark with water. Dilute this solution 10-fold before use.

### 5.11 Standard glucose solution

#### 5.11.1 Samples containing 200 g/kg to 1 000 g/kg starch

Prepare three independent standard glucose solutions ( $c = 0,0194 \text{ mol/l}$ ). In each 100 ml volumetric flask dissolve 350 mg anhydrous glucose ( $\text{C}_6\text{H}_{12}\text{O}_6$ ) to the nearest mg in water. Dilute to the mark with water.

#### 5.11.2 Samples containing 40 g/kg to 200 g/kg starch

Prepare three independent standard glucose solutions ( $c = 0,0039 \text{ mol/l}$ ). In each 500 ml volumetric flask, dissolve  $350 \text{ mg} \pm 1 \text{ mg}$  of anhydrous glucose ( $\text{C}_6\text{H}_{12}\text{O}_6$ ) in water. Dilute to the mark with water.

Prepare fresh standard glucose solutions daily.

### 5.12 Amyloglucosidase solution, 160 U/ml AMG.

Dissolve, in a mixture of 9 ml of water and 1 ml of sodium acetate buffer (5.7), 267 mg of amyloglucosidase (AMG) [EC 3.2.1.3 (*Aspergillus niger*, Roche Diagnostics, No. 1 202 367, 6 U/mg)].<sup>1)</sup> With respect to storage and the use of the enzymes, follow carefully the recommendations of the enzyme supplier.

When another enzyme is used, the activity should be estimated as given in Note 1. The mass of the enzyme should be adapted to the activity found.

NOTE 1 Different enzyme suppliers use different definitions for the units of activities of enzymes. In this International Standard the following definition for the unit of activity for AMG has been used: 1 unit of amyloglucosidase will release 1  $\mu\text{mol}$  of glucose from glycogen in 1 min at 25 °C and pH = 4,75.

NOTE 2 10 ml of enzyme solutions is enough for about 75 determinations.

**5.13 D-Glucose UV test set**, for quantifying glucose enzymatically, with the hexokinase method (R-Biopharm, No. 716251<sup>2)</sup>, according to the manufacturer, the unused kits may be stored for 1 year at 4 °C), as given in 5.13.1 to 5.13.3.

#### 5.13.1 Buffer/substrate solution (Bottle 1)

Dissolve the content of Bottle 1 in 45 ml of freshly prepared distilled water. Store this reagent in a cool (4 °C) and dark place for not longer than for 4 weeks. Use the required amount of this solution at ambient temperature.

#### 5.13.2 Enzyme solution (Bottle 2)

This solution is ready for use.

1) Roche Diagnostics No. 1 202 367 is an example of a suitable product available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this product.

2) R-Biopharm AG No. 716251 is an example of a suitable product available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this products.

### 5.13.3 Colouring solution

Mix 22,5 ml of buffer/substrate solution (5.13.1) with 95 ml of water and 0,45 ml of enzyme solution (5.13.2) and homogenize. This amount of reagent is enough for a series of about 40 measurements. If more or less measurements are to be carried out, the mixing volumes may be adapted.

Only use fresh prepared colouring solutions at ambient temperature.

## 6 Apparatus

Usual laboratory apparatus and, in particular, the following.

- 6.1 Analytical balance**, capable of weighing to the nearest 0,1 mg.
- 6.2 Centrifuge**, with an acceleration of  $(180 \pm 10) g$  and of  $3\,000 g$ , suitable for screw cap tubes of 12 ml (the value of  $g$  is given for the bottom of the tube).
- 6.3 Water bath**, adjusted to  $(100 \pm 2) ^\circ\text{C}$ .
- 6.4 Water bath**, adjusted to  $(60 \pm 1) ^\circ\text{C}$ .
- 6.5 pH-meter**, calibrated, with a combined glass electrode, capable of measuring the pH to the nearest 0,01 pH unit.
- 6.6 Microlitre pipettes**, calibrated and adjustable from 40  $\mu\text{l}$  to 200  $\mu\text{l}$ , 200  $\mu\text{l}$  to 1 000  $\mu\text{l}$ , and 1 ml to 5 ml, and a **dispenser**, adjustable from 1 ml to 5 ml.
- 6.7 Spectrometer**, with a flow cuvette, capable of measuring at 340 nm.
- 6.8 Rotary shaker**, speed 50 r/min, for screw-cap-closed centrifuge tubes (6.11).
- 6.9 Dispenser/diluter** (e.g. Hook & Tucker Compudil D<sup>3</sup>).
- 6.10 Heating oven**, with forced air circulation.
- 6.11 Centrifuge tubes**, 100 mm  $\times$  16 mm, made of glass, with screw cap fitted with PTFE covered rubber seals.
- 6.12 Measuring flasks**, of capacity 100 ml, wide-neck, equipped with a ground glass joint 14/23.
- It shall be possible to measure the pH in the flask with the combined glass electrode.
- 6.13 Tube mixer**, (e.g. Vortex mixer<sup>3</sup>), for the alternative method for the disintegration and solubilization of the starch, as described in 8.4.2. The following apparatus is also needed.
- 6.14 Shaking water bath**, adjusted to  $(100 \pm 2) ^\circ\text{C}$ , with a stroke of 2 cm and a shaking frequency of 150 to 200 strokes per minute. The water bath should be equipped with a holder in which the tubes fit horizontally in the water.
- 6.15 Centrifuge tubes**, made of glass, of capacity at least 20 ml, with a screw cap fitted with PTFE covered rubber seal.
- 6.16 Glass pearls**, 3 mm in diameter.

---

3) Hook and Tucker Compudil D and Vortex are examples of suitable products available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of these products.



## 7 Preparation of test sample

Prepare the test sample in accordance with ISO 6498.

## 8 Procedure

### 8.1 Test portion

Carry out the determination in duplicate.

Weigh approximately 200 mg of the test sample (Clause 7), to the nearest 0,1 mg, in a centrifuge tube (6.11). In a second centrifuge tube, weigh, to the nearest 0,1 mg, another 200 mg as the duplicate of the pretreated sample (Clause 7).

Repeat the procedure for each tube.

### 8.2 Extraction of the sugars

Add to each tube 10 ml of ethanol (5.2). Extract the solubilized sugars by rotating the suspension continuously for 10 min in the rotary shaker (6.8). Then centrifuge for 10 min in the centrifuge (6.2) with a radial acceleration of  $(180 \pm 10) g$  and discard the supernatant. Repeat this extraction procedure once again.

### 8.3 Sample blank iTeh STANDARD PREVIEW

From this stage in the procedure, in each analysis series a sample blank shall be analysed. Use an empty tube as a sample blank.

### 8.4 Disintegration of the starch ISO 15914:2004 <https://standards.iteh.ai/catalog/standards/sist/cd5c8cc9-ac42-4da1-ae81-667c8b249b4e/iso-15914-2004>

#### 8.4.1 General

The disintegration/solubilization of starch may be carried out in two ways, i.e. according to 8.4.2 or according to 8.4.3. When the alternative procedure (see 8.4.3) is used, the larger glass reagent tubes with screw caps (6.15) shall be used.

#### 8.4.2 Procedure I

With the dispenser under continuous vortex mixing on a vortex mixer (6.13), add 10,0 ml of aqueous DMSO solution (5.8) to the tube and keep mixing until a lump-free suspension is obtained. Close the tube with the screw cap.

**NOTE** Vigorous homogenization during the addition of DMSO is necessary to prevent the formation of microgel and/or lumps. Microgel and lumps will result in incorrect starch contents

- due to microgel and lumps producing an inhomogeneous sample solution, and
- the microgel and lumps will not completely be hydrolysed by AMG into glucose, so an incorrect (mostly too low) starch content will be determined.

Immediately after homogenization, attach the tube to the rotary shaker (6.8) and disintegrate the starch under continuous rotation for 30 min in an oven (6.10) set at 100 °C. Cool the tube and add, with an adjustable pipette (6.6), 1,7 ml of concentrated hydrochloric acid (5.3) and mix well. Close the tube and place it in the rotary shaker. Partly hydrolyse the sample by placing the tube under continuous rotation for 30 min in an oven (6.10) set at 60 °C.

Proceed as in 8.4.4.