
**Microbiology of food and animal feeding
stuffs — Horizontal method for the
detection of *Escherichia coli* O157**

*Microbiologie des aliments — Méthode horizontale pour la recherche des
Escherichia coli O157*

iTeh Standards
(<https://standards.iteh.ai>)
Document Preview

ISO 16654:2001

<https://standards.iteh.ai/catalog/standards/iso/f8b4d94-6a9f-4800-9da6-2d347cb40119/iso-16654-2001>



Reference number
ISO 16654:2001(E)

PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

iTeh Standards
(<https://standards.iteh.ai>)
Document Preview

ISO 16654:2001

<https://standards.iteh.ai/catalog/standards/iso/f8b4d94-6a9f-4800-9da6-2d347cb40119/iso-16654-2001>

© ISO 2001

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.ch
Web www.iso.ch

Printed in Switzerland

Contents

Page

Foreword.....	iv
Introduction	v
1 Scope	1
2 Normative references	1
3 Term and definition	1
4 Principle.....	2
5 Culture media, reagents and antisera	2
6 Apparatus and glassware	7
7 Sampling.....	8
8 Preparation of test sample.....	8
9 Procedure (see annex A).....	8
9.1 Test portion and initial suspension	8
9.2 Enrichment	8
9.3 Immunomagnetic separation (IMS).....	8
9.4 Plating out onto selective agars and identification of <i>E. coli</i> O157 colonies.....	9
9.5 Confirmation.....	10
9.6 Further characterization.....	11
10 Quality assurance.....	11
10.1 Test strains for quality assurance purposes	11
10.2 Culture method	11
11 Expression of results	11
12 Test report	11
Annex A (normative) Diagram of procedure	12
Bibliography	13

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 3.

Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this International Standard may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

International Standard ISO 16654 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 9, *Microbiology*.

Annex A forms a normative part of this International Standard.

ITeH Standards
(<https://standards.iteh.ai>)
Document Preview

ISO 16654:2001

<https://standards.iteh.ai/catalog/standards/iso/f8b4d94-6a9f-4800-9da6-2d347cb40119/iso-16654-2001>

Introduction

Because of the large variety of food and feed products, this horizontal method may not be appropriate in every detail for certain products. In this case, different methods specific to these products may be used if absolutely necessary for justified technical reasons. Nevertheless, every attempt should be made to apply this horizontal method as far as possible.

When this International Standard is next reviewed, account will be taken of all information then available regarding the extent to which this horizontal method has been followed and the reasons for deviations from this method in the case of particular products.

The harmonization of test methods cannot be immediate, and for certain groups of products International Standards and/or national standards may already exist that do not comply with this horizontal method. It is hoped that when such standards are reviewed they will be changed to comply with this International Standard so that eventually the only remaining departures from this horizontal method will be those necessary for well-established technical reasons.

iTeh Standards
(<https://standards.itih.ai>)
Document Preview

[ISO 16654:2001](https://standards.itih.ai/catalog/standards/iso/f8b4d94-6a9f-4800-9da6-2d347cb40119/iso-16654-2001)

<https://standards.itih.ai/catalog/standards/iso/f8b4d94-6a9f-4800-9da6-2d347cb40119/iso-16654-2001>

Microbiology of food and animal feeding stuffs — Horizontal method for the detection of *Escherichia coli* O157

WARNING — *Escherichia coli* O157 can cause severe life-threatening illness and has a low infective dose. Laboratory-acquired infections have been reported.

In order to safeguard the health of laboratory personnel, it is essential that the whole of this method be carried out only by skilled personnel using good laboratory practices and preferably working in a containment facility. Relevant national Health and Safety Regulations relating to this organism must be adhered to.

Care must be taken in the disposal of all infectious materials.

1 Scope

This International Standard specifies a horizontal method for the detection of *Escherichia coli* serogroup O157.

Subject to the limitations discussed in the introduction, this International Standard is applicable to products intended for human consumption or for animal feeding stuffs.

2 Normative references

The following normative documents contain provisions which, through reference in this text, constitute provisions of this International Standard. For dated references, subsequent amendments to, or revisions of, any of these publications do not apply. However, parties to agreements based on this International Standard are encouraged to investigate the possibility of applying the most recent editions of the normative documents indicated below. For undated references, the latest edition of the normative document referred to applies. Members of ISO and IEC maintain registers of currently valid International Standards.

ISO 6887-1, *Microbiology of food and animal feeding stuffs — Preparation of test samples, initial suspension and decimal dilutions for microbiological examination — Part 1: General rules for the preparation of the initial suspension and decimal dilutions*.

ISO 7218, *Microbiology of food and animal feeding stuffs — General rules for microbiological examinations*.

3 Term and definition

For the purposes of this International Standard, the following term and definition applies.

3.1

Escherichia coli O157

E. coli O157

microorganisms which form typical colonies on the surface of the plating-out medium used in this International Standard, and which produce indole and agglutinate specifically with antiserum against the O157 antigen

NOTE 1 Sorbitol-positive *E. coli* O157 strains are not detected on CT-SMAC (5.2) media.

NOTE 2 Some indole-negative mutations have been found.

4 Principle

The detection of *Escherichia coli* O157 necessitates four successive stages (see annex A).

- Enrichment** of the test portion homogenized in modified tryptone soya broth containing novobiocin (mTSB + N) with incubation at $41,5\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$ for 6 h and subsequently for a further 12 h to 18h.
- Separation and concentration** of microorganisms by means of immunomagnetic particles coated with antibodies to *E. coli* O157.
- Isolation** by subculture of the immunomagnetic particles with adhering bacteria onto cefixime tellurite sorbitol MacConkey agar (CT-SMAC) and the user's choice of a second selective isolation agar.
- Confirmation** of sorbitol-negative colonies from CT-SMAC and colonies typical of *E. coli* O157 on the second isolation agar, by indole production and agglutination with *E. coli* O157 antiserum.

NOTE Further characterization, by for example pathogenic markers, of the positive isolates can be obtained by forwarding them to an appropriate reference laboratory.

5 Culture media, reagents and antisera

For current laboratory practices, see ISO 7218.

5.1 Enrichment medium: Modified tryptone soya broth with novobiocin (mTSB + N)

See reference [1].

5.1.1 Modified tryptone soya broth (mTSB)

5.1.1.1 Composition

Enzymatic digest of casein	17,0 g
Enzymatic digest of soya	3,0 g
D(+)-glucose	2,5 g
Bile salts No. 3	1,5 g
Sodium chloride	5,0 g
Dipotassium hydrogen phosphate (K_2HPO_4)	4,0 g
Water	1 000 ml

5.1.1.2 Preparation

Dissolve the components or the dehydrated complete medium in the water, by heating if necessary. Adjust the pH, using the pH-meter (6.6), if necessary, so that after sterilization it is $7,4 \pm 0,2$ at $25\text{ }^{\circ}\text{C}$.

Dispense the medium in appropriate amounts in flasks or bottles (6.7).

Sterilize for 15 min in the autoclave (6.1) set at $121\text{ }^{\circ}\text{C}$.

5.1.2 Novobiocin solution

5.1.2.1 Composition

Novobiocin	0,45 g
Water	100 ml

5.1.2.2 Preparation

Dissolve the novobiocin in the water and sterilize by membrane filtration.

Prepare on the day of use.

5.1.2.3 Preparation of the complete medium

Immediately before use, add 1 ml or 4 ml of novobiocin solution (5.1.2) to either 225 ml or 900 ml of cooled mTSB (5.1.1).

The final concentration of novobiocin is 20 mg per litre of mTSB.

5.2 First selective isolation medium: Cefixime tellurite sorbitol MacConkey agar (CT-SMAC)

See reference [2].

5.2.1 Base medium

5.2.1.1 Composition

Enzymatic digest of casein	17,0 g
Enzymatic digest of animal tissues	3,0 g
Sorbitol	10,0 g
Bile salts No. 3	1,5 g
Sodium chloride	5,0 g
Neutral Red	0,03 g
Crystal Violet	0,001 g
Agar	9 g to 18 g ^a
Water	1 000 ml

^a Depending on the gel strength of the agar.

5.2.1.2 Preparation

Dissolve the basic components or the complete dehydrated base in the water by boiling. Adjust the pH (6.6), if necessary, so that after sterilization it is $7,1 \pm 0,2$ at 25 °C.

Sterilize for 15 min in the autoclave (6.1) set at 121 °C.

5.2.2 Potassium tellurite solution

5.2.2.1 Composition

Potassium tellurite for bacteriological use	0,25 g
Water	100 ml

5.2.2.2 Preparation

Dissolve the potassium tellurite in the water and sterilize by membrane filtration.

This solution may be stored at room temperature for up to 1 month, but discard it if a white precipitate forms.

5.2.3 Cefixime solution

5.2.3.1 Composition

Cefixime	5,0 mg
Water	100,0 ml

5.2.3.2 Preparation

Dissolve the cefixime in the water and sterilize by membrane filtration.

NOTE Cefixime may need to be dissolved in ethanol.

This solution may be stored at $3\text{ }^{\circ}\text{C} \pm 2\text{ }^{\circ}\text{C}$ for 1 week.

5.2.4 Complete medium

5.2.4.1 Composition

Base medium (5.2.1)	1 000 ml
Potassium tellurite solution (5.2.2)	1,0 ml
Cefixime solution (5.2.3)	1,0 ml

5.2.4.2 Preparation

Either cool the freshly sterilized base medium (5.2.1) to between $44\text{ }^{\circ}\text{C}$ and $47\text{ }^{\circ}\text{C}$ (6.5), or melt it by steaming the previously sterilized and solidified base medium, then cool to between $44\text{ }^{\circ}\text{C}$ and $47\text{ }^{\circ}\text{C}$.

Add 1 ml of the tellurite solution and 1 ml of the cefixime solution to 1000 ml of the base medium. Mix and pour about 15 ml amounts into sterile Petri dishes (6.15). Allow to solidify.

The final concentration of tellurite is 2,5 mg/l and cefixime 0,05 mg/l.

Immediately before use, dry the agar plates, preferably with the lids removed and with the agar surfaces facing downwards, in an oven set at a temperature between $25\text{ }^{\circ}\text{C}$ and $50\text{ }^{\circ}\text{C}$ (6.2), until the droplets have disappeared from the surface of the medium. Do not dry them any further. The agar plates may also be dried in a laminar-flow safety cabinet for 30 min with half-open lids, or overnight with the lids in place.

If prepared in advance, the undried plates may be stored in the dark in plastic bags or other moisture-retentive containers, in a refrigerator at $3\text{ }^{\circ}\text{C} \pm 2\text{ }^{\circ}\text{C}$ for up to 2 weeks.

5.3 Second selective isolation medium

Use any other solid selective medium, at the choice of the laboratory, complementary to CT-SMAC agar and especially appropriate for the isolation of *Escherichia coli* O157.

Immediately before use, dry the agar plates, preferably with the lids removed and with the agar surfaces facing downwards, in an oven set at a temperature between $25\text{ }^{\circ}\text{C}$ and $50\text{ }^{\circ}\text{C}$ (6.2), until the droplets have disappeared