

iTeh STANDARD PREVIEW
(standards.iteh.ai)

SIST EN 15477:2009

<https://standards.iteh.ai/catalog/standards/sist/d51c8d24-70db-4f04-b9bf-d34824f5d3a6/sist-en-15477-2009>

EUROPEAN STANDARD
NORME EUROPÉENNE
EUROPÄISCHE NORM

EN 15477

January 2009

ICS 65.080

Supersedes CEN/TS 15477:2006

English Version

Fertilizers - Determination of the water-soluble potassium content

Engrais - Détermination de la teneur en potassium soluble dans l'eau

Düngemittel - Bestimmung von wasserlöslichem Kalium

This European Standard was approved by CEN on 30 November 2008.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and United Kingdom.

[SIST EN 15477:2009](https://standards.iteh.ai/catalog/standards/sist/d51c8d24-70db-4f04-b9bf-d34824f5d3a6/sist-en-15477-2009)

<https://standards.iteh.ai/catalog/standards/sist/d51c8d24-70db-4f04-b9bf-d34824f5d3a6/sist-en-15477-2009>



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: rue de Stassart, 36 B-1050 Brussels

Contents

Page

Foreword.....	3
1 Scope	4
2 Normative references	4
3 Terms and definitions	4
4 Principle.....	4
5 Reagents.....	4
6 Apparatus	5
7 Sampling and sample preparation	5
8 Procedure	5
9 Calculation and expression of the result	7
10 Precision.....	8
11 Test report	9
Annex A (informative) Results of the inter-laboratory tests	10
Bibliography	11

iTech STANDARD PREVIEW
(standards.itech.ai)

SIST EN 15477:2009
<https://standards.itech.ai/catalog/standards/sist/d51c8d24-70db-4f04-b9bf-d34824f5d3a6/sist-en-15477-2009>

Foreword

This document (EN 15477:2009) has been prepared by Technical Committee CEN/TC 260 “Fertilizers and liming materials”, the secretariat of which is held by DIN.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by July 2009, and conflicting national standards shall be withdrawn at the latest by July 2009.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

This document supersedes CEN/TS 15477:2006.

This document has been prepared under a mandate given to CEN by the European Commission and the European Free Trade Association.

According to the CEN/CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland and the United Kingdom.

(standards.iteh.ai)

[SIST EN 15477:2009](https://standards.iteh.ai/catalog/standards/sist/d51c8d24-70db-4f04-b9bf-d34824f5d3a6/sist-en-15477-2009)

<https://standards.iteh.ai/catalog/standards/sist/d51c8d24-70db-4f04-b9bf-d34824f5d3a6/sist-en-15477-2009>

EN 15477:2009 (E)**1 Scope**

This European Standard specifies a method for the determination of water-soluble potassium, which is applicable to all potassium fertilizers listed in Annex I of the Regulation (EC) No 2003/2003 [3].

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN 1482-2, *Fertilizers and liming materials — Sampling and sample preparation — Part 2: Sample preparation*

EN 12944-1:1999, *Fertilizers and liming materials and soil improvers — Vocabulary — Part 1: General terms*

EN 12944-2:1999, *Fertilizers and liming materials and soil improvers — Vocabulary — Part 2: Terms relating to fertilizers*

EN ISO 3696:1995, *Water for analytical laboratory use — Specification and test methods (ISO 3696:1987)*

3 Terms and definitions

ITeh STANDARD PREVIEW

(standards.iteh.ai)

For the purposes of this document, the terms and definitions given in EN 12944-1:1999 and EN 12944-2:1999 apply.

[SIST EN 15477:2009](https://standards.iteh.ai/catalog/standards/sist/d51c8d24-70db-4f04-b9bf-d34824f5d3a6/sist-en-15477-2009)

<https://standards.iteh.ai/catalog/standards/sist/d51c8d24-70db-4f04-b9bf-d34824f5d3a6/sist-en-15477-2009>

4 Principle

The potassium in the sample to be analyzed is dissolved in water. After eliminating or fixing the substances that might interfere with the quantitative determination, the potassium is precipitated in a slightly alkaline medium in the form of potassium tetraphenylborate.

5 Reagents**5.1 General**

Use only reagents of recognized analytical grade and distilled or demineralized water (grade 3 according to EN ISO 3696:1995).

5.2 Formaldehyde, clear formaldehyde solution with a mass fraction of 25 % to 35 % formaldehyde.

5.3 Potassium chloride, p. a.

5.4 Sodium hydroxide solution, $c = 10 \text{ mol/l}$.

Care should be taken to ensure that only potassium free sodium hydroxide is used.

5.5 Indicator solution

Dissolve 0,5 g of phenolphthalein in ethanol at 90 % and make the volume up to 100 ml.

5.6 EDTA solution

Dissolve 4 g of the dihydrated disodium salt of ethylenediaminetetraacetic acid in water in a 100 ml graduated flask. Make up the volume and mix.

Store the reagent in a plastics container.

5.7 STPB solution

Dissolve 32,5 g of sodium tetrphenylborate in 480 ml of water, add 2 ml of the sodium hydroxide solution (5.4) and 20 ml of a magnesium chloride solution (100 g of $MgCl_2 \cdot 6H_2O$ per litre).

Stir for 15 min and filter through a fine, ashless filter.

Store this reagent in a plastics container.

5.8 Liquid for washing

Dilute 20 ml of the STPB solution (5.7) to 1 000 ml with water.

5.9 Bromine water, saturated bromine solution in water.

6 Apparatus

6.1 Graduated flasks, capacity 1 000 ml.

6.2 Beaker, capacity 250 ml and 600 ml.

6.3 Filter crucibles, porosity 5 μm to 20 μm .

6.4 Oven, regulated at (120 ± 10) °C.

6.5 Desiccator

7 Sampling and sample preparation

Sampling is not part of the method specified in this document. A recommended sampling method is given in EN 1482-1.

Sample preparation shall be carried out in accordance with EN 1482-2. Grinding is recommended for homogeneity reasons.

8 Procedure

8.1 Test portion

Weigh to the nearest 0,001 g 10 g of the prepared sample (5 g for potassium salts with a mass fraction of potassium oxide of more than 50 %). Place this test portion in a 600 ml beaker with approximately 400 ml of water.

Bring to a boil and allow it to boil for 30 min. Cool, transfer quantitatively into a 1 000 ml graduated flask, make up the volume, mix and filter into a dry receiver. Discard the first 50 ml of the filtrate (see 8.6).

EN 15477:2009 (E)**8.2 Preparation of the aliquot part for precipitation**

Transfer by pipette an aliquot part of the filtrate containing 25 mg to 50 mg of potassium (see Table 1) and place it in a 250 ml beaker. If required make up to 50 ml with water.

To remove any interference, add 10 ml of the EDTA solution (5.6), several drops of the phenolphthalein solution (5.5) and stir in, drop by drop, sodium hydroxide solution (5.4) until it turns red, then finally add a few more drops of sodium hydroxide to ensure an excess (usually 1 ml of sodium hydroxide is sufficient to neutralize the sample and ensure an excess).

To eliminate most of the ammonia, (see 8.6) boil gently for 15 min.

If necessary, add water to make the volume up to 60 ml.

Bring the solution to the boil, remove the beaker from the heat and add 10 ml of formaldehyde (5.2). Add several drops of phenolphthalein and, if necessary, some more sodium hydroxide until a distinct red colour appears. Cover the beaker with a watch glass and place it on a steam bath for 15 min.

8.3 Weighing the crucible

Dry the filter crucible to a constant mass (about 15 min) in the oven at 120 °C.

Allow the crucible to cool in a desiccator and then weigh it.

8.4 Precipitation

Remove the beaker from the steam bath, stir in drop-by-drop 10 ml of the STPB solution (5.7). This addition takes about 2 min. Wait for at least 10 min before filtering.

8.5 Filtering and washing

Filter under vacuum into the weighed crucible, rinse the beaker with the liquid for washing (5.8), wash the precipitate three times with the liquid for washing (60 ml in all of the liquid for washing), and twice with 5 ml to 10 ml of water.

Dry the precipitate thoroughly.

8.6 Drying and weighing

Wipe the outside of the crucible with a filter paper. Place the crucible with its contents in the oven for 1,5 h at 120 °C. Allow the crucible to cool in a desiccator to ambient temperature and weigh immediately.

If the filtrate is dark in colour, transfer by pipette, an aliquot part containing at the most, 100 mg of K_2O and place in a 100 ml graduated flask. Add bromine water (5.9) and bring to a boil to eliminate any surplus bromine. After cooling make up the volume, filter and quantitatively determine the potassium in an aliquot part of the filtrate.

Where there is little or no ammoniacal nitrogen present there is no need to boil for 15 min.

iTeh STANDARD PREVIEW

(standards.iteh.ai)

SIST EN 15477:2009

<https://standards.iteh.ai/catalog/standards/sist/d51c8d24-70db-4f04-b9bf->[d34824f5d3a6/sist-en-15477-2009](https://standards.iteh.ai/catalog/standards/sist/d51c8d24-70db-4f04-b9bf-d34824f5d3a6/sist-en-15477-2009)

8.7 Aliquot parts to be taken as samples and conversion factors

Table 1 — Aliquot parts and conversion factors

K ₂ O in the fertilizer %	K in the fertilizer %	Sample for analysis g	Sample of the extract solution for the dilution ml	Dilution to ml	Aliquot part to be taken as a sample for precipitation ml	Conversion factor <i>F</i> $\frac{\% \text{ K}_2\text{O}}{\text{g TPBK}}$	Conversion factor <i>F'</i> $\frac{\% \text{ K}}{\text{g TPBK}}$
5 - 10	4,2 – 8,3	10	-	-	50	26,280	21,812
10 - 20	8,3 – 16,6	10	-	-	25	52,560	43,624
20 - 50	16,6 – 41,5	10{	either –	250	10	131,400	109,060
			or 50		50	131,400	109,060
more than 50	more than 41,5	5{	either –	250	10	262,800	218,120
			or 50		50	262,800	218,120

8.8 Blank test

iTeh STANDARD PREVIEW
(standards.itech.ai)

For each series of determinations, carry out a blank test using only the reagents in the proportions used in the analysis and allow for this when calculating the final result.

[SIST EN 15477:2009](#)

8.9 Control test <https://standards.itech.ai/catalog/standards/sist/d51c8d24-70db-4f04-b9bf-d34824f5d3a6/sist-en-15477-2009>

In order to obtain a control for the Method of analysis, carry out a determination on an aliquot part of an aqueous solution of potassium chloride, containing at the most 40 mg of K₂O.

9 Calculation and expression of the result

9.1 Dilution according to Table 1

Calculate the K₂O content, $w_{\text{K}_2\text{O}}$, as mass fraction in percent of the fertilizer according to equation (1):

$$w_{\text{K}_2\text{O}} = (m_1 - m_2) \times F \quad (1)$$

Calculate the K content, w_{K} , as mass fraction in percent of the fertilizer according to equation (2):

$$w_{\text{K}} = (m_1 - m_2) \times F' \quad (2)$$

where

m_1 is the mass of the precipitate from the sample, in grams;

m_2 is the mass of the precipitate from the blank, in grams;

F and F' conversion factors (see Table 1).