
**Tobacco and tobacco products —
Determination of organochlorine pesticide
residues — Gas chromatographic method**

*Tabac et produits du tabac — Dosage des résidus de pesticides
organochlorés — Méthode par chromatographie en phase gazeuse*

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[ISO 4389:2000](https://standards.iteh.ai/catalog/standards/sist/51978da7-acac-4d36-b3d4-f065cc5b285/iso-4389-2000)

[https://standards.iteh.ai/catalog/standards/sist/51978da7-acac-4d36-b3d4-
f065cc5b285/iso-4389-2000](https://standards.iteh.ai/catalog/standards/sist/51978da7-acac-4d36-b3d4-f065cc5b285/iso-4389-2000)



PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 4389:2000

<https://standards.iteh.ai/catalog/standards/sist/51978da7-acac-4d36-b3d4-f065cc5b285/iso-4389-2000>

© ISO 2000

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.ch
Web www.iso.ch

Printed in Switzerland

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 3.

Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this International Standard may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

International Standard ISO 4389 was prepared by Technical Committee ISO/TC 126, *Tobacco and tobacco products*.

This third edition cancels and replaces the second edition (ISO 4389:1997). As a result of extensive examination, the scope of this edition has been extended to tobacco products. Technical Corrigendum 1:1998 has been included.

Annexes A and B of this International Standard are for information only.

<https://standards.iteh.ai/catalog/standards/sist/51978da7-acac-4d36-b3d4-f065cc5b285/iso-4389-2000>

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 4389:2000

<https://standards.iteh.ai/catalog/standards/sist/51978da7-acac-4d36-b3d4-f065cc5b285/iso-4389-2000>

Tobacco and tobacco products — Determination of organochlorine pesticide residues — Gas chromatographic method

1 Scope

This International Standard specifies a method for the gas chromatographic determination of pesticide residues in leaf tobacco, manufactured tobacco and tobacco products.

The method is applicable to the determination of the organochlorine pesticides listed in Table 1. The method is particularly recommended for determination of the substances within the detection limits listed in Table 1.

NOTE 1 The method has been demonstrated to be free from errors that may arise from interfering peaks on the chromatogram that originate from non-organochlorine pesticide substances, when applied to various styles of leaf tobacco. However, the preceding statement cannot be assumed to apply to all forms and variants of tobacco and tobacco products and care should be taken in the interpretation of any unexpected, apparently positive results. Some analysts advocate the use of mass spectrometric confirmation of the chemical structure of compounds detected by chromatography in such circumstances.

NOTE 2 ISO 1750 contains the systematic chemical names and structures corresponding to the common names given in Table 1.

2 Normative references

<https://standards.iteh.ai/catalog/standards/sist/51978da7-acac-4d36-b3d4-f065cc5b2285/iso-4389-2000>

<https://standards.iteh.ai/catalog/standards/sist/51978da7-acac-4d36-b3d4-f065cc5b2285/iso-4389-2000>

The following normative documents contain provisions which, through reference in this text, constitute provisions of this International Standard. For dated references, subsequent amendments to, or revisions of, any of these publications do not apply. However, parties to agreements based on this International Standard are encouraged to investigate the possibility of applying the most recent editions of the normative documents indicated below. For undated references, the latest edition of the normative document referred to applies. Members of ISO and IEC maintain registers of currently valid International Standards.

ISO 648, *Laboratory glassware — One-mark pipettes.*

ISO 1042, *Laboratory glassware — One-mark volumetric flasks.*

ISO 3696, *Water for analytical laboratory use — Specification and test methods.*

ISO 4874, *Tobacco — Sampling of batches of raw material — General principles.*

ISO 8243, *Cigarettes — Sampling.*

3 Principle

The pesticide residues from a dried and milled sample, mixed with Florisil®¹⁾, are extracted by *n*-hexane in a special Soxhlet extractor. They are determined without any further clean-up with a gas chromatographic apparatus equipped with an electron-capture detector.

4 Reagents

All reagents shall be suitable for pesticide residue analysis. All solvents shall be checked for purity before use by carrying out a blank determination using exactly the same procedure (extraction and gas chromatography) as used for the test sample. The chromatogram obtained from the solvents shall have a baseline without noticeable peaks that could interfere with those from the pesticide residues being determined.

4.1 Water, degassed, in accordance with at least grade 2 of ISO 3696.

4.2 *n*-Hexane.

4.3 Toluene.

4.4 Reference substances: certified reference materials of minimum purity 95 % (mass fraction) of the substances listed in Table 1.

NOTE *trans*-Chlordane is used as an indicator for chlordane (technical mixture). If α -endosulfan is detected by this method, it is advisable to determine residues of the sum of α -endosulfan, β -endosulfan and endosulfan sulfate by a method suitable for such determinations.

4.5 Internal standard stock solution

Use Mirex,²⁾ an obsolete pesticide which has been superseded (see reference [2]).

NOTE Mirex is a generic name for dodecachloropentacyclo[5.2.1.0^{2,6}.0^{3,9}.0^{5,8}]decane.

Weigh, to the nearest 0,000 1 g, 0,02 g of Mirex into a 100 ml volumetric flask. Dilute to the mark with *n*-hexane (4.2).

4.6 Internal standard solution

Pipette 5 ml of the internal standard stock solution (4.5) into a 200 ml volumetric flask and dilute to the mark with *n*-hexane (4.2) to give a solution containing approximately 5 μ g/ml of Mirex. Store the internal standard solution at between 0 °C and +4 °C and exclude light. Under these conditions the internal standard solution is stable for at least 6 months.

1) Florisil® is an example of a suitable product available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this product.

2) Mirex is an example of a suitable product available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this product.

Table 1 — List of suitable pesticides with detection limits

Substance	Common name (ISO 1750)	Detection limit µg/g
Aldrin	aldrin	0,02
<i>trans</i> -Chlordane	chlordane	0,02
<i>p,p'</i> -DDE	—	0,02
<i>o,p'</i> -DDT	—	0,04
<i>p,p'</i> -DDT	DDT	0,06
Dieldrin	dieldrin	0,02
α -Endosulfan	endosulfan	0,03
HCB	hexachlorobenzene	0,02
α -HCH or α -BHC	HCH or BHC	0,02
β -HCH or β -BHC	HCH or BHC	0,02
γ -HCH or γ -BHC	gamma-HCH (lindane) or gamma-BHC	0,01
δ -HCH or δ -BHC	HCH or BHC	0,02
Heptachlor	heptachlor	0,02
Heptachlor epoxide	—	0,02
<i>o,p'</i> -TDE or <i>o,p'</i> -DDD	—	0,03
<i>p,p'</i> -TDE or <i>p,p'</i> -DDD	TDE	0,02
<i>o,p'</i> -DDE	—	0,03

4.7 Standard pesticide solutions

Store all pesticide solutions at between 0 °C and +4 °C and exclude light. Under these conditions the solutions are stable for at least 6 months.

4.7.1 Individual standard stock solutions

Weigh, to the nearest 0,000 1 g, 0,02 g of each pesticide reference substance (4.4) into individual 100 ml volumetric flasks. Dilute to the mark with *n*-hexane (4.2) to give individual standard stock solutions containing approximately 200 µg/ml of the individual pesticide.

In the case of β -HCH, the standard stock solution should be made by dissolving the pesticide in toluene (4.3) because of reduced solubility in *n*-hexane.

4.7.2 Mixed stock solution A

Pipette 5 ml of each individual standard stock solution (4.7.1) into a single 200 ml volumetric flask and dilute to the mark with *n*-hexane (4.2) [or toluene (4.3) for β -HCH] to give a solution containing approximately 5 $\mu\text{g/ml}$ of each pesticide.

4.7.3 Mixed stock solution B

Pipette 1 ml of mixed stock solution A (4.7.2) into a 10 ml volumetric flask and dilute to the mark with *n*-hexane (4.2) to give a solution containing approximately 0,5 $\mu\text{g/ml}$ of each pesticide.

4.7.4 Standard calibration solution

Pipette 1 ml of both the mixed stock solution A (4.7.2) and the internal standard solution (4.6) into a 100 ml volumetric flask and dilute to the mark with *n*-hexane (4.2) to give a solution containing approximately 0,05 $\mu\text{g/ml}$ of each pesticide and internal standard.

4.8 Florisil®, 150 μm to 250 μm .

NOTE Florisil® is a special, selected variety of magnesium silicate. The nominal aperture size of 150 μm to 250 μm corresponds to a mesh size range designated as 60 mesh to 100 mesh.

4.8.1 The quality of the Florisil® is one of the most critical features of the test method. The activity of the Florisil® must be sufficient to retain impurities present in the extract from the sample while allowing the pesticide residues to be eluted. The Florisil® shall first be pre-treated as described in 4.8.2. Only Florisil® that passes the subsequent verification test described in 4.8.3 shall be used.

4.8.2 Heat sufficient Florisil® for the verification test for at least 5 h in a quartz crucible in a muffle furnace at 550 °C. Allow the Florisil® to cool in a desiccator that contains no desiccant and transfer to a round-bottom flask. Add 5 ml of water (4.1) for every 100 g of Florisil®. Mix thoroughly in a rotating flask for approximately 1 h. Allow the Florisil® to equilibrate by storing in a tightly closed glass container for at least 48 h before proceeding as described in 4.8.3.

4.8.3 The activity level of the Florisil® is checked by the extraction of dieldrin from *n*-hexane solution. The solution shall have a concentration equivalent to that of an extract from tobacco containing 1,0 $\mu\text{g/g}$ of this pesticide. The activity level of the pretreated Florisil® is correct if the recovery of dieldrin is better than 95 %.

The activity of the Florisil® shall be checked each time a new portion is prepared.

5 Apparatus

It is essential to clean all glassware very thoroughly before use and to avoid the use of plastics containers and stopcock grease, otherwise impurities may be introduced into the solvents. All volumetric flasks and pipettes shall comply with class A of ISO 1042 and class A of ISO 648 respectively.

Usual laboratory apparatus and the following items.

5.1 Rotary evaporator.


5.2 Tobacco mill, with 2 mm mesh.

5.3 Oven, with ventilation.

5.4 Muffle furnace.

5.5 Heating mantles.

5.6 Soxhlet extractor, for continuous extraction (see Figure 1).



Key

- 1 Tobacco + Florisil®
- 2 Florisil®
- 3 Filter disk porosity 1

NOTE Distillation rate 200 ml/h.

Figure 1 — Apparatus used for tobacco extraction

5.7 Reflux condenser.

5.8 Desiccator.

5.9 Quartz crucible.

5.10 Gas chromatograph

Operate the gas chromatograph in accordance with the manufacturer's instructions. The injection port, oven and detector shall each be equipped with a separate heating unit.

The following conditions have been found to be satisfactory on a particular make of instrument and are given for guidance. If other conditions are used they should be validated prior to use.

a) Temperatures

The injection port temperature shall be between 180 °C and 210 °C. The detector temperature shall be between 290 °C and 340 °C. If any other conditions are used they shall be sufficient to achieve satisfactory separation of all components and similar to that given in the specimen chromatogram (see Figure A.1 for an example). A suitable temperature programme is

- initial temperature 40 °C
- initial time 2 min
- temperature profile 1 20 °C/min from 40 °C to 150 °C
- temperature profile 2 3 °C/min from 150 °C to 270 °C
- final time 15 min at 270 °C
- total GC run time 62,5 min

iTeh STANDARD PREVIEW
(standards.iteh.ai)

b) Gas flow rates

Gas flow rates should be set according to the instrument manufacturer's guidance and the analyst's experience. Suitable gas flow conditions are <https://standards.iteh.ai/catalog/standards/sist/51978da7-acac-4d36-b3d4-f065cc5b285/iso-4389-2000>

- carrier gas helium, 4 ml/min
- make-up gas nitrogen, 30 ml/min
- septum purge 5 ml/min
- split vent 30 ml/min

c) Injection mode

Use 2 µl splitless with split valve closed for 1 min after injection.

5.10.1 Injection device

Use an automated injector or any suitable alternative means of injection.

For manual injection, the use of a microsyringe capable of injecting 1 µl to 5 µl portions is recommended. Before solutions are injected with the syringe, rinse it at least ten times with *n*-hexane then five times with the solution. After injection, rinse the syringe five times with *n*-hexane.

5.10.2 Column

A fused silica capillary column of length 30 m and of internal diameter 0,32 mm is recommended; stationary phase DB-5³⁾ (5 % methyl phenyl silicone); thickness of the stationary phase 0,25 µm. The performance of the column should be sufficient to achieve satisfactory separation of all components and similar to that given in the specimen chromatogram (see Figure A.1 for an example).

5.10.3 Detector

An electron-capture detector shall be used with a sensitivity sufficient to detect (twice baseline noise) a 2 µl injection of a 0,001 5 µg/ml solution of *p,p'*-DDT.

6 Sampling and preparation of test sample

6.1 Sampling

Sample the tobacco in accordance with ISO 4874 or all the cigarettes in accordance with ISO 8243. Pay particular attention to ensuring that the test sample is representative of the product as received.

6.2 Preparation of test sample

Dry the tobacco in the oven (5.3) set at 50 °C for 2 h to obtain a mass fraction of moisture of approximately 5 % after drying.

Grind the tobacco through a 2 mm mesh (5.2), taking care to avoid heating above 50 °C. Alternatively the tobacco may be received in a milled form in which case ensure that the mass fraction of moisture is less than 10 %.

Store the tobacco in sealed containers and exclude light. If samples are kept for longer than one month prior to analysis, they shall be stored in a freezer at a temperature below 8 °C.

7 Procedure

7.1 Test portion

Weigh, to the nearest 0,01 g, 5 g tobacco test portions into 50 ml beakers. Add 5 g of pretreated Florisil® (4.8) and mix thoroughly. Carry out the procedures described in 7.2 and 7.3.

7.2 Extraction

Add 5 g of pretreated Florisil® (4.8) to a Soxhlet extractor. Transfer the test portion as prepared in 7.1 without mixing so that two separate layers are formed.

For recovery determinations, appropriate pesticide standard solutions should be added at this stage, by means of a pipette, to the top of the test portion layer.

Transfer 60 ml of the *n*-hexane (4.2) and 1 ml of the internal standard solution (4.6) to a suitable round-bottom flask of capacity 150 ml to 250 ml.

Assemble the extraction apparatus, ensuring good seals at all joints, and turn on the heating mantles (5.5).

³⁾ DB-5 is an example of a suitable product available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this product.