

---

---

**Soil quality — Determination of the  
effects of pollutants on soil flora —  
Screening test for emergence of lettuce  
seedlings (*Lactuca sativa* L.)**

*Qualité du sol — Détermination des effets des polluants sur la flore du  
sol — Essai de détection de l'émergence des plantules de laitue  
(*Lactuca sativa* L.)*

iTeh STANDARD PREVIEW  
(standards.iteh.ai)

[ISO 17126:2005](https://standards.iteh.ai/catalog/standards/sist/075cdfb3-08d9-4c4e-a025-a5896495e44b/iso-17126-2005)

<https://standards.iteh.ai/catalog/standards/sist/075cdfb3-08d9-4c4e-a025-a5896495e44b/iso-17126-2005>



Reference number  
ISO 17126:2005(E)

**PDF disclaimer**

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

ISO 17126:2005

<https://standards.iteh.ai/catalog/standards/sist/075cdfb3-08d9-4c4e-a025-a5896495e44b/iso-17126-2005>

© ISO 2005

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Case postale 56 • CH-1211 Geneva 20  
Tel. + 41 22 749 01 11  
Fax + 41 22 749 09 47  
E-mail [copyright@iso.org](mailto:copyright@iso.org)  
Web [www.iso.org](http://www.iso.org)

Published in Switzerland

# Contents

Page

Foreword .....	iv
Introduction .....	v
1 Scope .....	1
2 Normative references .....	1
3 Terms and definitions .....	1
4 Principle .....	2
5 Materials .....	2
6 Apparatus .....	2
7 Procedure .....	3
7.1 Testing of samples of soil and other test materials .....	3
7.2 Testing of chemicals .....	3
7.3 Temperature and light regime .....	4
7.4 Water content .....	4
7.5 Reference substance .....	4
7.6 Preparation and start of test dishes .....	4
7.7 Test duration .....	5
7.8 Measurements .....	5
7.9 Recordings .....	5
8 Expression of results .....	5
9 Validity of the test .....	6
10 Test report .....	6
Bibliography .....	7

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 17126 was prepared by Technical Committee ISO/TC 190, *Soil quality*, Subcommittee SC 4, *Biological methods*.

## iTeh STANDARD PREVIEW (standards.iteh.ai)

[ISO 17126:2005](https://standards.iteh.ai/catalog/standards/sist/075cdfb3-08d9-4c4e-a025-a5896495e44b/iso-17126-2005)

<https://standards.iteh.ai/catalog/standards/sist/075cdfb3-08d9-4c4e-a025-a5896495e44b/iso-17126-2005>

## Introduction

For the assessment of the suitability of soil to sustain living organisms, there is a need for simple, rapid, inexpensive biological test methods as a complement to chemical analysis. The method described in this International Standard has been developed for the testing of contaminated soil as well as other contaminated samples. It is cost effective and can be conducted within a short period of time. Furthermore, the test organism is easily available, it does not require advanced equipment for measurements or for growing plants, and it can be conducted by any skilled laboratory technician without special training.

This International Standard is based on US EPA 600/3-88-029 (1989)<sup>[1]</sup>.

## iTeh STANDARD PREVIEW (standards.iteh.ai)

[ISO 17126:2005](https://standards.iteh.ai/catalog/standards/sist/075cdfb3-08d9-4c4e-a025-a5896495e44b/iso-17126-2005)

<https://standards.iteh.ai/catalog/standards/sist/075cdfb3-08d9-4c4e-a025-a5896495e44b/iso-17126-2005>

## **iTeh STANDARD PREVIEW (standards.iteh.ai)**

ISO 17126:2005

<https://standards.iteh.ai/catalog/standards/sist/075cdfb3-08d9-4c4e-a025-a5896495e44b/iso-17126-2005>

# Soil quality — Determination of the effects of pollutants on soil flora — Screening test for emergence of lettuce seedlings (*Lactuca sativa* L.)

## 1 Scope

This International Standard specifies test procedures for the determination of effects of contaminated soils or other contaminated samples on the emergence of lettuce seeds.

This International Standard is applicable to contaminated soils, soil materials, compost, sludge and chemical testing. It is applicable to the measurement of effects of substances deliberately added to the soil and to the comparison of soils of known and unknown quality.

This International Standard is not applicable to volatile contaminants.

## 2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 10381-6, *Soil quality — Sampling — Part 6: Guidance on the collection, handling and storage of soil for the assessment of aerobic microbial processes in the laboratory*

ISO 10390, *Soil quality — Determination of pH*

ISO 11265, *Soil quality — Determination of the specific electrical conductivity*

ISO 11267:1999, *Soil quality — Inhibition of reproduction of Collembola (*Folsomia candida*) by soil pollutants*

ISO 11274, *Soil quality — Determination of the water-retention characteristic — Laboratory methods*

ISO 11465, *Soil quality — Determination of dry matter and water content on a mass basis — Gravimetric method*

## 3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

### 3.1

#### **seedling emergence**

appearance of the seedling (i.e. visible seedling) above the surface of the cover material

### 3.2

#### **EC<sub>x</sub>**

concentration of test material (or test substance) estimated to reduce the seedling emergence by *x* % as compared to the control

### 3.3

#### **test material**

material to be tested

EXAMPLES      Soils, soil materials, compost, sludge.

### 3.4

#### **test mixture**

mixture of test material (or test substance) and growth medium

## 4 Principle

Lettuce seeds are exposed to the test material under investigation in a geometric dilution series with test material and growth medium. Incubation takes place under controlled conditions of light and temperature, and lasts usually 5 days (120 h). It is also possible to use this International Standard for chemical testing. In this case, seeds are planted in control pots and in pots containing soil to which the test chemical has been added. At the end of the test, the number of seedlings visible above the sand are counted and recorded. The effect on seedling emergence is expressed as EC50 (possibly EC20), calculated from numbers of emerged seedlings in the control pots (pure growth medium) and in pots containing the test material (or test substance).

## 5 Materials

**5.1 Biological material**, in this case lettuce seeds, *Lactuca sativa* L.

Seeds coated with insecticides and/or fungicides ("dressed" seeds) should be avoided.

After purchase, examine the seeds and remove any trash, empty hulls and damaged seeds. A uniform emergence is dependent on uniform seed size. To reduce variability of emergence, the seed batch may be sized before use by means of four sieves with oblong holes (see 6.4) placed on top of each other. In this case, select for testing the fraction with the largest number of seeds.

Pack the seeds in small portions in air-tight containers. The storage time of the seeds should not exceed the expiration date given by the supplier.

Storage at 4 °C is recommended but good emergence may also be accomplished by storage in the dark at 18 °C.

The seeds should not be soaked in water before testing.

**5.2 Growth medium**, in this case washed, fine quartz sand, e.g. with grain size 0,4 mm to 0,8 mm.

**5.3 Cover material**, in this case washed, coarse quartz sand, e.g. with grain size 0,7 mm to 1,2 mm (possibly 0,8 mm to 1,4 mm).

The coarse quality of the cover material ensures air exchange between the growth medium and the surroundings.

## 6 Apparatus

Standard laboratory equipment (pH-meter, thermometer, pipettes, etc.) including the following.

**6.1 Balance**, with an accuracy of 0,1 g.

**6.2 Lower parts of plastic Petri dishes** (diameter 15 cm), or other containers with similar surface area, for use as test containers.



**6.3 Re-sealable polyethylene bags** that fit the test containers (20 cm × 25 cm for a 15 cm Petri dish).

**6.4 Sieves for seeds**, with oblong mesh dimensions of 0,75 mm × 10 mm, 0,8 mm × 10 mm, 0,85 mm × 10 mm and 0,9 mm × 10 mm.

**6.5 Sieve** for contaminated soil, stainless steel, with mesh size 2 mm.

**6.6 Controlled environment chamber:**

**6.7 Magnifier**

## 7 Procedure

### 7.1 Testing of samples of soil and other test materials

Usually the test material is not dried before the test. If necessary, upon reception air-dry it at room temperature to a water content that enables sieving. Immediately thereafter, sieve the test material through a stainless steel sieve (6.5) and store in the dark at  $4\text{ °C} \pm 2\text{ °C}$  until testing (in accordance with ISO 10381-6). Preferentially, storage should not exceed three months but if prolonged storage is necessary, it shall be stored at  $-18\text{ °C}$ . For sieving, a mesh size of 2 mm is preferable but if this is not possible, coarser sieves (e.g. 5 mm) may be used.

Determine the following properties of the sieved test material and register them prior to the test:

- water content (in accordance with ISO 11465);
- water-holding capacity (in accordance with ISO 11274, or alternatively Annex C of ISO 11267:1999);
- conductivity (in accordance with ISO 11265);
- pH (in accordance with ISO 10390).

The water content and water-holding capacity of the test material and the water-holding capacity of the growth medium are used for calculation of the amounts of water to be used for the test. Before the test, weigh and mix the moist test material and the dry growth medium.

Use a geometric dilution series between test material and growth medium (minimum of 5 concentrations), for which the dilution factor should not exceed two. The range of concentrations should include those at which 0 (or minimum) and 100 % emergence are expected, e.g. based on a preliminary test (7.6.1).

The calculations are based on dry mass, and the concentrations are calculated and expressed as grams dry mass test material per gram dry mass test mixture (i.e. test material and growth medium).

**NOTE** A commonly used dilution factor is  $\sqrt[4]{10}$  which is approximately 1,8, resulting in concentrations of e.g. 10, 18, 32, 56, 100.

### 7.2 Testing of chemicals

#### 7.2.1 General

For testing of chemicals, the test is basically conducted with growth medium only and the chemical is added to the growth medium. The concentration of the chemical is calculated based on the dry mass of the growth medium using the data reported in Clause 8.