

INTERNATIONAL
STANDARD

ISO
8552

IDF
132

First edition
2004-05-01

**Milk — Estimation of psychrotrophic
microorganisms — Colony-count
technique at 21 °C (Rapid method)**

*Lait — Estimation des micro-organismes psychrotrophes — Technique
par comptage des colonies à 21 °C (Méthode rapide)*

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[ISO 8552:2004](https://standards.iteh.ai/catalog/standards/sist/dfc622c2-e9e6-40c9-8a99-1313ceb66750/iso-8552-2004)

<https://standards.iteh.ai/catalog/standards/sist/dfc622c2-e9e6-40c9-8a99-1313ceb66750/iso-8552-2004>



Reference numbers
ISO 8552:2004(E)
IDF 132:2004(E)

© ISO and IDF 2004

PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. Neither the ISO Central Secretariat nor the IDF accepts any liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies and IDF national committees. In the unlikely event that a problem relating to it is found, please inform the ISO Central Secretariat at the address given below.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[ISO 8552:2004](#)

<https://standards.iteh.ai/catalog/standards/sist/dfc622c2-e9e6-40c9-8a99-1313ceb66750/iso-8552-2004>

© ISO and IDF 2004

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO or IDF at the respective address below.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

International Dairy Federation
Diamant Building • Boulevard Auguste Reyers 80 • B-1030 Brussels
Tel. + 32 2 733 98 88
Fax + 32 2 733 04 13
E-mail info@fil-idf.org
Web www.fil-idf.org

Published in Switzerland

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 8552|IDF 132 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 5, *Milk and milk products*, and the International Dairy Federation (IDF), in collaboration with AOAC International. It is being published jointly by ISO and IDF and separately by AOAC International.

ITIH STANDARD PREVIEW
(standards.iteh.ai)

[ISO 8552:2004](https://standards.iteh.ai/catalog/standards/sist/dfc622c2-e9e6-40c9-8a99-1313ceb66750/iso-8552-2004)

<https://standards.iteh.ai/catalog/standards/sist/dfc622c2-e9e6-40c9-8a99-1313ceb66750/iso-8552-2004>

Foreword

IDF (the International Dairy Federation) is a worldwide federation of the dairy sector with a National Committee in every member country. Every National Committee has the right to be represented on the IDF Standing Committees carrying out the technical work. IDF collaborates with ISO and AOAC International in the development of standard methods of analysis and sampling for milk and milk products.

Draft International Standards adopted by the Action Teams and Standing Committees are circulated to the National Committees for voting. Publication as an International Standard requires approval by at least 50 % of IDF National Committees casting a vote.

ISO 8552|IDF 132 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 5, *Milk and milk products*, and the International Dairy Federation (IDF), in collaboration with AOAC International. It is being published jointly by ISO and IDF and separately by AOAC International.

All work was carried out by the Joint ISO/IDF/AOAC Action Team on *Microbiological harmonization*, of the Standing Committee on *Microbiological methods of analysis*, under the aegis of its project leader, Dr J. Floor (ZA).

This first edition of ISO 8552|IDF 132 cancels and replaces the first edition of IDF 132A:1991, which has been editorially and technically revised.

ITeH STANDARD PREVIEW
(standards.iteh.ai)

[ISO 8552:2004](https://standards.iteh.ai/catalog/standards/sist/dfc622c2-e9e6-40c9-8a99-1313ceb66750/iso-8552-2004)

<https://standards.iteh.ai/catalog/standards/sist/dfc622c2-e9e6-40c9-8a99-1313ceb66750/iso-8552-2004>

Introduction

The rapid method described in this International Standard is essentially an approximate method because the incubation conditions are such as to permit organisms other than psychrotrophs to be counted if they grow sufficiently quickly at 21 °C.

Nevertheless, comparative trials in a number of laboratories have shown that the results obtained by the method described here (based on that of Reference [1]) correlate well with those obtained by the method described in ISO 6730|IDF 101 (with incubation at 6,5 °C for 10 days). The latter method (which will be replaced by the horizontal method ISO 17410) should be used if a more accurate enumeration of psychrotrophic microorganisms is required.

iTeh STANDARD PREVIEW **(standards.iteh.ai)**

[ISO 8552:2004](#)

<https://standards.iteh.ai/catalog/standards/sist/dfc622c2-e9e6-40c9-8a99-1313ceb66750/iso-8552-2004>

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 8552:2004

<https://standards.iteh.ai/catalog/standards/sist/dfc622c2-e9e6-40c9-8a99-1313ceb66750/iso-8552-2004>

Milk — Estimation of psychrotrophic microorganisms — Colony-count technique at 21 °C (Rapid method)

1 Scope

This international Standard specifies a rapid method for estimating the number of psychrotrophic microorganisms by means of the colony-count technique at 21 °C.

The method is applicable to raw and pasteurized milk.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 6730, *Milk — Enumeration of colony-forming units of psychrotrophic micro-organisms — Colony count technique at 6,5 degrees C¹⁾*

ISO 7218, *Microbiology of food and animal feeding stuffs — General rules for microbiological examinations*

ISO 8261|IDF 122, *Milk and milk products — General guidance for the preparation of test samples, initial suspensions and decimal dilutions for microbiological examination*

ISO/TS 11133-1, *Microbiology of food and animal feeding stuffs — Guidelines on preparation and production of culture media — Part 1: General guidelines on quality assurance for the preparation of culture media in the laboratory*

ISO/TS 11133-2, *Microbiology of food and animal feeding stuffs — Guidelines on preparation and production of culture media — Part 2: Practical guidelines on performance testing of culture media*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

psychrotrophic microorganisms

bacteria, yeasts and moulds forming countable colonies when incubated aerobically at 6,5 °C for 10 days under the conditions specified in ISO 6730

1) Equivalent to IDF 101A:1991.

4 Principle

- 4.1 Poured plates are prepared using a specified culture medium and a specified quantity of an appropriate dilution of the test sample.
- 4.2 The plates are incubated aerobically at 21 °C for 25 h.
- 4.3 The number of microorganisms per millilitre of sample is calculated from the number of colonies obtained on plates chosen at dilution levels so as to give a significant result.

5 Diluents and culture medium

5.1 General

See ISO 7218 and ISO/TS 11133-1.

5.2 Diluents

See ISO 8261|IDF 122.

5.3 Culture medium — Plate count milk agar

5.3.1 Composition

iTeh STANDARD PREVIEW

(standards.iteh.ai)

Yeast extract	2,5 g
Enzymatic digestion of casein	5,0 g
Skimmed milk powder ^a	1,0 g
Glucose, anhydrous (C ₆ H ₁₂ O ₆)	1,0 g
Agar	9 g to 18 g ^b
Water	1 000 ml

^a The skimmed milk powder shall be free from inhibitory substances.

^b Depending on the gel strength of the agar.

5.3.2 Preparation

5.3.2.1 Preparation from commercial dehydrated medium

Follow the manufacturer's instructions but, in all cases, add the skimmed milk powder even if the manufacturer considers such an addition unnecessary. Adjust the pH, if necessary, so that after sterilization it is 7,0 ± 0,2 at 25 °C.

5.3.2.2 Preparation from dehydrated basic components

Dissolve and disperse in the water, in the following order, the yeast extract, the enzymatic digestion of casein, the glucose and, finally, the skimmed milk powder. Heating the water will assist this procedure. Add the agar and heat to boiling while stirring frequently until the agar is completely dissolved.

Adjust the pH, if necessary, so that after sterilization it is 7,0 ± 0,2 at 25 °C.

5.3.2.3 Distribution, sterilization and storage

Dispense the prepared medium into test tubes (6.6), in quantities of 12 ml to 15 ml per tube, or into flasks or bottles (6.7) of capacity not greater than 500 ml. Sterilize for 15 min in an autoclave (6.10) set at 121 °C.

If the medium is to be used immediately, cool it in a water bath (6.4) to between 44 °C and 47 °C before use.

If not, store the medium in the dark at 3 °C ± 2 °C for no longer than three months (see ISO 7218). Before commencing the microbiological examination and in order to avoid any delay when pouring the medium, completely melt the stored medium, then cool it in a water bath (6.4) at between 44 °C and 47 °C before use (see 8.2.4).

With regard to the temperature check of the medium and other requirements, see ISO 7218.

5.3.3 Performance testing for the quality assurance of the culture medium

Test the performance of the medium in accordance with ISO/TS 11133-2.

6 Apparatus and glassware

Usual microbiological equipment (see ISO 7218) and, in particular, the following.

6.1 Glassware

Disposable glassware is an acceptable alternative to re-usable glassware if it has suitable specifications. Re-usable glassware shall be capable of undergoing repeated sterilization and shall be chemically inert.

6.2 Incubator, capable of operating at 21 °C ± 1 °C.

6.3 pH meter, having an accuracy of calibration of ± 0,1 pH unit at 25 °C.

6.4 Water bath, capable of operating at between 44 °C and 47 °C.

6.5 Colony-counting equipment, consisting, for example, of an illuminated base with a dark background, fitted with a magnifying lens to be used at a magnification of at least 2×, and a mechanical or electronic digital counter.

6.6 Test tubes, of approximate capacity 20 ml, with suitable stoppers.

6.7 Flasks or bottles, of appropriate capacity but not greater than 500 ml, with suitable stoppers.

6.8 Graduated pipettes, of nominal capacity 1 ml.

6.9 Petri dishes, made of glass or plastic, of diameter 90 mm to 100 mm.

6.10 Apparatus for dry sterilization (oven) or wet sterilization (autoclave).

See ISO 7218.

7 Sampling

It is important that the laboratory receive a sample which is truly representative and has not been damaged or changed during transport or storage.

Sampling is not part of the method specified in this International Standard. A recommended sampling method is given in ISO 707.