



# SLOVENSKI STANDARD SIST ISO 3960:2003

01-maj-2003

F Ugh]bg\_Y]b'y]j Ug\_Ya Uy cVY]b'c`U!`8c`c Yj Ub^Y'dYfc\_g]XbY[ UyHYj ]U

Animal and vegetable fats and oils -- Determination of peroxide value

Corps gras d'origines animale et végétale -- Détermination de l'indice de peroxyde

ITEH STANDARD PREVIEW  
(standards.iteh.ai)

Ta slovenski standard je istoveten z: **ISO 3960:2001**

[SIST ISO 3960:2003](#)

<https://standards.iteh.ai/catalog/standards/sist/5032e547-3dc8-4062-8a5b-fb975de188f8/sist-iso-3960-2003>

**ICS:**

67.200.10	Üæ dā • \ ^ Á Á ã ç \ ^ { æ [ à ^ Á Á  æ	Animal and vegetable fats and oils
-----------	---	---------------------------------------

**SIST ISO 3960:2003**

**en**

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

SIST ISO 3960:2003

<https://standards.iteh.ai/catalog/standards/sist/5032e547-3dc8-4062-8a5b-fb975de188f8/sist-iso-3960-2003>

# INTERNATIONAL STANDARD

**ISO  
3960**

Third edition  
2001-12-01

---

---

## **Animal and vegetable fats and oils — Determination of peroxide value**

*Corps gras d'origines animale et végétale — Détermination de l'indice de peroxyde*

**iTeh STANDARD PREVIEW  
(standards.iteh.ai)**

[SIST ISO 3960:2003](https://standards.iteh.ai/catalog/standards/sist/5032e547-3dc8-4062-8a5b-fb975de188f8/sist-iso-3960-2003)

<https://standards.iteh.ai/catalog/standards/sist/5032e547-3dc8-4062-8a5b-fb975de188f8/sist-iso-3960-2003>



Reference number  
ISO 3960:2001(E)

© ISO 2001

**PDF disclaimer**

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

## iTeh STANDARD PREVIEW (standards.iteh.ai)

[SIST ISO 3960:2003](https://standards.iteh.ai/catalog/standards/sist/5032e547-3dc8-4062-8a5b-fb975de188f8/sist-iso-3960-2003)

<https://standards.iteh.ai/catalog/standards/sist/5032e547-3dc8-4062-8a5b-fb975de188f8/sist-iso-3960-2003>

© ISO 2001

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Case postale 56 • CH-1211 Geneva 20  
Tel. + 41 22 749 01 11  
Fax + 41 22 749 09 47  
E-mail [copyright@iso.ch](mailto:copyright@iso.ch)  
Web [www.iso.ch](http://www.iso.ch)

Printed in Switzerland

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 3.

Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this International Standard may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

International Standard ISO 3960 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 11, *Animal and vegetable fats and oils*.

This third edition cancels and replaces the second edition (ISO 3960:1998), which has been technically revised.

Annex A of this International Standard is for information only.

ITeH STANDARD PREVIEW  
(standards.iteh.ai)

SIST ISO 3960:2003  
<https://standards.iteh.ai/catalog/standards/sist/5032e547-3dc8-4062-8a5b-fb975de188f8/sist-iso-3960-2003>

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

SIST ISO 3960:2003

<https://standards.iteh.ai/catalog/standards/sist/5032e547-3dc8-4062-8a5b-fb975de188f8/sist-iso-3960-2003>

# Animal and vegetable fats and oils — Determination of peroxide value

## 1 Scope

This International Standard specifies a method for the determination of the peroxide value of animal and vegetable fats and oils.

## 2 Normative references

The following normative documents contain provisions which, through reference in this text, constitute provisions of this International Standard. For dated references, subsequent amendments to, or revisions of, any of these publications do not apply. However, parties to agreements based on this International Standard are encouraged to investigate the possibility of applying the most recent editions of the normative documents indicated below. For undated references, the latest edition of the normative document referred to applies. Members of ISO and IEC maintain registers of currently valid International Standards.

ISO 661, *Animal and vegetable fats and oils — Preparation of test sample*

ISO 3696:1987, *Water for analytical laboratory use — Specification and test methods*

## 3 Term and definition

[SIST ISO 3960:2003](https://standards.iteh.ai/catalog/standards/sist/5032e547-3dc8-4062-8a5b-b9751e89110a/iso-3960-2001)

[https://standards.iteh.ai/catalog/standards/sist/5032e547-3dc8-4062-8a5b-](https://standards.iteh.ai/catalog/standards/sist/5032e547-3dc8-4062-8a5b-b9751e89110a/iso-3960-2001)

For the purposes of this International Standard, the following term and definition applies.

### 3.1

#### peroxide value

quantity of those substances in the sample, expressed in terms of active oxygen, which oxidize potassium iodide under the conditions specified in this International Standard, divided by the mass of the test portion

NOTE 1 Peroxide value is usually expressed in industry as milliequivalents per kilogram.

NOTE 2 Peroxide value may also be expressed (in SI units) as millimoles per kilogram. The value expressed in millimoles per kilogram is half that expressed in milliequivalents per kilogram (see clause 10).

## 4 Principle

A test portion, in solution in acetic acid and isooctane, is treated with a solution of potassium iodide. The liberated iodine is titrated with a standard volumetric sodium thiosulfate solution.

## 5 Reagents

Use only reagents of recognized analytical grade, unless otherwise stated. All reagents and the water shall be free of dissolved oxygen.

**5.1 Water**, complying with grade 3 of ISO 3696.

**5.2 Glacial acetic acid**, freed from oxygen by purging with a current of pure and dry inert gas (carbon dioxide or nitrogen).

## ISO 3960:2001(E)

**WARNING — Glacial acetic acid is moderately toxic by ingestion and inhalation. It is a strong irritant to skin and tissue.**

**5.3 Isooctane**, freed from oxygen by purging with a current of pure and dry inert gas (carbon dioxide or nitrogen).

**WARNING — Isooctane is flammable and a fire risk. Explosive limits in air are 1,1 % (by volume) to 6,0 % (by volume). It is toxic by ingestion and inhalation. A properly operating fume hood should be used when working with this solvent.**

**5.4 Acetic acid/isooctane solution** [60:40 (by volume)], prepared by mixing 3 volumes of glacial acetic acid (5.2) with 2 volumes of isooctane (5.3).

**5.5 Potassium iodide solution**, saturated, recently prepared and free from free iodine and iodates.

Make sure the solution remains saturated as indicated by the presence of undissolved crystals. Store in the dark. Test daily by adding 2 drops of starch solution (5.8) to 0,5 ml of the potassium iodide solution in 30 ml of the acetic acid/isooctane solution (5.4). If a blue colour is formed which requires more than 1 drop of sodium thiosulfate solution (5.7) to discharge the colour, discard the potassium iodide solution and prepare a fresh solution.

**5.6 Sodium thiosulfate solution**,  $c(\text{Na}_2\text{S}_2\text{O}_3) = 0,1 \text{ mol/l}$ , standardized just before use.

Dissolve 24,9 g of sodium thiosulfate pentahydrate ( $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$ ) in distilled water and dilute to 1 litre.

**5.7 Sodium thiosulfate solution**,  $c(\text{Na}_2\text{S}_2\text{O}_3) = 0,01 \text{ mol/l}$ , standardized just before use.

Prepare by diluting previous solution (5.6)

**5.8 Starch solution**, 5 g/l.

Mix 1 g of starch and a small amount of cold distilled water. Add this mixture, while stirring, to 200 ml of boiling water. Add 250 mg of salicylic acid as preservative and boil for 3 min. Immediately remove from the heat and cool.

If long storage is required, keep the solution in a refrigerator at between 4 °C and 10 °C. A fresh starch solution shall be prepared when the endpoint of the titration from blue to colourless fails to be sharp. If stored under refrigeration, the starch solution should be stable for about 2 to 3 weeks.

The sensitivity of the starch solution may be tested as follows. To 5 ml of the starch solution in 100 ml of water, add 0,05 % of potassium iodide solution (5.5) and 1 drop of 0,05 % sodium hypochlorite solution. It is essential that the deep blue colour produced be discharged by 0,05 ml of sodium thiosulfate solution (5.6).

## 6 Apparatus

All equipment used shall be free from reducing or oxidizing substances. Do not grease ground glass surfaces.

Usual laboratory apparatus and, in particular, the following.

**6.1 Conical flasks**, of 250 ml capacity, with ground glass stoppers.

## 7 Sampling

It is important that the laboratory receive a sample that is truly representative and has not been damaged or changed during transport or storage.

Sampling is not part of the method specified in this International Standard. A recommended sampling method is given in ISO 5555 <sup>[1]</sup>.

Ensure that the sample is taken and stored away from strong light, kept cold, and contained in completely filled dark-coloured glass containers, hermetically sealed with ground glass stoppers.



## 8 Preparation of test sample

Ensure that the packaging of the fat or oil is not damaged, and is well closed. When parameters other than the peroxide value have to be investigated, the test portion for the peroxide value shall be taken first from the laboratory sample.

Prepare the test sample in accordance with ISO 661.

## 9 Procedure

### 9.1 General

Carry out the test in artificial light or diffuse daylight.

NOTE If it is required to check that the repeatability requirement (see 11.2) is met, carry out two single determinations in accordance with 9.2 and 9.3.

### 9.2 Test portion

Rinse the conical flask (6.1) with a current of pure dry inert gas (carbon dioxide or nitrogen). Into the flask, weigh a mass of the sample to the accuracy given in Table 1 and in accordance with the expected peroxide value.

**Table 1 — Mass of test portion and accuracy of weighing**

Expected peroxide value meq/kg	Mass of test portion g	Weighing accuracy g
0 to 12	5,0 to 2,0	± 0,01
12 to 20	2,0 to 1,2	± 0,01
20 to 30	1,2 to 0,8	± 0,01
30 to 50	0,8 to 0,5	± 0,001
50 to 90	0,5 to 0,3	± 0,001

### 9.3 Determination

Add 50 ml of acetic acid/isooctane solution (5.4) to the conical flask and replace the stopper. Swirl the flask until the sample has dissolved. Using a suitable volumetric pipette, add 0,5 ml of saturated potassium iodide solution (5.5) and replace the stopper. Allow the solution to react for 1 min ± 1 s, thoroughly shaking it at least three times during this period, then immediately add 30 ml of distilled water.

Titrate the solution with sodium thiosulfate solution (5.7), adding it gradually and with constant, vigorous agitation, until the yellow iodine colour has almost disappeared. Add about 0,5 ml of starch solution (5.8) and continue the titration with constant agitation, especially near the endpoint, to liberate all of the iodine from the solvent layer, adding the sodium thiosulfate solution (5.7) drop by drop until the blue colour just disappears.

There is a 15 s to 30 s delay in neutralizing the starch indicator for peroxide values of 70 meq/kg and greater, due to the tendency of isooctane to float on the surface of the aqueous medium and the time necessary to mix adequately the solvent and the aqueous titrant, thus liberating the last traces of iodine. A small amount [0,5 % to 1,0 % (*m/m*)] of a high HLB emulsifier (e.g. Tween 60)<sup>1)</sup> may be added to the reaction mixture to retard the phase separation and decrease the time lag in the liberation of iodine.

1) Tween 60 is an example of a suitable product available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this product.