# INTERNATIONAL STANDARD

ISO 10539

First edition 2002-03-15

# Animal and vegetable fats and oils — Determination of alkalinity

Corps gras d'origines animale et végétale — Détermination de l'alcalinité

# iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 10539:2002 https://standards.iteh.ai/catalog/standards/sist/251f6fa6-5afc-4bde-b599-621a903ae02d/iso-10539-2002



### **PDF** disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

# iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 10539:2002 https://standards.iteh.ai/catalog/standards/sist/251f6fa6-5afc-4bde-b599-621a903ae02d/iso-10539-2002

#### © ISO 2002

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.ch
Web www.iso.ch

Printed in Switzerland

# **Foreword**

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 3.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this International Standard may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 10539 was prepared by Technical Committee ISO/TC 34, Food products, Subcommittee SC 11, Animal and vegetable fats and oils.

Annex A of this International Standard is for information only.

(standards.iteh.ai)

ISO 10539:2002 https://standards.iteh.ai/catalog/standards/sist/251f6fa6-5afc-4bde-b599-621a903ae02d/iso-10539-2002

# iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 10539:2002 https://standards.iteh.ai/catalog/standards/sist/251f6fa6-5afc-4bde-b599-621a903ae02d/iso-10539-2002

# Animal and vegetable fats and oils — Determination of alkalinity

# 1 Scope

This International Standard specifies a method for the determination of the alkalinity of animal and vegetable fats and oils without distinguishing between the various constituents. The method is not applicable to dry melted animal fats, nor to oils and fats with an acidity greater than 60 % (mass fraction) as determined in accordance with ISO 660.

NOTE Fats and oils can contain alkaline constituents either naturally (e.g. calcium soaps from bones) or accidentally (e.g. sodium soaps in imperfectly refined fats and oils).

# 2 Normative references

The following normative documents contain provisions which, through reference in this text, constitute provisions of this International Standard. For dated references, subsequent amendments to, or revisions of, any of these publications do not apply. However, parties to agreements based on this International Standard are encouraged to investigate the possibility of applying the most recent editions of the normative documents indicated below. For undated references, the latest edition of the normative document referred to applies. Members of ISO and IEC maintain registers of currently valid International Standards.

ISO 660:1996, Animal and vegetable fats and bils - Determination of acid value and acidity

ISO 661:1989, Animal and vegetable fats and oils Preparation of test sample

https://standards.iteh.ai/catalog/standards/sist/251f6fa6-5afc-4bde-b599-ISO 3696:1987, Water for analytical laboratory use Specification and test methods

### 3 Term and definition

For the purposes of this International Standard, the following term and definition apply.

# 3.1

# alkalinity

proportion of alkali in a fat or oil determined by titration with hydrochloric acid, in accordance with the method specified in this International Standard

# 4 Principle

A test portion is dissolved in warm aqueous acetone and titrated with hydrochloric acid.

# 5 Reagents

Use only reagents of recognized analytical grade, unless otherwise stated.

**5.1** Water, conforming to grade 3 of ISO 3696.

### 5.2 Aqueous acetone

Mix 970 ml of acetone, 20 ml of water and 10 ml of bromophenol blue solution (2 g/l in acetone). Add sodium hydroxide solution [c(NaOH) = 0,1 mol/l] to give a blue colour, then add hydrochloric acid [c(HCI) = 0,1 mol/l] until a yellowish green colour appears.

**5.3 Hydrochloric acid standard volumetric solution**, c(HCI) = 0.01 mol/l, standardized not more than 7 days before use.

# 6 Apparatus

Usual laboratory apparatus and, in particular, the following.

- **6.1** Borosilicate glass apparatus, for aqueous acetone (5.2).
- **6.2** Conical flasks, with wide necks, and of capacities 400 ml to 500 ml.

# 7 Sampling

It is important that the laboratory receive a sample which is truly representative and has not been damaged or changed during transport or storage.

Sampling is not part of the method specified in this International Standard. A recommended sampling method is given in ISO 5555.

# 8 Preparation of test sample

Prepare the test sample in accordance with ISO 661.

# 9 Procedure

# iTeh STANDARD PREVIEW

# 9.1 Preparation of the apparatus (standards.iteh.ai)

Rinse each conical flask (6.2) and stirrer to be used with successive 20 ml amounts of aqueous acetone (5.2) until the aqueous acetone does not change colour. Allow the flasks and stirrer to dry.

https://standards.iteh.ai/catalog/standards/sist/251f6fa6-5afc-4bde-b599-621a903ae02d/iso-10539-2002

# 9.2 Test portion

From the test sample (clause 8) weigh, to the nearest 0,1 g, a test portion of up to 40 g (see Table 1) according to the expected result and colour of the sample. Melt a solid test portion at below the boiling point of acetone. Add 100 ml of the acetone (5.2) at 40 °C and stir. Allow to settle until the solution splits in two layers; in the presence of soap, the upper layer is blue-coloured.

Table 1 — Mass of test portion

| Colour of sample | Expected result               | Mass |
|------------------|-------------------------------|------|
|                  | mg sodium oleate/kg of sample | g    |
| Light            | up to 500                     | 40   |
| Light            | over 500 to 1 000             | 10   |
| Light            | over 1 000                    | 4    |
| Dark             | not relevant                  | 2    |
| Very dark        | not relevant                  | 1    |

For test portions of less than 10 g, take replicate portions totalling at least 5 g for titration and sum the masses and titres to give the result for one determination.

### 9.3 Determination

Titrate the warm (30 °C to 40 °C) solution with the hydrochloric acid (5.3) while stirring until the colour of the indicator changes from blue (or blue green) to the yellowish green of the acetone (5.2) added. Allow the mixture to settle in order to observe the colour clearly.

NOTE An explosion-resistant hotplate with magnetic stirrer and burette support is particularly useful.

#### 9.4 Number of determinations

Carry out two determinations on the same test sample, treating the replications for small test portions as one determination (see Table 1).

# 10 Expression of results

Alkalinity may be expressed as a mass, in milligrams, of sodium oleate per kilogram of fat or oil, or as a percentage (mass fraction) of sodium hydroxide:

a) the alkalinity in terms of sodium hydroxide is given by the expression

$$A = \frac{4Vc}{m}$$

b) the alkalinity in terms of sodium oleate is given by the expression

$$S = 304\ 000\ \frac{Vc}{m}$$
 iTeh STANDARD PREVIEW (standards.iteh.ai)

where

ISO 10539:2002

https://standards.iteh.ai/catalog/standards/sist/251f6fa6-5afc-4bde-b599-

- A is the alkalinity expressed as a percentage (mass fraction) of sodium hydroxide;
- S is the alkalinity expressed as milligrams of sodium oleate per kilogram of sample;
- m is the mass of the test portion, in grams;
- c is the exact concentration of the standard volumetric hydrochloric acid solution used, in moles per litre;
- V is the volume of hydrochloric acid standard volumetric solution used to titrate the test portion, in millilitres.

Take as the result the mean of two determinations, provided that the requirement for repeatability (see 11.2) is met.

# 11 Precision

### 11.1 Interlaboratory test

Details of an interlaboratory test are summarized in annex A. The values derived from this interlaboratory test may not be applicable to concentration ranges other than those given.

### 11.2 Repeatability

The absolute difference between two independent single test results, obtained using the same method on identical test material in the same laboratory by the same operator using the same equipment within a short interval of time, will in not more than 5 % of cases exceed the repeatability limit r given in Table A.1 for the relevant level of the mean value determined.

# 11.3 Reproducibility

The absolute difference between two single test results, obtained using the same method on identical test material in different laboratories by different operators using different equipment, will in not more than 5 % of cases exceed the reproducibilty limit *R* given in Table A.1 for the relevant level of the mean value determined.

# 12 Test report

The test report shall specify:

- all information necessary for the complete identification of the sample;
- the sampling method used, if known;
- the test method used, with reference to this International Standard;
- all operating conditions not specified in this International Standard, or regarded as optional, together with details of any incidents which may have influenced the test results;
- the test results obtained or, if the repeatability has been checked, the final result obtained.

# iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 10539:2002 https://standards.iteh.ai/catalog/standards/sist/251f6fa6-5afc-4bde-b599-621a903ae02d/iso-10539-2002

# Annex A (informative)

# Results of an interlaboratory trial

An international collaborative test has been carried out on the method given in this International Standard. The test, involving fifteen laboratories in six countries, was organized by AFNOR and the results obtained were subjected to statistical analysis in accordance with ISO 5725-2 to give the precision data reported in Table A.1.

Table A.1 — Precision data

|  | Sample D | Sample E | Sample H |
|--|----------|----------|----------|
| Number of laboratories   | 15       | 15       | 15       |
| Number of laboratories retained after eliminating outliers         | 14       | 11       | 12       |
| Mean alkalinity (mg/kg)  | 8,3      | 216,3    | 12,5     |
| Repeatability standard deviation, $s_r$ (mg/kg)                    | 0,7      | 0,6      | 0,5      |
| Repeatability coefficient of variation (%)                         | 8,1      | 0,3      | 3,6      |
| Repeatability limit, r (mg/kg)                                     | 1,9      | 1,8      | 1,3      |
| Reproducibility standard deviation, s <sub>R</sub> (mg/kg) TANDARI | PR23VIE  | 8,4      | 9,5      |
| Reproducibility coefficient of variation (%) (standards.i          | teh.¾i)  | 3,9      | 76,1     |
| Reproducibility limit, R (mg/kg)                                   | 6,4      | 23,7     | 27,0     |

ISO 10539:2002

https://standards.iteh.ai/catalog/standards/sist/251f6fa6-5afc-4bde-b599-621a903ae02d/iso-10539-2002