INTERNATIONAL STANDARD

ISO 3033-1

First edition 2005-09-01

Oil of spearmint —

Part 1:

Native type (Mentha spicata L.)

Huile essentielle de menthe crépue (ou menthe verte) —

Partie 1: Type native (Mentha spicata L.)

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 3033-1:2005 https://standards.iteh.ai/catalog/standards/sist/b733b36b-3c93-45bf-919f-235f2e41177e/iso-3033-1-2005



PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 3033-1:2005 https://standards.iteh.ai/catalog/standards/sist/b733b36b-3c93-45bf-919f-235f2e41177e/iso-3033-1-2005

© ISO 2005

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 3033-1 was prepared by Technical Committee ISO/TC 54, Essential oils.

This first edition of ISO 3033-1, together with ISO 3033-2:2005, ISO 3033-3:2005 and ISO 3033-4:2005, cancels and replaces ISO 3033:1988, which has been technically revised.

ISO 3033 consists of the following parts, under the general title *Oil of spearmint*:

- Part 1: Native type (Mentha spicata L.) ISO 3033-1:2005 https://standards.iteh.a/catalog/standards/sist/b733b36b-3c93-45bf-919f-
- Part 2: Chinese type (80 % and 60 %) (Mentha viridis L. var. crispa Benth.), redistilled oil
- Part 3: Indian type (Mentha spicata L.), redistilled oil
- Part 4: Scotch variety (Mentha × gracilis Sole)

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 3033-1:2005 https://standards.iteh.ai/catalog/standards/sist/b733b36b-3c93-45bf-919f-235f2e41177e/iso-3033-1-2005

Oil of spearmint —

Part 1.

Native type (Mentha spicata L.)

Scope

This part of ISO 3033 specifies certain characteristics of the oil of spearmint, native type (Mentha spicata L.), in order to facilitate assessment of its quality.

Normative references

following referenced documents indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO/TR 210, Essential oils — General tules of or 1:2005

packaging, conditioning and storage hai/catalog/standards/sist/4713b3Appearance19f-235f2e41177e/iso-3033-1-2005

ISO/TR 211, Essential oils — General rules for labelling and marking of containers

ISO 212, Essential oils — Sampling

ISO 279, Essential oils — Determination of relative density at 20 °C — Reference method

ISO 280, Essential oils — Determination of refractive index

ISO 592, Essential oils — Determination of optical

ISO 875, Essential oils — Evaluation of miscibility in ethanol

ISO 1271, Essential oils — Determination of carbonyl value — Free hydroxylamine method

ISO 11024-1, Essential oils — General guidance on chromatographic profiles — Part 1: Preparation of chromatographic profiles for presentation in standards

ISO 11024-2, Essential oils — General guidance on chromatographic profiles — Part 2: Utilization of chromatographic profiles of samples of essential oils

Terms and definitions

For the purposes of this document, the following terms and definitions apply.

oil of spearmint, native type

essential oil obtained by steam distillation from the fresh above-ground parts of the flowering plant of Mentha spicata L., of the Lamiaceae family

For information on the CAS number, see ISO/TR 21092.

.iteh.ai)

Requirements

Clear mobile liquid.

4.2 Colour

From colourless to pale yellow.

4.3 Odour

Characteristic odour of carvone with a herbaceous note.

Relative density at 20 °C, d_{20}^{20}

Minimum: 0,921 0

0,938 0 Maximum:

4.5 Refractive index at 20 °C

Minimum: 1.484 0

1.4910 Maximum:

4.6 Optical rotation at 20 °C

Between -59° and -48°.

4.7 Miscibility in ethanol, 70 % (volume fraction), at 20 °C

It shall not be necessary to use more than 3 volumes of ethanol, 70 % (volume fraction), to obtain a clear solution with 1 volume of essential oil.

4.8 Carbonyl value

Minimum: 224, corresponding to a carbonyl compound content of 60 %, expressed as carvone.

4.9 Chromatographic profile

Analysis of the essential oil shall be carried out by gas chromatography. In the chromatogram obtained, the representative and characteristic components shown in Table 1 shall be identified. The proportions of these components, indicated by the integrator, shall be as shown in Table 1. This constitutes the chromatographic profile of the essential oil.

Table 1 — Chromatographic profile

rable i — omomatograpine prome		
Component	Minimum %tps://stan	Maximum ^{Ls} dards.ite%ai/catalo
Limonene ^a	9	15 ^{23512e4} 1
3-Octanol	0,6	1,4
Menthone b	_	0,2
trans-Sabinene hydrate	0,5	1,0
cis-Dihydrocarvone	1,0	2,5
Carvone ^c	60,0	70,0
trans-Dihydrocarvyl acetate	0,1	0,6
cis-Carvyl acetate	0,1	0,6
cis-Jasmone	0,2	0,7
β-Bourbonene	1,0	2,0
Viridiflorol	0,1	0,5

^a The limonene found is regarded to be predominantly L-limonene based on the physical testing. It is believed that there might be a small amount of D-limonene present but the exact quantity is unknown.

NOTE The chromatographic profile is normative, contrary to typical chromatograms given for information in Annex A.

4.10 Flashpoint

Information on the flashpoint is given in Annex B.

5 Sampling

See ISO 212.

Minimum volume of test sample: 50 ml.

NOTE This volume allows each of the tests specified in this part of ISO 3033 to be carried out at least once.

6 Test methods

6.1 Relative density at 20 °C, d_{20}^{20}

See ISO 279.

6.2 Refractive index at 20 °C

standards.iteh.ai)

6.2 Ontical rotation at 20 °C

6.3 Optical rotation at 20 °C O 3033-1:2005

ystand**sele/iso**b**592**b36b-3c93-45bf-919f-177e/iso-3033-1-2005

6.4 Miscibility in ethanol, 70 % (volume fraction), at 20 °C

See ISO 875.

6.5 Carbonyl value

See ISO 1271.

Test portion: 1 g.

Reflux time: 3 h.

Relative molar mass of carvone: 150,21.

6.6 Chromatographic profile

See ISO 11024-1 and ISO 11024-2.

7 Packaging, labelling, marking and storage


See ISO/TR 210 and ISO/TR 211.

^b The menthone found is regarded to be predominantly L-menthone based on the physical testing. It is believed that there might be a small amount of D-menthone present but the exact quantity is unknown.

^C The carvone found is regarded to be predominantly L-carvone based on the physical testing. It is believed that there might be a small amount of D-carvone present but the exact quantity is unknown.

Annex A (informative)

Typical chromatograms of the analysis by gas chromatography of the essential oil of spearmint, native type (*Mentha spicata* L.)



Peak identification

1 Myrcene + 3-octanol

2 Limonene

3 trans-Sabinene hydrate

4 Menthone

5 cis-Dihydrocarvone

6 Carvone

7 trans-Dihydrocarvyl acetate

8 cis-Carvyl acetate

9 *cis*-Jasmone

10 β-Bourbonene11 Viridiflorol

Operating conditions

Column: fused silica capillary; length 30 m; internal diameter 0,25 mm

Stationary phase: poly(dimethyl siloxane) (DB-1®)

Film thickness: 0,25 µm

Oven temperature: isothermal at 75 °C for 5 min, then temperature programming from 75 °C to 100 °C at a rate of 5 °C/min, then from 100 °C to 220 °C at a rate of 6 °C/min

and isothermal at 220 °C for 8,34 min Injector temperature: 230 °C


Detector temperature: 260 °C

Detector: flame ionization type Carrier gas: helium

Volume injected: 0,1 µl Carrier gas flow rate: 1 ml/min

Split ratio: 1/100

Figure A.1 — Typical chromatogram taken on an apolar column



Peak identification

Myrcene Column: fused silic

2 Limonene

3 3-Octanol

4 Menthone

4 Menthone

5 *trans*-Sabinene hydrate

6 β -Bourbonene

7 cis-Dihydrocarvone

8 trans-Dihydrocarvyl acetate

9 Carvone

10 cis-Carvyl acetate

11 cis-Jasmone

12 Viridiflorol

Operating conditions

Column: fused silica capillary; length 30 m; internal diameter 0,25 mm

Stationary phase: poly(ethylene glycol) (DB-WAX®)

Film thickness: 0,25 µm

Oven temperature: isothermal at 75 °C for 5 min, then temperature programming from 75 °C to 100 °C at a rate of 5 °C/min, then from 100 °C to 220 °C at a rate of 6 °C/min

and isothermal at 220 °C for 8,34 min

Injector temperature: 230 °C

Detector temperature: 260 °C

Detector: flame ionization type

Carrier gas: helium

Volume injected: 0,1 µl

Carrier gas flow rate: 1 ml/min

Split ratio: 1/100

Figure A.2 — Typical chromatogram taken on a polar column

Annex B (informative)

Flashpoint

B.1 General information

For safety reasons, transport companies, insurance companies, and people in charge of safety services require information on the flashpoints of essential oils, which in most cases are flammable products.

A comparative study of the relevant methods of analysis (see ISO/TR 11018) concluded that it was difficult to recommend a single apparatus for standardization purposes, given that:

there is wide variation in the chemical composition of essential oils;

Consequently, it was decided to give a mean value for the flashpoint in an informative annex to each International Standard, in order to meet the requirements of the interested parties.

The equipment with which this value was obtained should be specified.

For further information, see ISO/TR 11018.

B.2 Flashpoint of the essential oil of spearmint, native type

The mean value is +58 °C.

 the volume of the sample needed for certain requirements would be too costly for high? NOTE P Obtained with "Setaflash" equipment. priced essential oils;

(standards.iteh.ai)

— as there are several different types of equipment which can be used for the determination, users cannot be expected to use one specified type only.

235f2e41177e/iso-3033-1-2005