INTERNATIONAL STANDARD

ISO 8968-5

IDF 20-5

First edition 2001-12-15

Milk — Determination of nitrogen content —

Part 5:

Determination of protein-nitrogen content

iTeh Shait — Détermination de la teneur en azote —
Partie 5: Détermination de la teneur en azote protéique
(standards.iteh.ai)

ISO 8968-5:2001

https://standards.iteh.ai/catalog/standards/sist/f682c597-17ae-420c-919c-8541a49885a7/iso-8968-5-2001



PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. Neither the ISO Central Secretariat nor the IDF accepts any liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies and IDF national committees. In the unlikely event that a problem relating to it is found, please inform the ISO Central Secretariat at the address given below.

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 8968-5:2001 https://standards.iteh.ai/catalog/standards/sist/f682c597-17ae-420c-919c-8541a49885a7/iso-8968-5-2001

© ISO and IDF 2001

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO or IDF at the respective address below.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.ch
Web www.iso.ch

Printed in Switzerland

International Dairy Federation
41 Square Vergote • B-1030 Brussels
Tel. + 32 2 733 98 88
Fax + 32 2 733 04 13

E-mail info@fil-idf.org Web www.fil-idf.org

Contents Page Foreword......iv 1 2 3 Term and definition1 4 5 Reagents ______2 Apparatus ______2 6 7 8 Preparation of test sample......3 9 9.1 9.2 Direct determination......3 9.3 Indirect determination4 Calculation and expression of results DARD PREVIEW 4 10 Calculation of protein-nitrogen content......4 10.1 10.2 Precision......5 11 11.1 Repeatability https://standards.iteh.ai/catalog/standards/sist/f682c597-17ae-420c-919c-5 11.2 11.3 12 Test report5

Bibliography 6

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 3.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this part of ISO 8968 IDF 20 may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

International Standard ISO 8968-5 IDF 20-5 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 5, *Milk and milk products*, and the International Dairy Federation (IDF), in collaboration with AOAC International. It is being published jointly by ISO and IDF and separately by AOAC International.

ISO 8968 IDF 20 consists of the following parts, under the general title Milk — Determination of nitrogen content:

— Part 1: Kjeldahl method

- ISO 8968-5:2001
- Part 2: Block-digestion method (Macro method) standards/sist/f682c597-17ae-420c-919c-8541a49885a7/iso-8968-5-2001
- Part 3: Block-digestion method (Semi-micro rapid routine method)
- Part 4: Determination of the non-protein-nitrogen content
- Part 5: Determination of the protein-nitrogen content

Foreword

IDF (the International Dairy Federation) is a worldwide federation of the dairy sector with a National Committee in every member country. Every National Committee has the right to be represented on the IDF Standing Committees carrying out the technical work. IDF collaborates with ISO and AOAC International in the development of standard methods of analysis and sampling for milk and milk products.

Draft International Standards adopted by the Action Teams and Standing Committees are circulated to the National Committees for voting. Publication as an International Standard requires approval by at least 50 % of National Committees casting a vote.

International Standard ISO 8968-5 | IDF 20-5 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 5, *Milk and milk products*, and the International Dairy Federation (IDF), in collaboration with AOAC International. It is being published jointly by ISO and IDF and separately by AOAC International.

All work was carried out by the Joint ISO/IDF/AOAC Action Team, *Nitrogen compounds*, under the aegis of its project leader, Mr D.M. Barbano (US).

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 8968-5:2001 https://standards.iteh.ai/catalog/standards/sist/f682c597-17ae-420c-919c-8541a49885a7/iso-8968-5-2001

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 8968-5:2001

https://standards.iteh.ai/catalog/standards/sist/f682c597-17ae-420c-919c-8541a49885a7/iso-8968-5-2001

Milk — Determination of nitrogen content —

Part 5:

Determination of protein-nitrogen content

WARNING — The use of this part of ISO 8968 | IDF 20 may involve the use of hazardous materials, operations, and equipment. This standard does not purport to address all the safety risks associated with its use. It is the responsibility of the user of this standard to establish appropriate safety and healthy practices and determine the applicability of local regulatory limitations prior to use.

1 Scope

This part of ISO 8968 IDF 20 specifies a method for the direct determination of the protein-nitrogen content of liquid milk, whole or skimmed.

An alternative indirect method using calculations is also described.

(standards.iteh.ai)

2 Normative references

The following normative documents contain provisions which, through reference in this text, constitute provisions of this part of ISO 8968 IDF 20. For dated references, subsequent amendments to, or revisions of, any of these publications do not apply. However, parties to agreements based on this part of ISO 8968 IDF 20 are encouraged to investigate the possibility of applying the most recent editions of the normative documents indicated below. For undated references, the latest edition of the normative document referred to applies. Members of ISO and IEC maintain registers of currently valid International Standards.

ISO 8968-1 IDF 20-1:2001, Milk — Determination of nitrogen content — Part 1: Kjeldahl method

ISO 8968-2 | IDF 20-2:2001, Milk — Determination of nitrogen content — Part 2: Block-digestion method (Macro method)

ISO 8968-4 | IDF 20-4:2001, Milk — Determination of nitrogen content — Part 4: Determination of the non-protein-nitrogen content

3 Term and definition

For the purposes of this part of ISO 8968 IDF 20, the following term and definition apply.

3.1

protein-nitrogen content

mass fraction of substances determined by the procedure specified in this part of ISO 8968 | IDF 20, directly or indirectly

NOTE The protein-nitrogen content is expressed as a percentage by mass.

4 Principle

Protein is precipitated from a test portion by addition of trichloroacetic acid solution such that the final concentration of trichloroacetic acid in the mixture is approximately 12 %. The protein precipitate is separated by filtration. (The filtrate contains the non-protein-nitrogen contents.) The nitrogen content of the filtrate is determined by the procedure described either in part 1 or part 2 of ISO 8968 | IDF 20.

5 Reagents

Use only reagents of recognized analytical grade, unless otherwise specified, and distilled or demineralized water or water of equivalent purity.

The reagents are as specified for the determination of total nitrogen described in part 1 or part 2 of ISO 8968 IDF 20, together with the following. A laboratory may decide which method to use.

5.1 Trichloroacetic acid solution (CCI₃COOH)

Dissolve 15,0 g of trichloroacetic acid in water in a 100 ml one-mark volumetric flask. Dilute to the mark with water. Do not use concentrations of trichloroacetic acid and volumes of solutions other than those specified.

The performance of the method with respect to mean value and between-laboratory performance characteristics will differ if other than the specified concentrations of trichloroacetic acid and volumes of solutions are used.

5.2 Hydrochloric acid standard volumetric solution, $c(HCI) = (0.1 \pm 0,000 5) \text{ mol/l.}$

It is recommended that this material be purchased prestandardized by the manufacturer to meet or exceed the above specification.

ISO 8968-5:2001

6 Apparatus

https://standards.iteh.ai/catalog/standards/sist/f682c597-17ae-420c-919c-8541a49885a7/iso-8968-5-2001

Usual laboratory apparatus and, in particular, the following, together with the apparatus specified in part 1 or part 2 of ISO 8968 | IDF 20.

- **6.1** Water bath, capable of being maintained at 38 °C \pm 2 °C.
- **6.2** Pipette, of capacities 5 ml.
- **6.3** Filter funnel, made of glass, of diameter 75 mm.
- **6.4** Filter paper, nitrogen free, of diameter 15 cm (e.g. Whatman No. 1¹⁾) or equivalent.
- **6.5** Automatic pipette or piston pump, capable of delivering 10 ml.

7 Sampling

Sampling is not part of the method specified in this International Standard. A recommended sampling method is given in ISO 707.

It is important that the laboratory receive a sample which is truly representative and has not been damaged or changed during transport or storage.

¹⁾ Whatman is an example of a suitable product available commercially. This information is given for the convenience of users of this part of ISO 8968 | IDF 20 and does not constitute an endorsement by ISO of this product.

8 Preparation of test sample

Warm the test sample in the water bath (6.1) set at 38 °C. Gently mix the test sample thoroughly by repeatedly inverting the sample bottle without causing frothing or churning. Cool the sample to room temperature immediately prior to weighing the test portion (9.1).

9 Procedure

9.1 Test portion

Pipette 5,0 ml \pm 0,1 ml of the prepared test sample (clause 8) either into a dry and clean Kjeldahl flask or digestion tube, preweighed to the nearest 0,1 mg. Weigh the test sample to the nearest 0,1 mg. Immediately add 5 ml \pm 0,1 ml of water to the flask or tube, rinsing any test sample on its neck into the flask or tube.

NOTE 1 The use of either a Kjeldahl flask or digestion tube depends on the laboratory's choice of method.

NOTE 2 For advice on test portion size to apply this method to dairy products other than milk, see annex A of ISO 8968-1 | IDF 20-1:2001.

9.2 Direct determination

9.2.1 Precipitation and filtration

Add 40 ml \pm 0,5 ml of trichloroacetic acid solution (5.1) to the Kjeldahl flask or digestion tube containing the test portion (9.1) and swirl to mix the contents. Let the flask or tube stand for approximately 5 min to allow the precipitate to settle. Pour the contents of the flask or tube through a filter paper (6.4) placed in a filter funnel (6.3). Collect the filtrate in a clean conical flask. Some of the precipitate will remain in the Kjeldahl flask or digestion tube and some will be collected on the filter paper. It is not necessary to remove all of the precipitate from the flask or tube.

https://standards.itch.ai/catalog/standards/sist/f682c597-17ae-420c-919c-

Immediately after pouring the mixture and so as not to allow any precipitate to dry on the neck of the flask or tube, add by means of an automatic pipette (6.5), 10 ml of the trichloroacetic acid solution (5.1). Use the solution to rinse any precipitate from the neck of the flask or tube down into it. Swirl to mix the contents. Pour the thus-obtained contents of the flask or tube through the same filter paper. Add the filtrate to that collected previously in the conical flask. Again, immediately rinse the neck of the flask or tube with a further 10 ml of trichloroacetic acid solution and swirl to mix the contents. Pour the contents of the flask or tube for the third time through the same filter paper, adding the filtrate to that collected previously in the conical flask.

The obtained filtrate shall be clear and free of particulate matter. At this point, the filtrate is no longer needed and may be discarded in an appropriate manner.

If duplicated tests of the same test sample are to be done, then two separate precipitation and filtration procedures shall be carried out for each test sample.

9.2.2 Preparation of the filtrate

Wearing gloves, carefully remove the filter paper from the filter funnel and fold the paper to enclose the precipitate. If any precipitate remains on either the inner or outer lip of the Kjeldahl flask or digestion tube, wipe with the folded filter paper so that any precipitate adheres to the paper and then drop the filter paper also into the Kjeldahl flask or digestion tube.

9.2.3 Digestion and distillation

Add the appropriate amount of boiling aids, potassium sulfate, coper catalyst solution and sulfuric acid either to the Kjeldahl flask or digestion tube and continue with the digestion and distillation by the procedure described in either part 1 or part 2 of ISO 8968 | IDF 20.

A laboratory may choose to use either part 1 or part 2 of ISO 8968 IDF 20.