INTERNATIONAL STANDARD

ISO 21150

First edition 2006-04-15

Cosmetics — Microbiology — Detection of *Escherichia coli*

Cosmétiques — Microbiologie — Détection d'Escherichia coli

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 21150:2006 https://standards.iteh.ai/catalog/standards/sist/6d7173f2-6964-43d8-98c9-d06dfb4b3f26/iso-21150-2006



PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 21150:2006 https://standards.iteh.ai/catalog/standards/sist/6d7173f2-6964-43d8-98c9-d06dfb4b3f26/iso-21150-2006

© ISO 2006

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Page

Contents

Forewo	ord	. i\		
Introdu	ıction	۱		
1	Scope	′		
2	Normative references	1		
3	Terms and definitions	1		
4	Principle	2		
5	Diluents and culture media			
5.1 5.2	General Diluent for the bacterial suspension (tryptone sodium chloride solution)			
5.2 5.3	Culture media			
6	Apparatus and glassware	6		
7	Strains of microorganisms			
8	Handling of cosmetic products and laboratory samples	6		
9	Procedure iTeh STANDARD PREVIEW	(
9.1	General recommendations	6		
9.2	Preparation of the initial suspension in the enrichment broth			
9.3 9.4	Incubation of the inoculated enrichment broth Detection and identification of Escherichia coli			
10	Expression of the results (detection of Escherichia cofi) 64-43d8-98c9-			
11	d06dfb4b3f26/iso-21150-2006 Neutralization of the antimicrobial properties of the product			
11.1	General			
11.2	Preparation of inoculum	9		
11.3	Validation of the detection method	9		
12	Test report			
Annex	Annex A (informative) Other enrichment broths1			
Annex B (informative) Neutralizers of antimicrobial activity of preservatives and rinsing liquids 14				
Ribliography 14				

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 21150 was prepared by Technical Committee ISO/TC 217, Cosmetics.

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 21150:2006 https://standards.iteh.ai/catalog/standards/sist/6d7173f2-6964-43d8-98c9-d06dfb4b3f26/iso-21150-2006

Introduction

Microbiological examinations of cosmetic products are to be carried out according to an appropriate microbiological risk analysis in order to ensure their quality and safety for consumers.

Microbiological risk analysis depends on several parameters such as:

- potential alteration of cosmetic products;
- pathogenicity of microorganisms;
- site of application of the cosmetic product (hair, skin, eyes, mucous membranes, etc.);
- type of users (adults, children under 3 years, etc.).

For cosmetics and other topical products, the detection of skin pathogens such as *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Candida albicans* may be relevant. The detection of other kinds of microorganisms might be of interest since these microorganisms (including indicators of faecal contamination, e.g. *Escherichia coli*) suggest hygienic failure during manufacturing process.

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 21150:2006 https://standards.iteh.ai/catalog/standards/sist/6d7173f2-6964-43d8-98c9-d06dfb4b3f26/iso-21150-2006

© ISO 2006 - All rights reserved

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 21150:2006 https://standards.iteh.ai/catalog/standards/sist/6d7173f2-6964-43d8-98c9-d06dfb4b3f26/iso-21150-2006

Cosmetics — Microbiology — Detection of Escherichia coli

1 Scope

This International Standard gives general guidelines for the detection and identification of the specified microorganism *Escherichia coli* in cosmetic products. Microorganisms considered as specified in this International Standard might differ from country to country according to national practices or regulations.

In order to ensure product quality and safety for consumers, it is advisable to perform an appropriate microbiological risk analysis, so as to determine the types of cosmetic products to which this International Standard is applicable. Products considered to present a low microbiological risk include those with low water activity, hydro-alcoholic products, extreme pH values, etc.

This International Standard specifies a method that is based on the detection of *Escherichia coli* in a non-selective liquid medium (enrichment broth), followed by isolation on a selective agar medium. Other methods may be appropriate depending on the level of detection required.

NOTE For the detection of *Escherichia coli*, subcultures can be performed on non-selective culture media followed by suitable identification steps (e.g. using identification kits).

Because of the large variety of cosmetic products within this field of application, this method might not be suited to some products in every detail (e.g. certain water-immiscible products). Other International Standards may be appropriate. Other methods (e.g. automated) can be substituted for the test presented here provided that their equivalence has been demonstrated or the method has been otherwise validated.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 21148:—1), Cosmetics — Microbiology — General instructions for microbiological examination

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

product

portion of an identified cosmetic product received in the laboratory for testing

3.2

sample

portion of the product (at least 1 g or 1 ml) which is used in the test to prepare the initial suspension

¹⁾ To be published.

3.3

initial suspension

suspension (or solution) of the sample in a defined volume of an appropriate enrichment broth

3.4

sample dilution

dilution of the initial suspension

3.5

specified microorganism

aerobic mesophilic bacteria or yeast which is undesirable in a cosmetic product because it can cause skin or eye infection, or it can be recognized as an indicator of hygienic failure in the manufacturing process

3.6

Escherichia coli

Gram-negative rod, motile, smooth colonies

NOTE 1 The main characteristics for identification are catalase positive, oxidase negative, fermentation of lactose, production of indole, growth on selective medium containing bile salts with characteristic colonies.

NOTE 2 Escherichia coli can be isolated from the moist environmental sources (air, water, soil) and is a faecal contamination indicator.

3.7

enrichment broth

non-selective liquid medium containing suitable neutralizers and/or dispersing agents and validated for the product under test

(standards.iteh.ai)

4 Principle

ISO 21150:2006

The first step of the procedure is to perform an enrichment by using a non-selective broth medium to increase the number of microorganisms without the risk of inhibition by the selective ingredients that are present in selective/differential growth media.

The second step (isolation) of the test is performed on a selective medium followed by identification tests.

The possible inhibition of microbial growth by the sample shall be neutralized to allow the detection of viable microorganisms ^[5]. In all cases and whatever the methodology, the neutralization of the antimicrobial properties of the product shall be checked and validated ^[6] ^[7] ^[8].

5 Diluents and culture media

5.1 General

Use the general instructions given in ISO 21148. When water is mentioned in this document, use distilled water or purified water as specified in ISO 21148.

The enrichment broth is used to disperse the sample and to increase the initial microbial population. It may contain neutralizers if the specimen to be tested has antimicrobial properties. The efficacy of the neutralization shall be demonstrated (see Clause 11). Information relative to suitable neutralizers is given in Annex B.

The following enrichment broth is suitable for checking the presence of *Escherichia coli* according to this International Standard provided that it is validated according to Clause 11.

Other diluents and culture media may be used if they have been demonstrated to be suitable for use.

5.2 Diluent for the bacterial suspension (tryptone sodium chloride solution)

The diluent is used for the preparation of bacterial suspension used for the validation procedure (see Clause 11).

5.2.1 Composition

Tryptone, pancreatic digest of casein	1,0 g
Sodium chloride	8,5 g
Water	1 000 ml

5.2.2 Preparation

Dissolve the components in water by mixing while heating. Dispense into suitable containers. Sterilize in the autoclave at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to 7.0 ± 0.2 when measured at room temperature.

5.3 Culture media

5.3.1 General iTeh STANDARD PREVIEW

Culture media may be prepared using the descriptions provided below or from dehydrated culture media, according to the instructions from the manufacturer. The instructions provided by the supplier of the media should be followed.

ISO 21150:2006

NOTE Ready-to-use media may, be used when their composition and/or growth-yields are comparable to those of the formulas given herein.

d06dfb4b3f26/iso-21150-2006

5.3.2 Agar medium for validation [soybean-casein digest agar medium (SCDA) or tryptic soy agar (TSA)]

5.3.2.1 Composition

Pancreatic digest of casein	15,0 g
Papaic digest of soybean meal	5,0 g
Sodium chloride	5,0 g
Agar	15,0 g
Water	1 000 ml

5.3.2.2 Preparation

Dissolve the components or the dehydrated complete medium in the water by mixing while heating. Dispense the medium into suitable containers. Sterilize in the autoclave at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to 7.3 ± 0.2 when measured at room temperature.

© ISO 2006 – All rights reserved

5.3.3 Enrichment broth

5.3.3.1 Eugon LT 100 broth

5.3.3.1.1 General

This medium contains ingredients

- which neutralize inhibitory substances present in the sample: lecithin and polysorbate 80,
- dispersing agent: octoxynol 9.

5.3.3.1.2 Composition

Pancreatic digest of casein	15,0 g
Papaic digest of soybean meal	5,0 g
L-cystine	0,7 g
Sodium chloride	4,0 g
Sodium sulfite	0,2 g

Glucose iTeh STAND 55RD PREVIEW

Egg lecithin (standar,ds.iteh.ai)

Polysorbate 80 ISO 5,0 9,2006

Octoxynol 9 https://standards.iteh.ai/catalog/standards/sist/6d7173f2-6964-43d8-98c9-d06dfb4b3f26089-21150-2006

Water 1 000 ml

5.3.3.1.3 Preparation

Dissolve the components, polysorbate 80, octoxynol 9 and egg lecithin, successively into boiling water until their complete dissolution. Dissolve the other components by mixing while heating.

Dispense the medium into suitable containers. Sterilize in the autoclave at 121 $^{\circ}$ C for 15 min. After sterilization and cooling down, the pH shall be equivalent to 7,0 \pm 0,2 when measured at room temperature.

5.3.3.2 Other enrichment broths

Other enrichment broths may be used as appropriate (see Annex A).

5.3.4 Selective agar medium for isolation of Escherichia coli

5.3.4.1 MacConkey agar medium

5.3.4.1.1 Composition

Pancreatic digest of gelatin	17,0 g
Pancreatic digest of casein	1,5 g
Peptic digest of animal tissue	1,5 g

Lactose	10,0 g
Bile salts mixture	1,5 g
Sodium chloride	5,0 g
Agar	13,5 g
Neutral red	30,0 mg
Crystal violet	1,0 mg
Water	1 000 ml

5.3.4.1.2 Preparation

Dissolve all solid components in the water and boil for 1 min to effect solution.

Dispense in suitable containers and sterilize at 121 °C for 15 min.

The pH, after sterilization and cooling down, shall be equivalent to 7.1 ± 0.2 when measured at room temperature.

5.3.5 Selective agar medium for confirmation of Escherichia coli

5.3.5.1 Levine eosin-methylene blue agar medium PREVIEW

5.3.5.1.1 Composition (standards.iteh.ai)

Pancreatic digest of gelatin

ISO 21150:210(0 g

https://standards.iteh.ai/catalog/standards/sist/6d7173f2-6964-43d8-98c9Potassium dihydrogen phosphate (KH2PQ4)6/iso-21250-g006

Agar

15,0 g

Lactose

10,0 g

Eosine Y 400 mg

Methylene blue 65 mg

Water 1 000 ml

5.3.5.1.2 Preparation

Dissolve the pancreatic digest of gelatin, the dibasic potassium phosphate, and the agar in the water, with warming, and allow to cool. Just prior to use, liquefy the gelled agar solution, add the remaining ingredients, as solutions, in the following amounts, and mix; for each 100 ml of the liquefied agar solution

- 5 ml of 20 % lactose solution.
- 2 ml of 2 % eosin Y solution, and
- 2 ml of 0,033 % methylene blue solution.

The finished medium may not be clear.

Dispense in suitable containers and sterilize at 121 °C for 15 min.

© ISO 2006 – All rights reserved 5