
**Tobacco and tobacco products —
Preparation and constitution of identical
samples from the same lot for
collaborative studies for the evaluation of
test methods**

*Tabac et produits du tabac — Préparation et constitution d'échantillons
identiques à partir d'un même lot pour la conduite d'essais comparatifs
portant sur la qualité des méthodes d'essai*

[iteh standards](https://standards.iteh.ai)
(<https://standards.iteh.ai>)
Document Preview

[ISO/TS 7821:2005](https://standards.iteh.ai/catalog/standards/iso/ebac41c3-5158-4a4f-af4d-eca5ce8d9cb2/iso-ts-7821-2005)

<https://standards.iteh.ai/catalog/standards/iso/ebac41c3-5158-4a4f-af4d-eca5ce8d9cb2/iso-ts-7821-2005>



PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

iTeh Standards
(<https://standards.iteh.ai>)
Document Preview

[ISO/TS 7821:2005](https://standards.iteh.ai/catalog/standards/iso/ebac41c3-5158-4a4f-af4d-eca5ce8d9cb2/iso-ts-7821-2005)

<https://standards.iteh.ai/catalog/standards/iso/ebac41c3-5158-4a4f-af4d-eca5ce8d9cb2/iso-ts-7821-2005>

© ISO 2005

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

In other circumstances, particularly when there is an urgent market requirement for such documents, a technical committee may decide to publish other types of normative document:

- an ISO Publicly Available Specification (ISO/PAS) represents an agreement between technical experts in an ISO working group and is accepted for publication if it is approved by more than 50 % of the members of the parent committee casting a vote;
- an ISO Technical Specification (ISO/TS) represents an agreement between the members of a technical committee and is accepted for publication if it is approved by 2/3 of the members of the committee casting a vote.

An ISO/PAS or ISO/TS is reviewed after three years in order to decide whether it will be confirmed for a further three years, revised to become an International Standard, or withdrawn. If the ISO/PAS or ISO/TS is confirmed, it is reviewed again after a further three years, at which time it must either be transformed into an International Standard or be withdrawn.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO/TS 7821 was prepared by Technical Committee ISO/TC 126, *Tobacco and tobacco products*, Subcommittee SC 1, *Physical and dimensional tests*.

This edition cancels and replaces ISO/TR 7821:1982, which has been technically revised.

Introduction

The repeatability and the reproducibility of a particular laboratory method are defined in ISO 3534-1, ISO 3534-2 and ISO 3534-3. These characteristics may be determined by means of interlaboratory tests as described in ISO 5725-2.

In addition to the description of the equipment and the test method to be used, often by reference to existing standards, the organization of such tests also includes the preparation of test materials, i.e. samples that will serve as media for the studies to be carried out.

It should be noted that, in this type of study, there is rarely any interest in the characteristics of the test material itself, provided that such characteristics are within the typical range for the materials commonly tested, and values are determined in this way only in order to obtain the quantitative data necessary for compiling a statistical report concerning the test method.

Thus, in most cases, every effort is made to obtain lots or samples which are as similar to each other as possible, in such a way as to reduce the residual variance of the tests and, in the same way, to increase the accuracy of the conclusions that may be drawn from the global analysis of the results.

Since the individual test samples do not necessarily need to represent the population from which they are taken, but only have to be identical to each other, the method for their preparation can be quite different from those needed when the aim of the study is to obtain knowledge of the original population.

In the case of certain industries, the chemical industry for example, it is relatively easy to make up samples which are practically identical to each other, as the structure of the material allows it to be divided up so as to conserve the microscopic or macroscopic homogeneity of the product (e.g. solutions, powders).

In the case of tobacco, however, it is completely different because the raw materials, and even the finished products, are affected by an intrinsic heterogeneity likely to cause considerable differences between samples if special precautions are not taken.

Tobacco, as a test medium, can occur in the following forms:

- powder;
- cut tobacco;
- cigarettes;
- leaves or strips.

The method for the preparation of samples of test media depends, in most cases, on the type of test to be carried out. Nevertheless, it is possible to lay down a few general guidelines applicable to a large number of collaborative studies, depending on the form in which the material occurs.

Tobacco and tobacco products — Preparation and constitution of identical samples from the same lot for collaborative studies for the evaluation of test methods

1 Scope

This Technical Specification provides a practical framework for the operations of dividing a lot (not necessarily homogeneous) of tobacco or tobacco products into several similar samples intended for collaborative studies, either within or between laboratories.

2 Preparation of samples

2.1 General

The division of a lot into samples that are similar to each other is very easy if the lot is homogeneous (or can be made so, according to its physical state), but may be more difficult if it is not. In this case, an appropriate sampling procedure has to be used in order that the influence of heterogeneity is as small as possible.

It is obviously not possible to describe in detail every case that may arise. The aim of the following examples, therefore, is to illustrate the principles of, and to give guidance on, the operations to be carried out, which, moreover, appear simple since

- on the one hand, the given aim is to make up samples that are as similar as possible, and
- on the other hand, to take into account the fact that two units will probably have characteristics which are most similar if they are sampled from points which are physically close together.

2.2 Powders

2.2.1 Principle

Consider a lot to be divided comprising N g. It is to be divided into n samples of N/n g. Each sample consists of a number of increments.

Each increment is taken separately from the lot, and should, in theory, be allocated at random to each of the n samples. In practice, it is sufficient to allocate the increments to each of the n samples in rotation, provided that the number of cycles of rotation (i.e. number of increments in each sample) is large.

2.2.2 Precautions to be taken

During the preparation of the samples, the following precautions should be taken:

- a) work in an atmosphere which has a relative humidity as close as possible to that corresponding to the equilibrium humidity of the material being handled;
- b) work as quickly as possible in order to limit the exchange of moisture between the atmosphere and the material if it is not possible to satisfy completely condition a).

2.2.3 Example

A gross sample of 1 kg of powder is to be divided into 10 samples of 100 g. The condition to be fulfilled for the collaborative studies is not that 10 samples of exactly 100 g each should be obtained, but that each sample of approximately 100 g should be similar, from the physical and chemical point of view, to each of the others.

a) Manual method

After mixing the gross sample as thoroughly as possible, place the pile on a tray or in a receptacle from which the material can be easily scooped out. Use a spoon, ladle or measure, of capacity between 5 g and 10 g for example. Provide 10 boxes capable of holding at least 100 g each.

Taking a succession of increments, using the spoon, ladle or measure, feed, in turn, box 1, box 2, ..., box 10.

Each sample is thus made up of at least 10 (10 to 20) increments.

When the initial pile is almost exhausted, i.e. when there is only enough left for one round of 10 increments, it is recommended that distribution be finished with a measure of smaller capacity.

This method guarantees that the 10 samples thus constituted are as similar as possible. Any mixing of the gross sample will cause variation of particle size and density through the pile from top to bottom, and this should be taken into account when taking increments.

b) Automatic method

Use a mechanical device such that, at each revolution, the quantity attributed to each sample is less than or equal to one-tenth of the total mass of the final sample.

2.3 Cut tobacco

2.3.1 Principle

Exactly the same principle as for powders is applicable. In practice, handling is more difficult as the tobacco can form tangles, especially if it is in long strands.

The temptation to make up the increments from wads or lumps should be resisted and, instead, they should be broken down so that they can be divided between all the samples.

As a general rule, each sample should be made up of at least 10 increments (and more if possible). In addition, increments should always be distributed to each sample in a standard rotation.

The increment should be adapted to the size of the lot and the number of samples to be obtained.

2.3.2 Special precautions

Cut tobacco always tends to settle, i.e. handling causes a certain amount of separation of the short strands which collect at the bottom of the receptacle.

In order to avoid causing an artificial heterogeneity between samples when the long strands have been exhausted, it is important to distribute the short strands equally between samples using an instrument such as a scoop or spoon, which prevents the loss of these strands at the time of distribution.