



SLOVENSKI STANDARD
SIST EN 16198:2013

01-maj-2013

Nadomešča:

SIST-TS CEN/TS 16198:2011

Gnojila - Kompleksometrijsko določevanje magnezija

Fertilizers - Determination of magnesium by complexometry

Düngemittel - Komplexometrische Bestimmung von Magnesium

Engrais - Dosage du magnésium par complexométrie

iTeh STANDARD PREVIEW
(standards.iteh.ai)

Ta slovenski standard je istoveten z: ~~SIST EN 16198:2012~~ EN 16198:2012

<https://standards.iteh.ai/catalog/standards/sist/6cd91767-d642-4552-8b0d-a58e91e25d44/sist-en-16198-2013>

ICS:

65.080

Gnojila

Fertilizers

SIST EN 16198:2013

en,fr,de

iTeh STANDARD PREVIEW
(standards.iteh.ai)

SIST EN 16198:2013

<https://standards.iteh.ai/catalog/standards/sist/6cd91767-d642-4552-8b0d-a58c91c25d44/sist-en-16198-2013>

EUROPEAN STANDARD

EN 16198

NORME EUROPÉENNE

EUROPÄISCHE NORM

November 2012

ICS 65.080

Supersedes CEN/TS 16198:2011

English Version

Fertilizers - Determination of magnesium by complexometry

Engrais - Dosage du magnésium par complexométrie

Düngemittel - Komplexometrische Bestimmung von
Magnesium

This European Standard was approved by CEN on 6 October 2012.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.

SIST EN 16198:2013

<https://standards.iteh.ai/catalog/standards/sist/6cd91767-d642-4552-8b0d-a58c91c25d44/sist-en-16198-2013>



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

Management Centre: Avenue Marnix 17, B-1000 Brussels

Contents

	Page
Foreword.....	3
1 Scope	4
2 Normative references	4
3 Terms and definitions	4
4 Principle.....	4
5 Sampling.....	4
6 Reagents.....	5
7 Apparatus	6
8 Preparation of the extraction solution.....	6
9 Procedure	6
9.1 Control test.....	6
9.2 Aliquot samples to be taken.....	7
9.3 Titration in the presence of Eriochrome black-T.....	7
9.4 Titration in the presence of calcein or of calcon carbonic acid	7
10 Calculation and expression of the result.....	7
11 Remarks.....	7
12 Precision.....	8
12.1 Inter-laboratory test.....	8
12.2 Repeatability.....	8
12.3 Reproducibility.....	8
13 Test report	9
Annex A (informative) Statistical results of the inter-laboratory test	10
Bibliography.....	12

ITeH STANDARD PREVIEW
 (standards.iteh.ai)

SIST EN 16198:2013

<https://standards.iteh.ai/catalog/standards/sist/6cd91767-d642-4552-8b0d-a58c91c25d44/sist-en-16198-2013>

Foreword

This document (EN 16198:2012) has been prepared by Technical Committee CEN/TC 260 “Fertilizers and liming materials”, the secretariat of which is held by DIN.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by May 2013, and conflicting national standards shall be withdrawn at the latest by May 2013.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

This document supersedes CEN/TS 16198:2011.

The following changes have been made to the former edition:

- a) the CEN Technical Specification has been adopted as a European Standard;
- b) the document has been editorially revised.

This document has been prepared under a mandate given to CEN by the European Commission and the European Free Trade Association.

According to the CEN/CENELEC Internal Regulations, the national standards organisations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

EN 16198:2012 (E)**1 Scope**

This European Standard specifies a method for the determination of magnesium in fertilizer extracts by complexometry.

The method is applicable to the following EC fertilizer extracts for which the determination of total magnesium and/or water-soluble magnesium is provided for according to the Regulation (EC) No 2003/2003, Annex I [3]:

- fertilizers listed in [3], Annex I: straight nitrogenous fertilizers, type 1b + 1c (calcium magnesium nitrate), type 7 (magnesium sulfonitrate), type 8 (nitrogenous fertilizers with magnesium) and straight potassic fertilizers, type 2 (enriched kainite), type 4 (potassium chloride containing magnesium), type 6 (potassium sulfate containing magnesium salt);
- fertilizers listed in [3], Annex I D relating to secondary nutrients.

2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN 1482-2, *Fertilizers and liming materials — Sampling and sample preparation — Part 2: Sample preparation*

EN 12944-1:1999, *Fertilizers and liming materials and soil improvers — Vocabulary — Part 1: General terms*

EN 12944-2:1999, *Fertilizers and liming materials and soil improvers — Vocabulary — Part 2: Terms relating to fertilizers*

EN 15960, *Fertilizers — Extraction of total calcium, total magnesium, total sodium and total sulfur in the forms of sulfates*

SIST EN 16198:2013
<https://standards.iteh.ai/catalog/standards/sist/6cd91767-d642-4552-8b0d-a58c91c25d44/sist-en-16198-2013>

EN 15961, *Fertilizers — Extraction of water soluble calcium, magnesium, sodium and sulfur in the form of sulfates*

3 Terms and definitions

For the purposes of this document, the terms and definitions given in EN 12944-1:1999 and EN 12944-2:1999 apply.

4 Principle

The magnesium is solubilised according to EN 15960 or EN 15961. First titration: with EDTA of Ca and Mg in the presence of Eriochrome black-T. Second titration: with EDTA of Ca in the presence of calcein or of calcane carbonic acid. Determination of magnesium by difference.

5 Sampling

Sampling is not part of the method specified in this document. A recommended sampling method is given in EN 1482-1.

Sample preparation shall be carried out in accordance with EN 1482-2. Grinding of the laboratory sample is recommended for homogeneity reasons.

6 Reagents

6.1 Magnesium standard solution, $c = 0,05 \text{ mol/l}$.

6.1.1 Dissolve 1,232 g of magnesium sulfate ($\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$) in the hydrochloric acid solution (6.11) and make up to 100 ml with the same acid,

or

6.1.2 Weigh 2,016 g of magnesium oxide, previously calcined to remove all traces of carbonation. Place it in a beaker with 100 ml of water.

Stir in approximately 120 ml of hydrochloric acid (6.12).

After dissolution, transfer quantitatively into a graduated 1 000 ml flask. Make up to volume and mix.

1 ml of these solutions should contain 1,216 mg of Mg (= 2,016 mg of MgO).

The laboratory is responsible for testing the strength of this standard solution.

6.2 EDTA solution, $c = 0,05 \text{ mol/l}$.

Weigh 18,61 g of the dihydrated disodium salt of ethylenediaminetetraacetic (EDTA) ($\text{C}_{10}\text{H}_{14}\text{N}_2\text{Na}_2\text{O}_8 \cdot 2\text{H}_2\text{O}$), place it in a beaker (7.4) and dissolve in 600 ml to 800 ml of water. Transfer the solution quantitatively into a graduated flask (7.3). Make up the volume and mix. Check this solution with the standard solution (6.1) by taking a sample of 20 ml of the latter and by titration according to the analytical procedure described in 9.2.

1 ml of the EDTA solution should correspond to 1,216 mg of Mg (= 2,016 mg of MgO) and to 2,004 mg of Ca (= 2,804 mg CaO) (see remarks 11.1 and 11.6).

6.3 Calcium standard solution, $c = 0,05 \text{ mol/l}$.

Weigh 5,004 g of dry calcium carbonate. Place it in a beaker with 100 ml of water. Progressively stir in 120 ml of hydrochloric acid (6.12).

Bring to the boil in order to drive off the carbon dioxide, cool, transfer quantitatively into a graduated 1 l flask, make up the volume with water and mix. Check this solution against the EDTA solution (6.2) following analytical procedure (9.3). 1 ml of this solution should contain 2,004 mg of Ca (= 2,804 mg of CaO) and should correspond to 1 ml of the EDTA solution (6.2).

6.4 Calcein indicator.

Carefully mix in a mortar 1 g of calcein with 100 g of sodium chloride. Use 10 mg of this mixture. The indicator changes from green to orange. Titration shall be carried out until an orange is obtained which is free from green tinges.

6.5 Calcon carbonic acid indicator.

Dissolve 400 mg of calcon carbonic in 100 ml of methanol. This solution may only be kept for approximately four weeks. Use three drops of this solution. The indicator changes from red to blue. Titration shall be carried out until a blue is obtained which is free from red tinges.

6.6 Eriochrome black-T indicator.

Dissolve 300 mg of Eriochrome black-T in a mixture of 25 ml of propanol-1 and 15 ml of triethanolamine. This solution may only be kept for approximately four weeks. Use three drops of this solution. This indicator turns from red to blue and titration shall be carried out until a blue is obtained which is free from red tinges. It

EN 16198:2012 (E)

changes colour only when magnesium is present. If necessary add 1 ml of the magnesium standard solution (6.1).

When both calcium and magnesium are present the EDTA first forms a complex with the calcium and then with the magnesium. In that case two elements are determined concurrently.

6.7 Potassium cyanide solution, aqueous solution of $w(\text{KCN}) = 2\%$.

SAFETY PRECAUTIONS — Do not pipette by mouth and see 11.7.

6.8 Solution of potassium hydroxide and potassium cyanide.

Dissolve 280 g of KOH and 66 g of KCN in water, make up the volume to 1 l and mix.

6.9 Buffer solution, pH = 10,5.

In a graduated flask (7.5), dissolve 33 g of ammonium chloride in 200 ml of water, add 250 ml of ammonia ($\rho_{20} = 0,91$ g/ml) make up the volume with water and mix. Check the pH of the solution regularly.

6.10 Diluted hydrochloric acid, one volume of hydrochloric acid ($\rho_{20} = 1,18$ g/ml) plus one volume of water.

6.11 Hydrochloric acid solution, approximately $c = 0,5$ mol/l.

6.12 Hydrochloric acid solution, approximately $c = 1$ mol/l.

6.13 Sodium hydroxide solution, $c = 5$ mol/l.

STANDARD PREVIEW
(standards.iteh.ai)

7 Apparatus

7.1 Magnetic or mechanical stirrer.

[SIST EN 16198:2013](https://standards.iteh.ai/catalog/standards/sist/6cd91767-d642-4552-8b0d-a58c91c25d44/sist-en-16198-2013)

<https://standards.iteh.ai/catalog/standards/sist/6cd91767-d642-4552-8b0d-a58c91c25d44/sist-en-16198-2013>

7.2 pH meter.

7.3 1 000 ml graduated flask.

7.4 1 000 ml beaker.

7.5 500 ml graduated flask.

7.6 400 ml beaker.

8 Preparation of the extraction solution

Prepare the extraction solution according to EN 15960 or EN 15961.

9 Procedure

9.1 Control test

Carry out a determination on aliquot parts of solutions (6.1 and 6.3) such that the Ca/Mg ratio is approximately equal to that of the solution to be analysed. To this end take v_1 ml of Ca standard solution (6.3) and $(v_2 - v_1)$ ml of standard solution (6.1), where v_1 and v_2 are the volumes, in millilitres, of the EDTA solution used in the two titrations performed on the solution to be analysed. This procedure is correct only if the solutions of EDTA, calcium and magnesium are exactly equivalent. If this is not the case, it is necessary to make corrections.

9.2 Aliquot samples to be taken

The aliquot part will as far as possible contain between 9 mg and 18 mg of magnesium (= 15 mg to 30 mg of MgO).

9.3 Titration in the presence of Eriochrome black-T

Pipette an aliquot part (9.2) of the solution to be analysed into a beaker (7.6). Neutralise the surplus acid with the sodium hydroxide solution (6.13) using the pH meter. Dilute with water to approximately 100 ml. Add 5 ml of the buffer solution (6.9). The pH measured shall be $10,5 \pm 0,1$. Add 2 ml of the potassium cyanide solution (6.7) and three drops of the Eriochrome black-T indicator (6.6). Titrate with the EDTA solution (6.2). Stir gently with the stirrer (7.1) (see 11.2, 11.3 and 11.4). Take v_2 as the volume, in millilitres, of the EDTA solution (6.2).

9.4 Titration in the presence of calcein or of calcon carbonic acid

Pipette an aliquot part of the solution to be analysed equal to that taken from the above titration (9.3) and place it in a beaker (7.6). Neutralise the surplus acid with the sodium hydroxide solution (6.13) using the pH meter. Dilute with water to about 100 ml. Add 10 ml of the KOH/KCN solution (6.8) and the indicator (6.4 or 6.5). Stir gently with the stirrer (7.1) titrate with the EDTA solution (6.2) (see 11.2, 11.3 and 11.4). Take v_1 as the volume, in millilitres, of the EDTA solution (6.2).

10 Calculation and expression of the result

For the EEC fertilizers to which the method is applicable (5 g of fertilizer in 500 ml of extract), calculate the MgO content as a mass fraction, w_{MgO} , in percent of the fertilizer according to Formula (1).

$$w_{\text{MgO}} = \frac{(v_2 - v_1) \times T}{M} \quad (1)$$

For the EEC fertilizers to which the method is applicable (5 g of fertilizer in 500 ml of extract), calculate the Mg content as a mass fraction, w_{Mg} , in percent of the fertilizer according to Formula (2).

$$w_{\text{Mg}} = \frac{(v_2 - v_1) \times T'}{M} \quad (2)$$

where

v_1 is the volume of the EDTA solution (6.2), in millilitres, used for the titration in the presence of calcein or calcon carbonic acid;

v_2 is the volume of the EDTA solution (6.2), in millilitres, used for the titration in the presence of Eriochrome black-T;

M is the mass of the sample present in the aliquot part taken, in grams;

T is $0,2016 \times \text{mol/l}$ of the EDTA solution (6.2) 0,05;

T' is $0,1216 \times \text{mol/l}$ of the EDTA solution (6.2) 0,05.

11 Remarks

11.1 The stoichiometric EDTA-metal ratio in the complexometric analyses is always 1:1 whatever the valency of the metal and in spite of the fact that EDTA is quadrivalent. The EDTA titration solution and the standard solutions will therefore be molar and not normal.