



# SLOVENSKI STANDARD

## SIST ISO 20666:2010

01-september-2010

---

**Kakovost vode - Določevanje kronične strupenosti z *Brachionus calyciflorus* v 48 urah**

Water quality - Determination of the chronic toxicity to *Brachionus calyciflorus* in 48 h

Qualité de l'eau - Détermination de la toxicité chronique vis-à-vis de *Brachionus calyciflorus* en 48 h

**STANDARD PREVIEW**  
**(standards.iteh.ai)**

**Ta slovenski standard je istoveten z: ISO 20666:2008**

SIST ISO 20666:2010  
<https://standards.iteh.ai/catalog/standards/sist/281d51a2-a7e4-49f1-89ac-9499d7df135e/sist-iso-20666-2010>

---

**ICS:**

13.060.70	Preiskava bioloških lastnosti vode	Examination of biological properties of water
-----------	------------------------------------	---

**SIST ISO 20666:2010**

**en,fr**

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

SIST ISO 20666:2010

<https://standards.iteh.ai/catalog/standards/sist/28fd3fa2-a7e4-49f1-89ac-9499d7df135e/sist-iso-20666-2010>

# INTERNATIONAL STANDARD

**ISO**  
**20666**

First edition  
2008-12-15

---

---

## Water quality — Determination of the chronic toxicity to *Brachionus* *calyciflorus* in 48 h

*Qualité de l'eau — Détermination de la toxicité chronique vis-à-vis de  
Brachionus calyciflorus en 48 h*

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

[SIST ISO 20666:2010](https://standards.iteh.ai/catalog/standards/sist/28fd3fa2-a7e4-49f1-89ac-9499d7dfl35e/sist-iso-20666-2010)

[https://standards.iteh.ai/catalog/standards/sist/28fd3fa2-a7e4-49f1-89ac-  
9499d7dfl35e/sist-iso-20666-2010](https://standards.iteh.ai/catalog/standards/sist/28fd3fa2-a7e4-49f1-89ac-9499d7dfl35e/sist-iso-20666-2010)



Reference number  
ISO 20666:2008(E)

© ISO 2008

**PDF disclaimer**

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

## iTeh STANDARD PREVIEW (standards.iteh.ai)

SIST ISO 20666:2010

<https://standards.iteh.ai/catalog/standards/sist/28fd3fa2-a7e4-49f1-89ac-9499d7df135e/sist-iso-20666-2010>

**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2008

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Case postale 56 • CH-1211 Geneva 20  
Tel. + 41 22 749 01 11  
Fax + 41 22 749 09 47  
E-mail [copyright@iso.org](mailto:copyright@iso.org)  
Web [www.iso.org](http://www.iso.org)

Published in Switzerland

## Contents

Page

Foreword.....	iv
Introduction .....	v
1 Scope .....	1
2 Normative references .....	1
3 Terms and definitions.....	2
4 Principle .....	2
5 Test environment.....	3
6 Reagents, test organisms and media .....	3
7 Apparatus .....	4
8 Treatment and preparation of samples .....	5
9 Procedure .....	6
10 Expression of results .....	9
11 Validity criteria .....	10
12 Test report .....	10
Annex A (informative) Preparation of the LC OLIGO medium .....	12
Annex B (informative) Precision data.....	14
Bibliography .....	15

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 20666 was prepared by Technical Committee ISO/TC 147, *Water quality*, Subcommittee SC 5, *Biological methods*.

## iTeh STANDARD PREVIEW (standards.iteh.ai)

[SIST ISO 20666:2010](https://standards.iteh.ai/catalog/standards/sist/28fd3fa2-a7e4-49f1-89ac-9499d7df135e/sist-iso-20666-2010)

<https://standards.iteh.ai/catalog/standards/sist/28fd3fa2-a7e4-49f1-89ac-9499d7df135e/sist-iso-20666-2010>

## Introduction

The evaluation of harmful effects on water quality has for several years involved the performance of biological tests. Rotifera, and especially the species *Brachionus calyciflorus*, are of interest from the ecotoxicological standpoint because they offer the advantage of breeding by parthenogenesis and of possessing a very short generation time: a single mother maintained under favourable conditions over 48 h reproduces several times. *Brachionus calyciflorus* is an organism of the zooplankton, which lives in fresh water. These animals are primary consumers and serve as prey for a large number of fish larvae and invertebrates.

The test specified in this International Standard is carried out over 48 h and therefore involves at least three reproductions from a single parent organism (see Reference [11]).

## iTeh STANDARD PREVIEW (standards.iteh.ai)

[SIST ISO 20666:2010](https://standards.iteh.ai/catalog/standards/sist/28fd3fa2-a7e4-49f1-89ac-9499d7df135e/sist-iso-20666-2010)

<https://standards.iteh.ai/catalog/standards/sist/28fd3fa2-a7e4-49f1-89ac-9499d7df135e/sist-iso-20666-2010>

## **iTeh STANDARD PREVIEW** **(standards.iteh.ai)**

SIST ISO 20666:2010

<https://standards.iteh.ai/catalog/standards/sist/28fd3fa2-a7e4-49f1-89ac-9499d7df135e/sist-iso-20666-2010>



# Water quality — Determination of the chronic toxicity to *Brachionus calyciflorus* in 48 h

**WARNING** — Persons using this International Standard should be familiar with normal laboratory practice. This International Standard does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices and to ensure compliance with any national regulatory conditions.

**IMPORTANT** — It is absolutely essential that tests conducted according to this International Standard be carried out by suitably trained staff.

## 1 Scope

This International Standard specifies a method for the determination of the chronic toxicity to rotifer *Brachionus calyciflorus*, based on population growth inhibition in 48 h.

The method is applicable to:

- a) chemical substances which are soluble or which can be maintained as stable suspensions or dispersions under the conditions of the test;
- b) industrial or sewage effluents, treated or untreated, if appropriate after decantation, filtration or centrifugation;
- c) fresh waters;
- d) aqueous extracts.

This International Standard is not applicable to the testing of unstable chemicals (hydrolysing, absorbing, etc.) in water unless exposure concentration is measured, nor to the testing of aquatic samples from the estuarine or marine environment.

## 2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 5667-16:1998, *Water quality — Sampling — Part 16: Guidance on biotesting of samples*

ISO 5814, *Water quality — Determination of dissolved oxygen — Electrochemical probe method*

ISO 10523, *Water quality — Determination of pH*

## ISO 20666:2008(E)

### 3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

#### 3.1

##### **control batch**

series of replicates containing **control solution** (3.2)

[ISO 20665:2008]

NOTE In this International Standard, eight replicates constitute the control batch.

#### 3.2

##### **control solution**

mixture of test medium and of food without sample under test

[ISO 20665:2008]

#### 3.3

##### **effective concentration producing $x$ % population growth inhibition**

$EC_x$

estimated concentration of the sample giving rise to  $x$  % **population growth inhibition** (3.4) with respect to the **control batch** (3.1)

#### 3.4

##### **population growth inhibition**

comparison of the total number of females (offspring and mothers) at the end of the test between the **control batch** (3.1) and the **test batch** (3.5)

#### 3.5

##### **test batch**

series of replicates filled with the same **test solution** (3.6)

[ISO 20665:2008]

NOTE In this International Standard, eight replicates constitute a test batch.

#### 3.6

##### **test solution**

mixture of test medium, of food and of sample under test

[ISO 20665:2008]

### 4 Principle

Female *Brachionus calyciflorus*, less than 2 h old at the beginning of the test, are exposed individually to a range of concentrations of the sample under test for a period of 48 h. The test focuses on the population growth of planktonic rotifers by parthenogenetic reproduction. At the end of the test, the number of female rotifers is determined and, by comparison with the control, the population growth inhibition percentages are determined for each concentration.

The data obtained allow, using a regression model, the calculation of the concentration which gives rise to  $x$  % population growth inhibition,  $EC_x$ , e.g.  $EC_{10}$ ,  $EC_{20}$  or  $EC_{50}$ .

## 5 Test environment

Carry out the test in the dark, in a thermostatically controlled room or chamber so as to obtain a temperature of  $(25 \pm 1) ^\circ\text{C}$  in the test containers.

Maintain the atmosphere free from toxic dusts or vapours. This is checked by producing control solutions.

## 6 Reagents, test organisms and media

Use only reagents of recognised analytical grade, unless otherwise specified.

### 6.1 Test organisms

Females of the species *Brachionus calyciflorus* (Monogonota, Rotifera) are obtained from a laboratory culture (see References [3], [12], [13]) or born from commercially available cysts<sup>1)</sup>. Sensitivity of the test for organisms should be performed with copper sulfate pentahydrate or sodium pentachlorophenolate (NaPCP) (see 6.5).

If cysts are used, employ first generation *Brachionus calyciflorus*, obtained by hatching of cysts under the following conditions.

Transfer the cysts to a container containing the test medium (6.3), e.g. 15 mg of cysts in approximately 10 ml of test medium. Incubate the container at  $(25 \pm 1) ^\circ\text{C}$  for 18 h to 24 h, under continuous lighting of intensity 1 000 lx to 4 000 lx (7.7).

A food supply is not necessary for the hatching of the cysts. A better multiplication rate of the rotifers is, however, achieved by adding, just after the emergence of the first neonates, algae in identical quantity to that indicated in 6.4. Alternatively, 100 µg of the inert food ROTIRICH<sup>1)</sup> can be added as pre-feeding supplement (Reference [18]).

The animals used for the test shall be less than 2 h old; the hatching should therefore be supervised as from 17 h of incubation, then every half hour.

The test is started when the number of young rotifers is considered sufficient to perform a complete test.

**EXAMPLE** For a test with five concentrations and one control (i.e. 48 rotifers), this condition is generally fulfilled about 1 h after the first hatching has been observed. Hatching time is quite stable within one laboratory, allowing hatching to be planned in advance to provide sufficient offspring during working hours.

**6.2 Pure water**, having a conductivity below 10 µS/cm<sup>2)</sup>.

**6.3 Test medium**, prepared by dissolving the following mineral substances in 1 l of pure water (6.2):

NaHCO <sub>3</sub>	96 mg
CaSO <sub>4</sub> ·2H <sub>2</sub> O	60 mg
MgSO <sub>4</sub>	60 mg
KCl	4 mg

This test medium corresponds to a synthetic water, of moderate hardness, i.e. 80 mg CaCO<sub>3</sub> to 100 mg CaCO<sub>3</sub> per litre (see Reference [14]). Thus prepared, the medium has a pH of  $7,6 \pm 0,3$ .

Store this solution in the dark at ambient temperature and use within 7 d of preparation.

1) Dehydrated rotifer cysts and ROTIRICH are examples of suitable products available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of these products.

2) 1 mS/m.