
**Cheese, cheese rind and processed
cheese — Determination of natamycin
content —**

Part 1:
**Molecular absorption spectrometric
method for cheese rind**

iTeh STANDARD PREVIEW

(standards.iteh.ai)

*Fromage, croûte de fromage et fromages fondus — Détermination de la
teneur en natamycine —*

*Partie 1: Méthode par spectrométrie d'absorption moléculaire pour
croûte de fromage*

<https://standards.iteh.ai/catalog/standards/sist/04c2d771-12a3-4fca-b141-a50b599a9e2e/iso-9233-1-2007>



Reference numbers
ISO 9233-1:2007(E)
IDF 140-1:2007(E)

PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. Neither the ISO Central Secretariat nor the IDF accepts any liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies and IDF national committees. In the unlikely event that a problem relating to it is found, please inform the ISO Central Secretariat at the address given below.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[ISO 9233-1:2007](https://standards.iteh.ai/catalog/standards/sist/04c2d771-12a3-4fca-b141-a50b599a9e2e/iso-9233-1-2007)

<https://standards.iteh.ai/catalog/standards/sist/04c2d771-12a3-4fca-b141-a50b599a9e2e/iso-9233-1-2007>



COPYRIGHT PROTECTED DOCUMENT

© ISO and IDF 2007

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO or IDF at the respective address below.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

International Dairy Federation
Diamant Building • Boulevard Auguste Reyers 80 • B-1030 Brussels
Tel. + 32 2 733 98 88
Fax + 32 2 733 04 13
E-mail info@fil-idf.org
Web www.fil-idf.org

Published in Switzerland

Contents

Page

Foreword.....	iv
Foreword.....	v
1 Scope	1
2 Terms and definitions.....	1
3 Principle.....	1
4 Reagents	2
5 Apparatus	2
6 Sampling.....	3
7 Preparation of test sample.....	3
7.1 Cheese rind	3
7.2 Cheese interior.....	3
8 Procedure	4
8.1 Test portion	4
8.2 Preparation of test solution.....	4
8.3 Determination.....	5
9 Calculation and expression of results.....	6
9.1 Calculation of natamycin mass fraction.....	6
9.2 Calculation of absorbance.....	6
9.3 Calculation of surface-area-related natamycin mass.....	7
9.4 Correction of results.....	7
9.5 Expression of results	7
10 Precision.....	7
10.1 Interlaboratory tests	7
10.2 Repeatability.....	8
10.3 Reproducibility.....	8
11 Test report	8
Annex A (informative) Examples	9
Annex B (informative) Results of interlaboratory trial.....	13
Bibliography	14

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization. International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 9233-1|IDF 140-1 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 5, *Milk and milk products* and the International Dairy Federation (IDF) and is being published jointly by ISO and IDF.

iTeh STANDARD PREVIEW

This first edition of ISO 9233-1|IDF 140-1, together with ISO 9233-2|IDF 140-2, cancel and replace the first edition of ISO 9233:1991, which has been technically revised.

ISO 9233|IDF 140 consists of the following parts, under the general title *Cheese, cheese rind and processed cheese — Determination of natamycin content*:

- Part 1: *Molecular absorption spectrometric method for cheese rind*
- Part 2: *High-performance liquid chromatographic method for cheese, cheese rind and processed cheese*

Foreword

IDF (the International Dairy Federation) is a worldwide federation of the dairy sector with a National Committee in every member country. Every National Committee has the right to be represented on the IDF Standing Committees carrying out the technical work. IDF collaborates with ISO in the development of standard methods of analysis and sampling for milk and milk products. Draft International Standards adopted by the Action Teams and Standing Committees are circulated to the National Committees for voting. Publication as an International Standard requires approval by at least 50% of IDF National Committees casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. IDF shall not be held responsible for identifying any or all such patent rights.

ISO 9233-1|IDF 140-1 was prepared by the International Dairy Federation (IDF) and Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 5, *Milk and milk products* and is being published jointly by IDF and ISO.

All work was carried out by the Joint ISO-IDF Action Team on *Selected food additives and vitamins* of the Standing Committee on *Analytical methods for additives and contaminants* under the aegis of its project leader, Mr. M. Carl (DE).

This first edition of ISO 9233-1|IDF 140-1, together with ISO 9233-2|IDF 140-2, cancel and replace the first edition of IDF 140A:1992, which has been technically revised.

ISO 9233|IDF 140 consists of the following parts, under the general title *Cheese, cheese rind and processed cheese — Determination of natamycin content*.

- *Part 1: Molecular absorption spectrometric method for cheese rind*
- *Part 2: High-performance liquid chromatographic method for cheese, cheese rind and processed cheese*

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 9233-1:2007

<https://standards.iteh.ai/catalog/standards/sist/04c2d771-12a3-4fca-b141-a50b599a9e2e/iso-9233-1-2007>

Cheese, cheese rind and processed cheese — Determination of natamycin content —

Part 1: Molecular absorption spectrometric method for cheese rind

1 Scope

This part of ISO 9233|IDF 140 specifies a method for the determination in cheese rind of natamycin mass fraction of above 0,5 mg/kg and surface-area-related natamycin mass of above 0,03 mg/dm².

NOTE It is possible that the method may be suitable for detecting migration of natamycin into the cheese.

2 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

2.1

natamycin content <https://standards.iteh.ai/catalog/standards/sist/04c2d771-12a3-4fca-b141-ed1579e9c1c1/iso-9233-1-2007>
mass fraction of substances determined by the procedure specified in this part of ISO 9233|IDF 140

NOTE The natamycin content is expressed in milligrams per kilogram.

2.2

surface-area-related natamycin mass in cheese rind

surface-area-related mass of substances determined by the procedure specified in this part of ISO 9233|IDF 140

NOTE The surface-area-related natamycin mass is expressed in milligrams of natamycin per square decimetre of cheese rind.

2.3

cheese rind

outer layer of the cheese of thickness 5 mm, excluding the coating layer, if present.

3 Principle

A known quantity of sample is extracted with methanol. The extract is diluted with water followed by cooling to between -15 °C and -20 °C to precipitate most of the fat, followed by filtration. The natamycin content or surface-area-related natamycin mass is determined in the filtrate (after concentration, if necessary) by molecular absorption spectrometry.

4 Reagents

Use only reagents of recognized analytical grade, unless otherwise specified, and only distilled or demineralized water or water of equivalent purity.

4.1 Methanol (CH₃OH).

4.2 Methanol, aqueous solution.

Mix 2 volumes of methanol (4.1) with 1 volume of water.

4.3 Natamycin standard solutions.

4.3.1 Natamycin standard stock solution, of concentration 500 mg/l.

Immediately before use, dissolve in methanol (4.1) a quantity of a natamycin preparation of known natamycin content, corresponding to 50 mg of pure natamycin (C₃₃H₄₇NO₁₃), in a 100 ml one-mark volumetric flask (5.1). Make up to the mark with water and mix.

4.3.2 Natamycin standard working solution, of concentration 5 mg/l.

Pipette 5,0 ml of natamycin standard stock solution (4.3.1) into a 50 ml one-mark volumetric flask (5.1). Dilute to the mark with aqueous methanol (4.2) and mix.

Pipette 5,0 ml of the thus diluted solution into another 50 ml one-mark volumetric flask (5.1). Dilute to the mark with aqueous methanol (4.2) and mix. The concentration of this natamycin standard working solution is 5 µg/ml.

This concentration shall be close to that of the test solution measured in 8.3.3. Adjust the standard working dilution by pipetting and diluting another quantity, if required.

5 Apparatus

Usual laboratory equipment and, in particular, the following.

5.1 One-mark volumetric flasks, of capacities 50 ml and 100 ml.

5.2 Slicer or similar apparatus, capable of cutting cheese portions of thickness 5 mm and of width about 30 mm (Figure A.1 shows an example).

5.3 Fine slicer, capable of cutting thin cheese slices of maximum thickness 1 mm (Figure A.2 shows an example).

5.4 Grinder or blender.

5.5 Sharp knife, capable of cutting cheese slices into small pieces.

5.6 Magnetic stirrer or shaking machine.

5.7 Conical flasks, of capacities 100 ml and 200 ml, made of coloured glass and fitted with ground-glass stoppers.

5.8 Syringes, disposable, of capacity 10 ml.

5.9 Membrane microfilters, of pore size 0,20 µm and 0,45 µm, resistant to attack by alcoholic solutions.

5.10 Folded paper filters, fast speed, of diameter 150 mm [e.g. S and S, No. 595 1/2¹].

5.11 Funnel, of diameter approximately 70 mm.

5.12 Freezer, capable of freezing at a temperature of between -15 °C and -20 °C.

5.13 Extraction cartridges, to concentrate the filtered extract, if necessary [e.g. Sep-pack C18¹) or Waters No. 51910¹].

5.14 Spectrometer, suitable for recording an ultraviolet (UV) spectrum between 300 nm and 340 nm, equipped with cells of optical pathlength 10 mm and a recorder.

5.15 Sample jar, of suitable capacity.

6 Sampling

A representative sample should have been sent to the laboratory. It should not have been damaged or changed during transport or storage.

Sampling is not part of the method specified in this part of ISO 9233|IDF 140. A recommended sampling method is given in ISO 707|IDF 50.

The laboratory sample shall be a whole cheese, or a segment of a cheese representative of the whole.

7 Preparation of test sample

7.1 Cheese rind <https://standards.iteh.ai/catalog/standards/sist/04c2d771-12a3-4fca-b141-a50b599a9e2e/iso-9233-1-2007>
ISO 9233-1:2007

If necessary, cut the test sample into sectors or smaller portions so that the width of the cheese rind is not more than about 30 mm. Using the slicer (5.2), remove the whole rind from all obtained sectors or portions by slicing off a maximum thickness of 5 mm.

From the rind obtained, cut, with a sharp knife (5.5), a rectangular piece of area between 2 dm² and 4 dm². Determine its surface area, in square decimetres, and its mass, in kilograms.

Grind (5.4) carefully the whole rind, including the weighed and measured piece, and mix thoroughly. Immediately transfer a quantity of the sample thus prepared to a sample jar (5.15).

After preparing each test sample, clean all tools that have been in contact with the sample with hot water and then with methanol (4.1). Dry all tools thoroughly, e.g. by using a stream of compressed air.

7.2 Cheese interior

After removing the rind (7.1), use the fine slicer (5.3) to remove a slice of maximum thickness 1 mm from the whole of the outer section of the test sample.

Cut all cheese slices into small pieces of about 50 mm² and mix thoroughly. Immediately transfer a quantity of the sample thus prepared to a sample jar (5.15).

1) Example of a suitable product available commercially. This information is given for the convenience of users of this part of ISO 9233|IDF 140 and does not constitute an endorsement by ISO or by IDF of this product.

After preparing each test sample, clean all tools that have been in contact with the test sample with hot water and then with methanol (4.1). Dry all tools thoroughly, e.g. by using a stream of compressed air.

8 Procedure

8.1 Test portion

8.1.1 Cheese rind

Weigh, to the nearest 10 mg, approximately 10,00 g of test sample (7.1) into a 200 ml conical flask (5.7).

8.1.2 Cheese interior

Weigh, to the nearest 10 mg, approximately 5,00 g of test sample (7.2) into a 100 ml conical flask (5.7).

8.2 Preparation of test solution

8.2.1 Cheese rind

8.2.1.1 Initial steps

Add 100 ml of methanol (4.1) to the test portion in the conical flask (8.1.1). Stir the contents of the conical flask for 90 min with a magnetic stirrer (5.6) or shake for 90 min in a shaking machine (5.6).

Add 50 ml water. Immediately place the conical flask in the freezer (5.12) for about 60 min.

8.2.1.2 Filtration

<https://standards.iteh.ai/catalog/standards/sist/04c2d771-12a3-4fca-b141-a50b599a9e2e/iso-9233-1-2007>

Filter the cold extract through a folded filter paper (5.10) while discarding the first 5 ml of filtrate. The filtration should be carried out while the suspension is still cold to avoid dissolution of the fat and consequently turbid filtrates.

Bring the filtrate to room temperature. Take a portion of the filtrate in a syringe (5.8). Filter through a membrane microfilter of pore size 0,45 µm (5.9) and then through a membrane microfilter of pore size 0,20 µm (5.9).

The minimum amount of test solution (filtrate) required is 3 ml for direct measurement (8.3.3) and 25 ml or 50 ml for measurement at 5 or 10 times concentration (8.3.4), respectively.

8.2.2 Cheese interior

8.2.2.1 Initial steps

Use a measuring cylinder to add 50 ml of methanol (4.1) to the test portion in the conical flask (8.1.2). Stir the contents of the conical flask for 90 min with a magnetic stirrer (5.6) or shake for 90 min in a shaking machine (5.6).

Use a measuring cylinder to add 25 ml of water. Immediately place the conical flask in the freezer (5.12) for about 60 min.

8.2.2.2 Filtration

Filter the solution as described in 8.2.1.2.

8.3 Determination

8.3.1 Determination and detection limits

The laboratory applying the method shall establish the limits of detection and determination under its own instrumental conditions using recognized calculation methods to verify that natamycin can be determined down to levels of 0,5 mg/kg and 0,03 mg/dm².

8.3.2 UV absorbance of natamycin standard working solution

Record the spectrum of the natamycin standard working solution (4.3.2) in the range 300 nm to 340 nm. Measure the absorbance of natamycin at its maximum at about 317 nm, at its minimum at about 311 nm, and at 329 nm exactly. Use aqueous methanol (4.2) as blank.

An example of a natamycin standard working solution spectrum is shown in Figure A.3.

Because natamycin is unstable in aqueous methanol, carry out the measurement as rapidly as possible.

8.3.3 Test solution

Record the spectra of the test solutions (8.2.1.2 or 8.3.4.2 and 8.2.2.2) with aqueous methanol (4.2) as blank in the range 300 nm to 340 nm. Additionally, record in the same range the spectrum of the cheese rind test solution (8.2.1.2 or 8.3.4.2) with the cheese interior test solution (8.2.2.2) as blank.

Record the absorbance of the cheese rind test solution (8.2.1.2 or 8.3.4.2) with the cheese interior test solution (8.2.2.2) as blank at its maximum absorbance at about 317 nm, at its minimum at about 311 nm, and at 329 nm exactly. Examples of test solution spectra are shown in Figure A.4.

If the natamycin content of the test sample (7.1) is so low that no detection is possible or almost impossible (signal-to-noise ratio less than 3), but its determination is nevertheless required, proceed in accordance with 8.3.4.

NOTE The presence of spices, particularly pepper, in the cheese can interfere with the result, which might also be shown by an obvious distortion of the absorbance curve. Examples of spectra of various test solutions of cheeses are given in Figure A.4.

8.3.4 Low natamycin content

8.3.4.1 Concentration

Decide whether a concentration of about 5 times or about 10 times is desired. Base that decision on the result obtained in 8.3.3 and on the required limit of determination.

Then pipette 25 ml or 50 ml (for concentration times 5 or concentration times 10, respectively) of test solution (8.2.1.2) into a beaker. Add, depending on the concentration desired, 50 ml or 100 ml of water, respectively, and mix.

Activate an extraction cartridge (5.13) by using 3 ml to 5 ml of methanol (4.1). Then wash with 10 ml of water.

Pass the diluted test solution through the cartridge at a speed of 3 ml/min to 5 ml/min with the aid of a syringe (5.8). Rinse the cartridge with 10 ml of water with the aid of a syringe (5.8). Elute the natamycin with 3 ml of methanol (4.1) with the aid of a syringe (5.8).