

INTERNATIONAL
STANDARD

ISO
11866-2

IDF
170-2

Second edition
2005-12-01

**Milk and milk products — Enumeration
of presumptive *Escherichia coli* —**

Part 2:

**Colony-count technique at 44 °C using
membranes**

iTeh STANDARD PREVIEW

(standards.iteh.ai)

Lait et produits laitiers — Dénombrement d'Escherichia coli présumés —

*Partie 2: Technique par comptage des colonies obtenues sur
membranes à 44 °C*

[ISO 11866-2:2005](#)

<https://standards.iteh.ai/catalog/standards/sist/a31e81dd-2619-4281-8125-16b46117ca2b/iso-11866-2-2005>



Reference numbers
ISO 11866-2:2005(E)
IDF 170-2:2005(E)

© ISO and IDF 2005

PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. Neither the ISO Central Secretariat nor the IDF accepts any liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies and IDF national committees. In the unlikely event that a problem relating to it is found, please inform the ISO Central Secretariat at the address given below.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[ISO 11866-2:2005](https://standards.iteh.ai/catalog/standards/sist/a31e81dd-2619-4281-8125-16b46117ca2b/iso-11866-2-2005)

<https://standards.iteh.ai/catalog/standards/sist/a31e81dd-2619-4281-8125-16b46117ca2b/iso-11866-2-2005>

© ISO and IDF 2005

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO or IDF at the respective address below.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

International Dairy Federation
Diamant Building • Boulevard Auguste Reyers 80 • B-1030 Brussels
Tel. + 32 2 733 98 88
Fax + 32 2 733 04 13
E-mail info@fil-idf.org
Web www.fil-idf.org

Published in Switzerland

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 11866-2|IDF 170-2 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 5, *Milk and milk products*, and the International Dairy Federation (IDF). It is being published jointly by ISO and IDF.

This edition of ISO 11866-2|IDF 170-2 cancels and replaces ISO 11866-3:1997, of which it constitutes a minor revision.

ISO 11866-1:1997 has been cancelled and replaced by ISO 7251:2005, *Microbiology of food and animal feeding stuffs — Horizontal method for the detection and enumeration of presumptive Escherichia coli — Most probable number technique*.

ISO 11866|IDF 170 consists of the following parts, under the general title *Milk and milk products — Enumeration of presumptive Escherichia coli*:

- *Part 1: Most probable number technique using 4-methylumbelliferyl- β -D-glucuronide (MUG)*
- *Part 2: Colony-count technique at 44 °C using membranes*

Foreword

IDF (the International Dairy Federation) is a worldwide federation of the dairy sector with a National Committee in every member country. Every National Committee has the right to be represented on the IDF Standing Committees carrying out the technical work. IDF collaborates with ISO in the development of standard methods of analysis and sampling for milk and milk products.

Draft International Standards adopted by the Action Teams and Standing Committees are circulated to the National Committees for voting. Publication as an International Standard requires approval by at least 50 % of the IDF National Committees casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. IDF shall not be held responsible for identifying any or all such patent rights.

ISO 11866-2|IDF 170-2 was prepared by the International Dairy Federation (IDF) and Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 5, *Milk and milk products*. It is being published jointly by IDF and ISO.

All work was carried out by the Joint ISO/IDF/AOAC Group of Experts on *Pathogenic contaminants* (E102), under the aegis of its chairman, Mrs R. Lodi (IT).

This edition of ISO 11866-2|IDF 170-2 cancels and replaces the former part 3 of IDF 170A:1999, while the former part 1 has been replaced by ISO 7251:2005

ITeH STANDARD PREVIEW
(standards.iteh.ai)
<https://standards.iteh.ai/catalog/standards/sist/a31e81dd-2619-4281-8125-16b46117ca2b/iso-11866-2-2005>

Milk and milk products — Enumeration of presumptive *Escherichia coli* —

Part 2: Colony-count technique at 44 °C using membranes

1 Scope

This part of ISO 11866|IDF 170 specifies a method for the enumeration of presumptive *Escherichia coli* by means of a colony-count technique at 44 °C.

The method is applicable to

- milk, liquid milk products,
- dried milk, dried sweet whey, dried buttermilk, lactose,
- acid casein, lactic casein and rennet casein,
- caseinate and dried acid whey,
- cheese and processed cheese,
- butter,
- frozen milk products (including edible ices), and
- custard, desserts and cream.

The method specified in this part of ISO 11866|IDF 170 is the preferred method for samples in which comparatively large numbers of presumptive *Escherichia coli* (more than 100 per gram or 10 per millilitre) are suspected.

CAUTION — Some pathogenic strains of *Escherichia coli* do not grow at 44 °C.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 7218, *Microbiology of food and animal feeding stuffs — General requirements and guidance for microbiological examinations*

ISO 8261|IDF 122, *Milk and milk products — General guidance for the preparation of test samples, initial suspensions and decimal dilutions for microbiological examination*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1 presumptive *Escherichia coli*
bacteria which at 44 °C form indole-positive (pink) colonies on cellulose acetate membranes overlaid on tryptone-bile agar, under the conditions specified in this part of ISO 11866|IDF 170

4 Principle

4.1 Resuscitation

A specified quantity of the test sample or initial suspension is inoculated onto cellulose acetate membranes overlaid on mineral-modified glutamate agar, then they are incubated at 37 °C for 4 h.

NOTE This procedure enables the presumptive *Escherichia coli* damaged by storage under frozen, dried or chill conditions, or damaged by heat or chemical processes, to be resuscitated. It also permits the diffusion of high concentrations of any fermentable carbohydrate present in the test sample which would otherwise interfere with indole production during the subsequent isolation stage.

4.2 Isolation

The membranes from the resuscitation stage on the mineral-modified glutamate agar are transferred to tryptone-bile agar. They are incubated at 44 °C for 18 h to 24 h.

4.3 Detection

The presence of presumptive *Escherichia coli* on the membrane is demonstrated by the production of indole by each colony.

4.4 Calculation

The number of colony-forming units (CFU) of presumptive *Escherichia coli* per gram or per millilitre of sample is calculated from the number of indole-positive colonies obtained on membranes at dilution levels chosen so as to give a significant result.

5 Dilution fluid, culture media and reagent

5.1 General

For current laboratory practice, see ISO 7218 and ISO 8261.

If the prepared culture media and reagents are not used immediately, they shall, unless otherwise stated, be stored in the dark at a temperature between 0 °C and +5 °C for no longer than 1 month, under conditions which do not produce any change in their composition.

5.2 Dilution fluid

See ISO 8261|IDF 122.

5.3 Culture media and reagent

5.3.1 Resuscitation medium: Mineral-modified glutamate agar

5.3.1.1 Composition

Sodium glutamate	6,35 g
Lactose	10,0 g
Sodium formate	0,25 g
L(-)Cystine	0,02 g
L(-)Aspartic acid	0,02 g
L(+)-Arginine	0,024 g
Thiamine	0,001 g
Nicotinic acid	0,001 g
Pantothenic acid	0,001 g
Magnesium sulfate heptahydrate (MgSO ₄ ·7H ₂ O)	0,100 g
Ammonium iron(III) citrate ^a	0,010 g
Calcium chloride dihydrate (CaCl ₂ ·2H ₂ O)	0,010 g
Dipotassium hydrogen phosphate (K ₂ HPO ₄)	0,90 g
Ammonium chloride	2,5 g
Agar	12 g to 18 g ^b
Water	1 000 ml
^a Iron content of at least 15 % (mass fraction).	
^b Depending on the gel strength of the agar.	

5.3.1.2 Preparation

Dissolve the ammonium chloride in the water. Add the other components and heat to boiling.

Adjust the pH, if necessary, so that after sterilization it is 6,7 at 25 °C.

Transfer 100 ml volumes of the medium to suitable containers.

Sterilize in the autoclave (6.1) set at 115 °C for 10 min.

5.3.1.3 Preparation of agar plates

Pour into sterile Petri dishes (6.12), 12 ml to 15 ml of the medium cooled to approximately 45 °C, and allow it to solidify. The plates may be stored at 0 °C to +5 °C for up to 4 days.

Immediately before use, dry the plates, preferably with the lids removed and the agar surfaces facing downwards, in the drying cabinet or the oven (6.3) set at 50 °C for 30 min or until the droplets have disappeared from the surface of the medium.

The agar should be dry enough not to allow excess moisture to appear within 15 min of spreading the inoculum (1 ml).

5.3.2 Selective medium: Tryptone-bile agar

5.3.2.1 Composition

Tryptone	20,0 g
Bile salts (refined)	1,5 g
Agar	12 g to 18 g ^a
Water	1 000 ml
^a Depending on the gel strength of the agar.	

5.3.2.2 Preparation

Dissolve the components in the water and heat to boiling.

Adjust the pH, if necessary, so that after sterilization it is 7,2 at 25 °C.

Transfer aliquots of up to 500 ml of the medium to suitable containers.

Sterilize the medium in the autoclave (6.1) set at 121 °C for 15 min.

5.3.2.3 Preparation of agar plates

Pour into sterile Petri dishes (6.12), 12 ml to 15 ml of the medium cooled to approximately 45 °C, and allow it to solidify. The plates may be stored at 0 °C to +5 °C for up to 4 days.

Immediately before use, dry the plates, preferably with the lids removed and the agar surfaces facing downwards, in the oven (6.3) set at 50 °C for 30 min or until the droplets have disappeared from the surface of the medium.

5.3.3 Indole detection reagent (Vracko and Sherris reagent)

5.3.3.1 Composition

4-Dimethylaminobenzaldehyde	5,0 g
Hydrochloric acid, $c(\text{HCl}) = 1 \text{ mol/l}$	100 ml

5.3.3.2 Preparation

Dissolve the 4-dimethylaminobenzaldehyde in the hydrochloric acid by heating, if necessary. The reagent may be stored in the dark at 0 °C to +5 °C for a maximum period of 3 months.

6 Apparatus and glassware

For general requirements, see ISO 7218 and ISO 8261 | IDF 122. Glassware shall be resistant to repeated sterilization.

Usual microbiological laboratory apparatus and, in particular, the following.

6.1 Autoclave, capable of operating at $115 \text{ °C} \pm 1 \text{ °C}$ and at $121 \text{ °C} \pm 1 \text{ °C}$.

For details, see ISO 7218.

- 6.2 Incubators**, capable of operating at $37\text{ °C} \pm 1\text{ °C}$ and at $44\text{ °C} \pm 0,5\text{ °C}$.
- 6.3 Drying cabinet or oven**, ventilated by convection, capable of operating at $50\text{ °C} \pm 1\text{ °C}$.
- 6.4 Refrigerator** (for storage of prepared media and reagent), capable of operating at 0 °C to 5 °C .
- 6.5 Cellulose acetate membranes**, $0,45\text{ }\mu\text{m}$ to $1,2\text{ }\mu\text{m}$ pore size and of 85 mm diameter.
- 6.6 Long-wave ultraviolet (UV) lamp**, of wavelength between 360 nm and 366 nm , fitted with a suitable filter to remove UV radiation below 310 nm .
- 6.7 Blunt-ended forceps**, sterile, of approximately 12 cm length.
- 6.8 pH-meter**, accurate to within $\pm 0,1\text{ pH}$ units at 25 °C .
- 6.9 Pipettes**, calibrated for bacteriological use, with 1 ml nominal capacity, graduated in divisions of $0,1\text{ ml}$ and with an outflow opening of 2 mm to 3 mm diameter.
- 6.10 Measuring cylinders**, for preparation of the media and reagent.
- 6.11 Bottles or flasks**, for sterilization and storage of culture media.
- 6.12 Petri dishes**, made of glass or plastic, of approximately 90 mm or approximately 100 mm diameter.
- 6.13 Spreaders**, made of glass or plastic, for example hockey sticks made from a glass rod of approximately $3,5\text{ mm}$ diameter and 20 cm length, bent at right angles about 3 cm from one end and with the cut ends made smooth by heating.

(standards.iteh.ai)

7 Sampling

ISO 11866-2:2005

<https://standards.iteh.ai/catalog/standards/sist/a31e81dd-2619-4281-8125-1041cc29b011/iso-11866-2:2005>

A representative sample should have been sent to the laboratory. It should not have been damaged or changed during transport or storage.

Sampling is not part of the method specified in this part of ISO 11866|IDF 170. A recommended sampling method is given in ISO 707|IDF 50.

8 Preparation of test sample

Prepare the test sample according to the method given in ISO 8261|IDF 122.

9 Procedure

NOTE If it is required to check whether the repeatability requirement is met (see Clause 11) carry out two single determinations in accordance with 9.1 to 9.5.

9.1 Test portion, initial suspension and further dilutions

Prepare the test portion, initial suspension (primary dilution) and further dilutions according to the method given in ISO 8261|IDF 122.