

Annulé

ISO

INTERNATIONAL ORGANIZATION FOR STANDARDIZATION

ISO RECOMMENDATION

R 455

iTeh STANDARD PREVIEW
ANALYSIS OF SOAP
(standards.iteh.ai)

DETERMINATION OF TOTAL CRUDE FATTY ACIDS

ISO/R 455:1965

<https://standards.iteh.ai/catalog/standards/sist/a3f9bbf5-3ce5-48db-a153-fca5c1aff1a/iso-r-455-1965>

1st EDITION

November 1965

COPYRIGHT RESERVED

The copyright of ISO Recommendations and ISO Standards belongs to ISO Member Bodies. Reproduction of these documents, in any country, may be authorized therefore only by the national standards organization of that country, being a member of ISO.

For each individual country the only valid standard is the national standard of that country.

Printed in Switzerland

Also issued in French and Russian. Copies to be obtained through the national standards organizations.

BRIEF HISTORY

The ISO Recommendation R 455, *Analysis of Soap. Determination of Total Crude Fatty Acids*, was drawn up by Technical Committee ISO/TC 91, *Surface Active Agents*, the Secretariat of which is held by the Association Française de Normalisation (AFNOR).

Work on this question by the Technical Committee began in 1961 and led, in 1962, to the adoption of a Draft ISO Recommendation.

In June 1963, this Draft ISO Recommendation (No. 583) was circulated to all the ISO Member Bodies for enquiry. It was approved, subject to a few modifications of an editorial nature, by the following Member Bodies:

Argentina	Hungary	Poland
Austria	Italy	Portugal
Canada	Japan	Romania
Chile	Korea, Rep. of	Spain
Colombia	Morocco	Switzerland
Czechoslovakia	Netherlands	United Kingdom
France	New Zealand	Yugoslavia
Germany	Norway	

One Member Body opposed the approval of the Draft: India.

The Draft ISO Recommendation was then submitted by correspondence to the ISO Council, which decided, in November 1965, to accept it as an ISO RECOMMENDATION.

ANALYSIS OF SOAP

DETERMINATION OF TOTAL CRUDE FATTY ACIDS

1. SCOPE

The purpose of this ISO Recommendation is to specify the method of determining total crude fatty acids in commercial soaps, excluding compounded products.

In the Annex, there is also described a method which should be used only for routine controls and for soaps in which the fatty acids are not contaminated with substances insoluble in water, derived from additives (silicates etc.).

2. TERMINOLOGY

Total crude fatty acids means the water-insoluble fatty material which is collected by decomposing the soap with a strong mineral acid under the operating conditions described. This term includes unsaponifiable matter, glycerides and any resinic acids contained in the soap, in addition to the so-called fatty acids.

3. PRINCIPLE

ISO/R 455:1965

The fatty acids are extracted with diethyl ether and titrated with a solution of sodium hydroxide in ethanol.

4. REAGENTS

4.1 Diethyl ether, pure.

4.2 Ethanol solution, 95 % by volume.

4.3 Acid solution:

either sulphuric acid, $d = 1.83$, diluted 1/5 ; *
or hydrochloric acid, $d = 1.19$, diluted 1/3. *

4.4 Sodium chloride solution, 10 g of sodium chloride dissolved in 100 ml of distilled water.

4.5 Sodium hydroxide ethanolic solution, analytical grade, recently standardized to approximately 0.5 N.

4.6 Methyl-orange solution, 0.2 g in 100 ml of distilled water.

4.7 Phenolphthalein solution, 1 g in 100 ml of the ethanol solution (4.2).

* Diluted p_1/p_2 means that p_1 parts by volume of the specified solution are diluted to give p_2 parts by volume of the final mixture.

5. APPARATUS

Ordinary laboratory equipment and in particular,

- (a) Porcelain or glass basins of about 250 ml.
- (b) Separating funnels, of about 500 ml.
- (c) Water-bath.
- (d) Oven regulated at 120°C.
- (e) Desiccator.
- (f) Analytical balance.

6. PROCEDURE

6.1 Test portion

Weigh 5 to 10 g of soap into the basin to an accuracy of 0.001 g.

6.2 Determination

Make a hot solution in 100 ml of distilled water. Pour this aqueous solution into a separating funnel, rinsing the basin with small quantities of distilled water.

Add a few drops of the methyl-orange (4.6) and then about 10 ml of the acid solution (4.3). The indicator should turn red. If it does not, add 2 ml more of the acid solution (4.3). Leave to cool to room temperature.

Add 100 ml of the diethyl ether (4.1). Shake the mixture vigorously for 5 min and allow to stand until the two phases are completely separated.

Draw off the acidified water into a second separating funnel. Make a second extract of this acid solution, mixing in the same way, with 50 ml of the diethyl ether (4.1).

Draw off the acidified water. Combine the ethereal solutions in the same separating funnel. Wash twice in succession, using 50 ml of the sodium chloride solution (4.4) each time (mixing for 1 min each time). Verify that the last wash is neutral to methyl-orange (4.6). If it is not, continue washing in the same way until the washing solution is neutral to methyl-orange (4.6).

After drawing off the last washing solution, filter the ethereal solution, if necessary, through a filter paper. Collect in a tared flask. Wash the filter with small portions of the diethyl ether (4.1). Distil off nearly all the diethyl ether by boiling gently.

Dissolve the residue in 20 ml of the ethanol solution (4.2). Neutralize the ethanolic solution of fatty acids with the ethanolic sodium hydroxide solution (4.5),* using 2 to 3 drops of phenolphthalein (4.7) as indicator. Note the volume used.**

Remove the ethanol by placing on a boiling water-bath. Heat the flask in the oven at 120°C until constant mass is reached, i.e., until the difference in weight, after drying in the oven for an additional 15 min, does not exceed 5 mg.*** Weigh the dry soap.

* Do not use ethanolic potassium hydroxide solution.

** If the fatty acid colour masks the end-point, this may be determined potentiometrically; this method will be standardized later.

*** Before weighing, leave the flask in the desiccator just long enough for it to reach ambient temperature.

7. EXPRESSION OF RESULTS

7.1 Method of calculation and formula

If E is the mass, in grammes, of the test portion,

m is the mass, in grammes, of dry soap,

n is the number of millilitres of ethanolic sodium hydroxide solution (4.5) used,

T is the exact normality of the ethanolic sodium hydroxide solution (4.5),

the percentage of total crude fatty acids in the soap is

$$\left[m - (n \times T \times 0.022) \right] \times \frac{100}{E}$$

Round the result to the nearest 0.1 per cent.

If M_0 is the nominal mass, in grammes, of the piece of soap,

M_1 is the mean mass, in grammes, of the piece of soap at the time of analysis,

the percentage of total crude fatty acids in the initial soap is

$$\left[m - (n \times T \times 0.022) \right] \times \frac{100}{E} \times \frac{M_1}{M_0}$$

iTeh STANDARD PREVIEW
(standards.iteh.ai)

7.2 Reproducibility

± 0.2 absolute.

[ISO/R 455:1965](https://standards.iteh.ai/catalog/standards/sist/a3f9bbf5-3ce5-48db-a153-fca5c1aff1a/iso-r-455-1965)

<https://standards.iteh.ai/catalog/standards/sist/a3f9bbf5-3ce5-48db-a153-fca5c1aff1a/iso-r-455-1965>

7.3 Notes

1. When analysis certificates refer only to "fatty acids", this always means "total crude fatty acids".
2. If resin is present, and if a test has been made for it, the result should preferably be expressed as "fatty and resin acids", but the term "fatty acids" by itself does not imply that the soap is free from resin.
3. If the determination of "pure fatty acids" is required, deduct from the percentage of "total crude fatty acids" the sum of the percentages of unsaponifiable matter, unsaponified matter and any resinic acids.

8. TEST REPORT

In addition to the results, the test report should state all the test conditions and details of procedure not mentioned in this ISO Recommendation, as well as any other factors which may have affected the results.

ANNEX A

WAX CAKE METHOD

A.1 PRINCIPLE

The soap is dissolved and the fatty acids are completely separated by acidification. They are then solidified with beeswax in the form of a cake, which is washed with boiling water, dried and weighed.

A.2 REAGENTS

A.2.1 Beeswax, paraffin wax or stearic acid (or mixture of these) described as "wax" in the text, free from impurities and previously dried by heating at approximately 120°C.

A.2.2 *Acid solution :*

either sulphuric acid, $d = 1.83$, diluted 1/5 ;
or hydrochloric acid, $d = 1.19$, diluted 1/3.

A.2.3 Methyl orange solution: 0.2 g in 100 ml of distilled water.

<https://standards.iteh.ai/catalog/standards/sist/a3f9bbf5-3ce5-48db-a153-fca5c1aff1a/iso-r-455-1965>

A.3 APPARATUS

Ordinary laboratory equipment, and in particular,

- (a) Porcelain or glass basin (C1), approximately 11 cm diameter.
- (b) Basin (C2), 8 to 9 cm diameter and 4 to 5 cm deep, preferably flat-bottomed and containing a small stirrer thermometer.
- (c) Water-bath.
- (d) Sand-bath.
- (e) Desiccator.
- (f) Analytical balance.

A.4 PROCEDURE

A.4.1 Test portion

Weigh approximately 10 g of soap with an accuracy of 0.01 g.

A.4.2 Determination

Place the test portion in the basin (C1) containing a stirrer. Add 100 ml of distilled water. Leave on a boiling water-bath until the soap has completely dissolved.

Add a few drops of the methyl-orange indicator (A.2.3) and then approximately 10 ml of the acid solution (A.2.2) until the indicator clearly changes colour. The fatty acids separate and come to the surface as a snow-like mass. Continue heating on the water-bath, stirring from time to time, until the fatty acids have collected on the surface as a clear layer.

Add about 20 g of wax (A.2.1), weigh with an accuracy of 0.01 g. When melting is complete, mix with the fatty acids by means of the stirrer so as to obtain a homogeneous layer. Remove the basin from the water-bath and allow to cool. The supernatant layer solidifies as a cake.

When the cake is hard enough, draw off the underlying water as completely as possible. Add approximately 100 ml of boiling distilled water, which melts the cake. Wash it by stirring in the distilled water added, and allow to cool again, so that the cake re-forms. Remove the washing water, and then the whole of the cake. Dry it as well as possible with filter paper.

After breaking the cake into pieces, place it in the previously tared small basin (C2) containing the stirrer thermometer.

Drive off the water by placing the basin and its contents on the sand-bath at low heat, stirring constantly with the thermometer, without exceeding 105°C.

Cease drying as soon as bubbles of water vapour are no longer given off.

Leave to cool in the desiccator and weigh.

A.5 EXPRESSION OF RESULTS

A.5.1 Method of calculation and formula

If E is the mass, in grammes, of the test portion,
 m is the mass, in grammes, of the wax cake after cooling,
 m_1 is the mass, in grammes, of the wax added,
 the percentage of total crude fatty acids in the soap is

$$(m - m_1) \times \frac{100}{E}$$

If M_0 is the nominal mass, in grammes, of the piece of soap,
 M_1 is the mean mass, in grammes, of the piece of soap at the time of analysis,
 the percentage content of total crude fatty acids in the initial soap is

$$(m - m_1) \times \frac{100}{E} \times \frac{M_1}{M_0}$$

A.5.2 Note

This result should be increased by 0.5 per cent absolute for soaps containing fatty acids of coconut or palm kernel oil, to allow for losses on drying.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO/R 455:1965

<https://standards.iteh.ai/catalog/standards/sist/a3f9bbf5-3ce5-48db-a153-fca5c1aff1a/iso-r-455-1965>