
**Animal and vegetable fats and oils —
Determination of acid value and acidity**

*Corps gras d'origines animale et végétale — Détermination de l'indice
d'acide et de l'acidité*

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 660:2009

<https://standards.iteh.ai/catalog/standards/sist/fb1aa8f-c05c-47a5-9b54-5bb94fd5c06e/iso-660-2009>



PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 660:2009

<https://standards.iteh.ai/catalog/standards/sist/fb1aa8f-c05c-47a5-9b54-5bb94fd5c06e/iso-660-2009>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2009

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 660 was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 11, *Animal and vegetable fats and oils*.

This third edition cancels and replaces the second edition (ISO 660:1996), which has been technically revised. It also incorporates the Amendment (ISO 660:1996/Amd 1:2003).

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 660:2009
<https://standards.iteh.ai/catalog/standards/sist/fb1aa8f-c05c-47a5-9b54-5bb94fd5c06e/iso-660-2009>

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 660:2009

<https://standards.iteh.ai/catalog/standards/sist/fb1aa8f-c05c-47a5-9b54-5bb94fd5c06e/iso-660-2009>

Animal and vegetable fats and oils — Determination of acid value and acidity

1 Scope

This International Standard specifies three methods (two titrimetric and one potentiometric) for the determination of the acidity in animal and vegetable fats and oils, hereinafter referred to as fats. The acidity is expressed preferably as acid value, or alternatively as acidity calculated conventionally.

This International Standard is applicable to refined and crude vegetable or animal fats and oils, soap stock fatty acids or technical fatty acids. The methods are not applicable to waxes.

Since the methods are completely non-specific, they cannot be used to differentiate between mineral acids, free fatty acids, and other organic acids. The acid value, therefore, also includes any mineral acids that may be present.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document applies.

ISO 660:2009

ISO 661, *Animal and vegetable fats and oils — Preparation of test sample*

ISO 3696, *Water for analytical laboratory use — Specification and test methods*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

acid value

number of milligrams of potassium hydroxide required to neutralize the free fatty acids present in 1 g of fat, when determined in accordance with the procedure specified in this International Standard

NOTE The acid value is expressed in milligrams per gram.

3.2

acidity

content of free fatty acids determined according to the procedure specified in this International Standard

NOTE The acidity is expressed as a percentage by mass. If the result of the determination is reported as acidity without further explanation, this is, by convention, the acidity based on the oleic acid content.

4 Principle

The sample is dissolved in a suitable solvent mixture, and the acids present are titrated with an ethanolic or methanolic solution of potassium or sodium hydroxide.

The methods specified in 9.1 and 9.2 are reference methods.

5 Reagents

WARNING — Attention is drawn to the regulations which specify the handling of hazardous substances. Technical, organizational and personal safety measures shall be followed.

Use only reagents of recognized analytical grade, unless otherwise specified.

5.1 Solvent A for solvent mixture (5.3): ethanol, volume fraction, $\varphi \approx 96\%$.

As a replacement, propan-2-ol, volume fraction, $\varphi \approx 99\%$, can be used.

5.2 Solvent B for solvent mixture (5.3): diethyl ether, peroxide-free.

As a replacement, *tert*-butyl methyl ether, light petroleum (boiling range 40 °C to 60 °C) or toluene can be used.

WARNING — Diethyl ether is very flammable and may form explosive peroxides. Use with great caution.

5.3 Solvent mixture, mix equal volumes of solvent A and B, (e.g. $\varphi_A = 50\text{ ml}/100\text{ ml}$ and $\varphi_B = 50\text{ ml}/100\text{ ml}$).

For hard or animal fats, a solvent mixture of one volume of solvent A (e.g. 25 ml) and three volumes of *tert*-butyl methyl ether or toluene (e.g. 75 ml) is recommended.

Neutralize, just before use, by adding potassium hydroxide solution in the presence of 0,3 ml of the phenolphthalein solution per 100 ml of solvent mixture.

For the titration with aqueous KOH, the solvent propan-2-ol can be used.

5.4 Ethanol or methanol, of minimum volume fraction, $\varphi = 95\%$.

5.5 Sodium hydroxide or potassium hydroxide, ethanolic or methanolic standard volumetric solutions, amount of substance concentration $c(\text{NaOH})$ or $c(\text{KOH}) = 0,1\text{ mol/l}$ and $0,5\text{ mol/l}$. The concentration shall be checked with a standard volumetric HCl solution.

NOTE The ethanolic/methanolic sodium/potassium hydroxide solution may be replaced by an aqueous sodium/potassium hydroxide solution, but only if the volume of water introduced does not lead to phase separation.

5.6 Phenolphthalein, solution in ethanol, mass concentration, $\rho = 1\text{ g}/100\text{ ml}$.

5.7 Thymolphthalein, solution in ethanol, mass concentration, $\rho = 2\text{ g}/100\text{ ml}$.

5.8 Alkali blue 6B, solution in ethanol, mass concentration, $\rho = 2\text{ g}/100\text{ ml}$.

For dark-coloured fats, **alkali blue** or **thymolphthalein** shall be used.

5.9 Water in accordance with ISO 3696, grade 3.

6 Apparatus

Usual laboratory equipment and, in particular, the following.

- 6.1 **Burette**, capacity 10 ml, graduated in 0,02 ml, ISO 385^[1] class A.
- 6.2 **Burette**, capacity 25 ml, graduated in 0,05 ml, ISO 385^[1] class A.
- 6.3 **Analytical balance**, capable of being read to the nearest 0,001 g.
- 6.4 **Automatic titration apparatus** (based on potentiometric electrode) or potentiometer.
- 6.5 **Combined pH electrode** for non-aqueous acid/base titrations.
- 6.6 **Graduated volumetric flasks**, volume 1 000 ml, ISO 1042^[2] class A.

7 Sampling

A representative sample should have been sent to the laboratory. It should not have been damaged or changed during transport or storage.

Sampling is not part of the method specified in this International Standard. A recommended sampling method is given in ISO 5555^[3].

iTeh STANDARD PREVIEW
(standards.iteh.ai)

8 Preparation of test sample

Prepare the test sample in accordance with ISO 661, except that if the sample contains volatile fatty acids, the test sample shall not be heated and filtered.

9 Procedure

9.1 Cold solvent method using indicator (Reference method)

9.1.1 Depending on the expected magnitude of the acid value, select the test portion mass and alkali concentration from Table 1.

9.1.2 According to Table 1 weigh the test portion into a 250 ml conical flask.

9.1.3 Add 50 ml to 100 ml of the neutralized solvent mixture (5.3) and dissolve the test portion if necessary with gentle warming.

For high melting point samples, use an ethanol-toluene mixture.

9.1.4 After the addition of an indicator (5.6, 5.7 or 5.8), titrate with constant swirling using standard potassium hydroxide solution (5.5). The endpoint of the titration is reached when the addition of a single drop of alkali produces a slight but definite colour change persisting for at least 15 s.

Table 1 — Test portion masses and alkali concentrations

Product group (examples)	Acid value approx.	Mass of test portion	Concentration of KOH	Accuracy of weighing of the test portion
		g	mol/l	g
Refined vegetable oils Animal fats	0 to 1	20	0,1	0,05
Crude vegetable oils Technical grade animal fats	1 to 4 4 to 15	10 2,5	0,1 0,1	0,02 0,01
Soap stock fatty acids	15 to 75	0,5 3,0	0,1 0,5	0,001
Technical fatty acids	> 75	0,2 1,0	0,1 0,5	0,001

9.2 Cold solvent method using potentiometric titration (Reference method)

9.2.1 According to Table 1, weigh the test portion into a 150 ml beaker.

9.2.2 Add 50 ml to 100 ml of the neutralized solvent mixture (5.3) and dissolve the sample, if necessary with gentle warming.

For high melting point samples, use an ethanol-toluene mixture

9.2.3 Introduce the combined electrode in the solvent mixture and connect it with the automatic titration apparatus.

9.2.4 Start the stirrer for at least 30 s and then titrate with constant swirling using standard potassium hydroxide solution (5.5).

9.2.5 As soon as the equivalence point is reached, record the amount of standard solution used.

9.3 Hot ethanol method using indicator

9.3.1 Under the conditions specified in this method, short-chain fatty acids, if present, are volatile.

9.3.2 Weigh into a flask a sufficient mass of the test sample as shown in Table 1, according to the colour and expected acid value.

9.3.3 Heat to boiling 50 ml of the ethanol containing 0,5 ml of the phenolphthalein indicator in a second flask. While the temperature of the ethanol is still above 70 °C, neutralize it carefully with a solution of 0,1 mol/l sodium or potassium hydroxide.

The endpoint of the titration is reached when the addition of a single drop of alkali produces a slight but definite colour change persisting for at least 15 s.

Larger volumes of ethanol and indicator may be necessary for dark-coloured fats. Moreover, for dark-coloured fats, alkali blue or thymolphthalein shall be used.

9.3.4 Add the neutralized ethanol to the test portion in the first flask and mix thoroughly. Bring the contents to the boil and titrate with the sodium or potassium hydroxide solution, agitating the flask contents vigorously during the titration.

10 Calculation

The acid value, w_{AV} , or the free fatty acid content, w_{FFA} , is reported as follows:

- to two decimal places for values between 0 up to and including 1;
- to one decimal place for values between 1 up to and including 100;
- as a whole number for values > 100 .

In addition to the following calculations, the approximate free fatty acid content (acidity) is calculated from:

$$w_{FFA} = 0,5 \times w_{AV}$$

10.1 Acid value

The acid value, w_{AV} , expressed as a mass fraction, is equal to

$$w_{AV} = \frac{56,1 \times c \ V}{m}$$

where

c is the exact concentration, in moles per litre, of the standard volumetric sodium or potassium hydroxide solution used;

V is the volume, in millilitres, of standard volumetric sodium or potassium hydroxide solution used;

m is the mass, in grams, of the test portion.

10.2 Acidity or free fatty acid content

The acidity or free fatty acid content, w_{FFA} , expressed as a percentage mass fraction, and according to fat type (see Table 2), is equal to

$$w_{FFA} = \frac{V \ c \ M \times 100}{1000 \times m}$$

where

V is the volume, in millilitres, of the standard volumetric sodium or potassium hydroxide solution used;

c is the concentration, in moles per litre, of the standard volumetric sodium or potassium hydroxide solution used;

M is the molar mass, in grams per mole, of the acid chosen for expression of the result (see Table 2) according to the fat type;

m is the mass, in grams, of the test portion.