



SLOVENSKI STANDARD

SIST ISO 10710:2011

01-junij-2011

Kakovost vode - Preskus zaviranja rasti z morsko in brakično makroalgo Ceranium tenuicorne

Water quality - Growth inhibition test with the marine and brackish water macroalga
Ceranium tenuicorne

iTeh STANDARD PREVIEW

Qualité de l'eau - Essai d'inhibition de croissance sur la macro algue d'eaux marine et
saumâtre Ceranium tenuicorne

[SIST ISO 10710:2011](https://standards.iteh.ai/catalog/standards/sist/446faf91-c1a0-4fff-ba51-1472a55b3e6a/sist-iso-10710-2011)

Ta slovenski standard je istoveten z: [ISO 10710:2010](https://standards.iteh.ai/catalog/standards/sist/446faf91-c1a0-4fff-ba51-1472a55b3e6a/sist-iso-10710-2011)

ICS:

13.060.70	Preiskava bioloških lastnosti vode	Examination of biological properties of water
-----------	------------------------------------	---

SIST ISO 10710:2011

en

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[SIST ISO 10710:2011](https://standards.iteh.ai/catalog/standards/sist/446faf91-c1a0-4fff-ba51-1472a53b3e6a/sist-iso-10710-2011)

<https://standards.iteh.ai/catalog/standards/sist/446faf91-c1a0-4fff-ba51-1472a53b3e6a/sist-iso-10710-2011>

INTERNATIONAL
STANDARD

ISO
10710

First edition
2010-06-15

**Water quality — Growth inhibition test
with the marine and brackish water
macroalga *Ceramium tenuicorne***

*Qualité de l'eau — Essai d'inhibition de croissance sur la macro algue
d'eaux marine et saumâtre Ceramium tenuicorne*

iTeh STANDARD PREVIEW
(standards.iteh.ai)

SIST ISO 10710:2011

<https://standards.iteh.ai/catalog/standards/sist/446faf91-c1a0-4ff-ba51-1472a53b3e6a/sist-iso-10710-2011>



Reference number
ISO 10710:2010(E)

© ISO 2010

ISO 10710:2010(E)

PDF disclaimer

This PDF file may contain embedded typefaces. In accordance with Adobe's licensing policy, this file may be printed or viewed but shall not be edited unless the typefaces which are embedded are licensed to and installed on the computer performing the editing. In downloading this file, parties accept therein the responsibility of not infringing Adobe's licensing policy. The ISO Central Secretariat accepts no liability in this area.

Adobe is a trademark of Adobe Systems Incorporated.

Details of the software products used to create this PDF file can be found in the General Info relative to the file; the PDF-creation parameters were optimized for printing. Every care has been taken to ensure that the file is suitable for use by ISO member bodies. In the unlikely event that a problem relating to it is found, please inform the Central Secretariat at the address given below.

iTeh STANDARD PREVIEW (standards.iteh.ai)

SIST ISO 10710:2011

<https://standards.iteh.ai/catalog/standards/sist/446faf91-c1a0-4fff-ba51-1472a53b3e6a/sist-iso-10710-2011>

**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2010

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Contents

Page

Foreword	iv
Introduction.....	v
1 Scope	1
2 Terms and definitions	1
3 Principle.....	2
4 Test organisms, nutrients, media, and materials.....	2
5 Cultivation	6
6 Apparatus	6
7 Procedure	7
7.1 Algae material	7
7.2 Preparation of dilution water.....	8
7.3 Choice of test concentrations	8
7.4 Start of test.....	9
7.5 Incubation.....	9
7.6 Measurement	9
7.7 Test with reference substances.....	9
8 Validity criteria.....	9
9 Evaluation of results	10
9.1 Calculations	10
9.2 Determination of $E_r C_x$	10
9.3 Expression of results	10
9.4 Interpretation of results	11
10 Reproducibility	11
11 Test report.....	11
Annex A (informative) Growth pattern of <i>Ceramium tenuicorne</i> in 7S and 20S.....	13
Annex B (informative) Results of ring test with <i>Ceramium tenuicorne</i> at 20 S.....	16
Bibliography.....	17

ISO 10710:2010(E)**Foreword**

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 10710 was prepared by Technical Committee ISO/TC 147, *Water quality*, Subcommittee SC 5, *Biological methods*.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

SIST ISO 10710:2011

<https://standards.iteh.ai/catalog/standards/sist/446faf91-c1a0-4fff-ba51-1472a53b3e6a/sist-iso-10710-2011>

Introduction

The red macroalga *Ceramium tenuicorne* belongs to Ceramiaceae, Rhodophyta. The species can be used as a model organism for the near coastal ecosystem. This species is found in temperate marine waters in both the northern and southern hemispheres and is thus relevant for large areas. As primary producers, they are a food source for many invertebrates and serve as living habitat for bacteria, invertebrates, and juvenile fish. They also serve as substrate for many oviparous fish species.

iTeh STANDARD PREVIEW (standards.iteh.ai)

[SIST ISO 10710:2011](https://standards.iteh.ai/catalog/standards/sist/446faf91-c1a0-4fff-ba51-1472a53b3e6a/sist-iso-10710-2011)

<https://standards.iteh.ai/catalog/standards/sist/446faf91-c1a0-4fff-ba51-1472a53b3e6a/sist-iso-10710-2011>

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[SIST ISO 10710:2011](https://standards.iteh.ai/catalog/standards/sist/446faf91-c1a0-4fff-ba51-1472a53b3e6a/sist-iso-10710-2011)

<https://standards.iteh.ai/catalog/standards/sist/446faf91-c1a0-4fff-ba51-1472a53b3e6a/sist-iso-10710-2011>

Water quality — Growth inhibition test with the marine and brackish water macroalga *Ceramium tenuicorne*

WARNING — Persons using this International Standard should be familiar with normal laboratory practice. This International Standard does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices and to ensure compliance with any national regulatory conditions.

IMPORTANT — It is absolutely essential that tests conducted according to this International Standard be carried out by suitably trained staff.

1 Scope

This International Standard specifies a method for the determination of the inhibition of growth of the macroalga *Ceramium tenuicorne* by substances and mixtures contained in seawater or by waste water with salinities between 4S and 32S. This method is applicable to substances that are easily soluble in water.

NOTE With modifications as described in ISO 14442^[4] and ISO 5667-16^[2], the inhibitory effects of poorly soluble organic and inorganic materials, volatile compounds, metals, waste water, marine water samples, and elutriates of sediments can be tested.

2 Terms and definitions

SIST ISO 10710:2011

<https://standards.iteh.ai/catalog/standards/sist/446fa91-c1a0-4ff-ba51-1472e73b7c6a/sist-iso-10710-2011>

For the purposes of this document, the following terms and definitions apply.

2.1

algal length

length from the first division to the most distant tip of the plant

NOTE The algal length is expressed in millimetres.

See Figure 1.

2.2

control medium

combination of dilution water and/or nutrient medium used in the test

[ISO 20079:2005^[5], 3.6]

2.3

control batch

control medium including organisms used for testing

[ISO 20079:2005^[5], 3.5]

2.4

effective concentration

$E_x C_x$

concentration of test sample at which an effect of x % is measured, if compared to the control

[ISO 20079:2005^[5], 3.9]

ISO 10710:2010(E)**2.5****growth medium**

mixture of natural seawater and nutrients in which algal plants are cultivated, which is used for pre-cultures

NOTE Adapted from ISO 8692:2004^[3], 3.3.

2.6**growth rate**

μ

(water quality) proportional rate of increase in algal length per day

NOTE See Clause 9.

2.7**salinity****practical salinity**

S

(seawater) ratio K_{15} of the electrical conductivity of the seawater sample, at the temperature of 15 °C and a pressure of one standard atmosphere, to that of a potassium chloride solutions in which the mass fraction of KCl is $32,435 6 \times 10^{-3}$, at the same temperature and pressure

NOTE Adapted from Reference [14], p. 12.

2.8**test medium**

mixture of seawater, nutrients and test sample

NOTE Adapted from ISO 8692:2004^[3], 3.5.

2.9**test sample**

(water quality) aqueous sample, e.g. chemical substance, mixture of chemicals or waste water, for which the inhibitory effects on the growth of algae are determined

NOTE Adapted from ISO 8692:2004^[3], 3.4.

3 Principle

Algal tips from monocultures of ceramium female gametophytes are grown in defined test conditions and in a defined medium containing a range of concentrations of the test sample. The test solutions are incubated for a period of 7 d after which the increase in length is measured and the growth rate is calculated. The growth inhibition is determined as a reduction in growth rate, relative to control cultures grown under identical conditions.

When toxicity of samples is to be compared to the toxicity of other chemicals or waste waters, tests can be performed in artificial seawater. If the purpose of the testing is to assess and to predict effects in a specific receiving water body, the tests can be conducted with algae adapted to the salinity in the receiving water body. In this case, natural seawater from an uncontaminated site of the same properties is used.

4 Test organisms, nutrients, media, and materials

Unless otherwise stated, use only reagents of recognized analytical grade and distilled or demineralized water or water of equivalent purity.

4.1 Test organisms

Use either of the following marine algal clones:

- a) *Ceramium tenuicorne* Kützing Waern (7S clone originating from the Baltic Sea);
- b) *C. tenuicorne* Kützing Waern (20S and 30S clone originating from the Oslo fjord).

This alga is a widely distributed macroalgae species (phylum Rhodophyta) in estuarine and coastal areas. The strains recommended are available in unialgal, non-axenic cultures¹⁾.

NOTE 1 This growth inhibition test is based on two clones, which were formerly regarded as two different species. These species were the marine *Ceramium strictum* Harvey sensu Kylin and the brackish water species *C. tenuicorne* Kützing Waern. Complete interfertility (References [11][12]) and DNA data (Reference [10]) have shown that the two entities belong to the same species, with *C. tenuicorne* as the valid name. The marine clone (former *C. strictum*) used in this test was isolated in 1973 and originates from the Oslo fjord (20S to 25S). It has been maintained as a laboratory culture for over 30 years. The brackish water clone was isolated in 1995 and originates from the Baltic Sea, 20 km south of the Askö laboratory in northern Baltic proper (6S to 7S). Cultures can be maintained in the medium specified in Clause 5. Regular subculturing is necessary.

NOTE 2 Among the red algae, changes occur between haploid and diploid generations. In the growth inhibition test, the female gametophytic generation is used since it has an even dichotomous growth pattern and the fastest growth rate. In nature it is difficult to distinguish between male and female plants. This can be done in the laboratory where spermatangia are found on the branches of the male plants and trichogynes can be seen on the tips of the claws on the female plants.

NOTE 3 The Baltic Sea clone can be adapted and used in tests in salinities between 4S and 12S. The marine clone can be used as test organism in salinities between 12S and 32S.

4.2 Natural and artificial seawater

4.2.1 General

SIST ISO 10710:2011
<https://standards.iteh.ai/catalog/standards/sist/446faf91-c1a0-4fff-ba51-1472a53b3e6a/sist-iso-10710-2011>

Natural seawater is used for the cultivation of the algae and either natural or artificial seawater should be used for testing. The type of seawater to be used depends on the objective of the test. When natural seawater is used, care shall be taken to ensure that it is not polluted. Take special care to avoid contamination of the water by inorganic or organic substances during preparation and storage. Equipment made of copper shall not be used.

4.2.2 Artificial seawater

Prepare the stock solutions for artificial seawater according to Table 1.

Start with about one third of the desired volume of water, add the weighed quantities of chemicals in accordance with Table 1 and make up to volume with water.

This stock solution with a salinity of 100S (equivalent to 10 % mass per volume) has a durability of at least six months, if stored in darkness at room temperature. Before use, the stock solution should be diluted with water to the desired salinity. Adjust the pH to $8,0 \pm 0,2$ with 1 mol/l HCl or 1 mol/l NaOH.

The artificial seawater shall be sterilized by autoclaving or sterile filtration (pore size, 0,2 μm) before use. Re-check the pH after sterilization, and adjust if necessary to $8,0 \pm 0,2$ with 1 mol/l HCl or 1 mol/l NaOH, before use.

1) Suitable suppliers are: a) ITM, Department of Applied Environmental Research of Science, Stockholm University, S-106 91 Stockholm, Sweden; b) University of Oslo, Department of Biology, P.O. Box 1066 Blindern, N-0316 Oslo, Norway. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of these suppliers.