

First edition  
2011-09-15

Corrected version  
2012-06-06

---

---

**Soil quality — Determination of soil  
microbial diversity —**

Part 2:

**Method by phospholipid fatty acid  
analysis (PLFA) using the simple PLFA  
extraction method**

iTeh STANDARD PREVIEW  
(standards.iteh.ai)

*Qualité du sol — Détermination de la diversité microbienne du sol —*

*Partie 2: Méthode par analyse des acides gras phospholipidiques  
(PLFA) en utilisant la méthode simple d'extraction des PLFA*

ISO/TS 29843-2:2011

<https://standards.iteh.ai/catalog/standards/sist/1809eba5-6d97-49de-8ad7-96835ca9285e/iso-ts-29843-2-2011>



Reference number  
ISO/TS 29843-2:2011(E)

© ISO 2011

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

[ISO/TS 29843-2:2011](https://standards.iteh.ai/catalog/standards/sist/1809eba5-6d97-49de-8ad7-96835ca9285e/iso-ts-29843-2-2011)

<https://standards.iteh.ai/catalog/standards/sist/1809eba5-6d97-49de-8ad7-96835ca9285e/iso-ts-29843-2-2011>



**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2011

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Case postale 56 • CH-1211 Geneva 20  
Tel. + 41 22 749 01 11  
Fax + 41 22 749 09 47  
E-mail [copyright@iso.org](mailto:copyright@iso.org)  
Web [www.iso.org](http://www.iso.org)

Published in Switzerland

<b>Contents</b>		Page
Foreword .....		iv
Introduction .....		v
1	Scope .....	1
2	Normative references .....	1
3	Symbols and abbreviated terms (except chemical products and reagents).....	1
4	Principle .....	1
5	Test materials .....	2
5.1	Soil.....	2
5.2	Reagents .....	2
5.3	Apparatus .....	4
6	Procedures .....	5
6.1	Lipid extraction (Bligh-Dyer extraction).....	5
6.2	Separation of lipids by SI column .....	5
6.3	Derivatization — Transmethylation — Clean-up .....	5
6.4	PLFA analysis .....	6
Bibliography .....		7

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

[ISO/TS 29843-2:2011](https://standards.iteh.ai/catalog/standards/sist/1809eba5-6d97-49de-8ad7-96835ca9285e/iso-ts-29843-2-2011)

<https://standards.iteh.ai/catalog/standards/sist/1809eba5-6d97-49de-8ad7-96835ca9285e/iso-ts-29843-2-2011>

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

In other circumstances, particularly when there is an urgent market requirement for such documents, a technical committee may decide to publish other types of document:

- an ISO Publicly Available Specification (ISO/PAS) represents an agreement between technical experts in an ISO working group and is accepted for publication if it is approved by more than 50 % of the members of the parent committee casting a vote;
- an ISO Technical Specification (ISO/TS) represents an agreement between the members of a technical committee and is accepted for publication if it is approved by 2/3 of the members of the committee casting a vote.

An ISO/PAS or ISO/TS is reviewed after three years in order to decide whether it will be confirmed for a further three years, revised to become an International Standard, or withdrawn. If the ISO/PAS or ISO/TS is confirmed, it is reviewed again after a further three years, at which time it must either be transformed into an International Standard or be withdrawn.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO/TS 29843-2 was prepared by Technical Committee ISO/TC 190, *Soil quality*, Subcommittee SC 4, *Biological methods*.

ISO/TS 29843 consists of the following parts, under the general title *Soil quality — Determination of soil microbial diversity*:

- *Part 1: Method by phospholipid fatty acid analysis (PLFA) and phospholipid ether lipids (PLEL) analysis*
- *Part 2: Method by phospholipid fatty acid analysis (PLFA) using the simple PLFA extraction method*

In this corrected version, the missing Equation (1) in 6.1 has been added.

## Introduction

Phospholipids are essential components of membranes of all living cells. Extracted from soil samples in fatty acid form (PLFA: phospholipid fatty acids) or ether-linked isoprenoid side chains (PLEL: phospholipid ether lipid), they provide quantitative and qualitative insights into the soil's viable/active microbial biomass. Cellular enzymes hydrolyse and release the phosphate group within minutes to hours following cell death (Reference [1]). The determination of total PLFA and PLEL provides a quantitative measure of the viable biomass of soil, i.e. microorganisms from all three domains of the biosphere (bacteria, fungi and archaeobacteria). PLFA and PLEL can also allow for taxonomic differentiation within complex microbial communities (References [2] and [3]). This approach is now well established in soil ecology and serves as a phenotypic, and thus complementary, tool to genotypic (molecular genetic) approaches for determining microbial diversity. Apart from taxonomic descriptions, the PLFA technique enables the determination of physiological changes within microbial consortia. For instance, the monoenic PLFA 16:1 $\omega$ 7c and 18:1 $\omega$ 7c are increasingly converted to the cyclopropyl fatty acids cy17:0 and cy19:0 in *Gram-negative* bacteria in response to environmental stress (Reference [4]).

Different methodologies are available for the determination of soil fatty acids. These methodologies present different levels of complexity when applied and provide different levels of resolution in the description of soil microbial communities. ISO/TS 29843-1 deals with the generally called "extended PLFA extraction method" while this part of ISO/TS 29843 deals with the generally called "simple PLFA extraction method" (References [5] and [6]).

This part of ISO/TS 29843, which deals with the simple PLFA extraction method, is accessible to most research and analytical laboratories involved in soil sciences. This methodology can be used for a wide range of soils. It provides a broad diversity measurement of a soil microbial community at the phenotypic level. It can be applied to biomass estimation and can be used to differentiate microbial communities among different soil samples (with the aid of an adapted statistical method). This method is especially adapted for detecting rapid changes in the soil microbial community structure. It can also be used to give a rough description of microbial groups present in soil samples (e.g. *Gram-positive* bacteria, actinomycetes, fungi). Table 1 (adapted from Table 1 in Reference [5]), presents a comparison of the sensitivity of the "extended PLFA" versus "simple PLFA" techniques.

ISO/TS 29843-2:2011  
**Table 1 — Comparison of the sensitivity of the "simple" and "extended" PLFA techniques in characterizing shifts in the composition of microbial communities**

Property	PLFA (simple)	PLFA (extended)
Ability to differentiate between two communities (with the aid of multivariate statistical methods)	Yes	Yes
Applicability for biomass estimation	Yes	Yes
Ability to register all single components of an entire community structure ("fingerprint")	No	Yes
Ability to register FAs other than EL-FAs	No	Yes
Estimation of number of FAs in soil samples	<50	200 to 400
Capacity to determine the linkage of the FAs to lipids in the molecule	Yes, EL	Yes, EL, NEL
Capacity to detect defined FAs in lower concentrations in the soil extract	No	Yes
Capacity to detect unusual FAs in the soil extract	No	Yes
Number of available signatures of FAs for defined organisms	Few	High numbers
Relationships of FAs widespread in the profile	High	Natura
Ability to identify the organisms causing the shift in microbial community	No	Basically yes

This method has been derived from the one first proposed in Reference [7] and later modified in Reference [1]. This revised method has been widely used (Reference [8]) and has also been discussed and compared to the extended PLFA extraction method in peer-reviewed articles (References [5] and [6]).

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

[ISO/TS 29843-2:2011](https://standards.iteh.ai/catalog/standards/sist/1809eba5-6d97-49de-8ad7-96835ca9285e/iso-ts-29843-2-2011)

<https://standards.iteh.ai/catalog/standards/sist/1809eba5-6d97-49de-8ad7-96835ca9285e/iso-ts-29843-2-2011>

# Soil quality — Determination of soil microbial diversity —

## Part 2: Method by phospholipid fatty acid analysis (PLFA) using the simple PLFA extraction method

### 1 Scope

This part of ISO/TS 29843 specifies a simple method for the extraction of only phospholipid fatty acids (PLFA) from soils.

ISO/TS 29843-1 specifies an extended method for the extraction and determination of both PLFA and PLEL from soils.

### 2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 10381-6, *Soil quality — Sampling — Part 6: Guidance on the collection, handling and storage of soil under aerobic conditions for the assessment of microbiological processes, biomass and diversity in the laboratory*  
[ISO/TS 29843-2:2011](https://www.iso.org/standard/52011.html)

ISO/TS 29843-1, *Soil quality — Determination of soil microbial diversity — Part 1: Method by phospholipid fatty acid analysis (PLFA) and phospholipid ether lipids (PLEL) analysis*

### 3 Symbols and abbreviated terms (except chemical products and reagents)

FAs: fatty acids

EL-FAs: ester-linked FAs

NEL-FAs: non-ester-linked FAs

FAME: fatty acid methyl ester(s)

$w_w$ : mass fraction of water in the soil, in grams of water per gram of dry soil (g/g)


GC: gas chromatography

FID: flame ionization detector

HPLC: high-performance liquid chromatography

### 4 Principle

Lipids are extracted using the extraction procedure in Reference [7]. Lipid extracts are fractionated on neutral lipids, glycolipids and phospholipids by liquid chromatography using an SI column. Phospholipids are transformed into fatty acid methyl esters (FAME) by mild alkaline hydrolysis. The different FAMEs are measured using gas chromatography (GC). A schematic overview of the procedures is given in Figure 1.



iTeh STANDARD PREVIEW  
(standards.iteh.ai)  
ISO/TS 29843-2:2011  
<https://standards.iteh.ai/catalog/standards/sist/1809eba5-6d97-49de-8ad7-96835ca9285e/iso-ts-29843-2-2011>

Figure 1 — Schematic overview of PLFA analysis according to the simple extraction method

## 5 Test materials

### 5.1 Soil

Collect soil samples and prepare them as specified in ISO 10381-6. Determine the soil mass fraction of water in the soil,  $w_w$ . If samples which have been sieved in the fresh state are not analysed immediately, they may be kept at  $-20\text{ °C}$  or stored in chloroform after lipid extraction (see 6.1).

### 5.2 Reagents

During the analysis, unless otherwise stated, use only reagents of recognized analytical grade or HPLC grade when specified.

#### 5.2.1 Organic solvents.

5.2.1.1 Acetone,  $C_3H_6O$  (HPLC grade).

5.2.1.2 Chloroform,  $CHCl_3$  (HPLC grade).

5.2.1.3 Hexane,  $C_6H_{14}$ .

5.2.1.4 Methanol,  $CH_3OH$  (HPLC grade).



5.2.1.5 Toluene,  $C_7H_8$ .

## 5.2.2 Chemicals.

5.2.2.1 2,6-Di-*tert*-butyl-4-methylphenol (BHT),  $C_{15}H_{24}O$ .

5.2.2.2 Citric acid,  $C_6H_8O_7 \cdot H_2O$ .

5.2.2.3 Trisodium citrate,  $C_6H_5Na_3O_7 \cdot 2H_2O$ .

5.2.2.4 Silicic acid hydrate,  $SiO_2 \cdot nH_2O$ .

5.2.2.5 Anhydrous sodium sulfate,  $Na_2O_4S$ .

5.2.2.6 Potassium hydroxide, KOH.

5.2.2.7 Acetic acid,  $C_2H_4O_2$ .

5.2.2.8 Sodium hydroxide, NaOH.

5.2.2.9 Nonadecanoic acid methyl ester,  $C_{20}H_{40}O_2$ .

5.2.2.10 Nitrogen gas,  $N_2$ .

## 5.2.3 Buffers and standards.

5.2.3.1 **CM (chloroform/methanol) solution**, to a chloroform and methanol solution with the ratio 1:2, add 2,6-di-*tert*-butyl-4-methylphenol (BHT) (0,005 %).

5.2.3.2 **CB (citrate buffer) solution** consisting of the following:

- citric acid monohydrate, 0,15 mol/l, 15,76 g of  $C_6H_8O_7 \cdot H_2O$  in 500 ml of  $H_2O$ ;
- trisodium citrate, 0,15 mol/l, 22,06 g of  $C_6H_5Na_3O_7 \cdot 2H_2O$  in 500 ml of  $H_2O$ ;
- for pH 4, add 59 ml of citric acid solution to 41 ml of trisodium citrate solution.

5.2.3.3 **BD (Bligh and Dyer) solvent**, to a chloroform/methanol:citrate buffer solution with the ratio 1:2:0,8, add 2,6-di-*tert*-butyl-4-methylphenol (BHT) (0,005 %).

EXAMPLE (100 ml of chloroform:200 ml of methanol:80 ml of CB) + BHT.

5.2.3.4 **Methanolic KOH solution**, 0,2 mol/l, 0,56 g of KOH in 50 ml of dry methanol (anhydrous sodium sulfate), freshly prepared.

5.2.3.5 **SE (solvent for extraction)**, hexane and chloroform with the ratio 4:1 (volume fraction).

5.2.3.6 **Acetic acid**, 1 mol/l, 58 ml/l. Add 58 ml of acetic acid to 750 ml of distilled water and fill up with distilled water to 1 l.

5.2.3.7 **Sodium hydroxide**, 0,3 mol/l, 12 g/l. Dissolve 12 g of sodium hydroxide in 750 ml of distilled water and fill up with distilled water to 1 l.