
Merjenje radioaktivnosti v okolju - Zrak: radon-222 - 2. del: Integrativna metoda za ugotavljanje povprečne potencialne koncentracije alfa energije njegovih kratkoživih razpadnih produktov (ISO 11665-2:2012)

Measurement of radioactivity in the environment - Air: radon-222 - Part 2: Integrated measurement method for determining average potential alpha energy concentration of its short-lived decay products (ISO 11665-2:2012)

Ermittlung der Radioaktivität in der Umwelt - Luft: Radon-222 - Teil 2: Integrierendes Messverfahren für die Bestimmung des Durchschnittswertes der potenziellen Alpha-Energiekonzentration der kurzlebigen Radon-Folgeprodukte (ISO 11665-2:2012)

<https://standards.iteh.ai/catalog/standards/sist/ea87a7e7-4667-4896-b049-1a0c1b1011665-2-140>

Mesurage de la radioactivité dans l'environnement - Air: radon 222 - Partie 2: Méthode de mesure intégrée pour la détermination de l'énergie alpha potentielle volumique moyenne de ses descendants à vie courte (ISO 11665-2:2012)

Ta slovenski standard je istoveten z: EN ISO 11665-2:2015

ICS:

13.040.99	Drugi standardi v zvezi s kakovostjo zraka	Other standards related to air quality
17.240	Merjenje sevanja	Radiation measurements

SIST EN ISO 11665-2:2015 en

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[SIST EN ISO 11665-2:2015](#)

<https://standards.iteh.ai/catalog/standards/sist/ea87a7e7-4667-4896-b049-160ca54578f5/sist-en-iso-11665-2-2015>

EUROPEAN STANDARD

EN ISO 11665-2

NORME EUROPÉENNE

EUROPÄISCHE NORM

September 2015

ICS 17.240

English Version

Measurement of radioactivity in the environment - Air:
radon-222 - Part 2: Integrated measurement method for
determining average potential alpha energy concentration
of its short-lived decay products (ISO 11665-2:2012)

Mesurage de la radioactivité dans l'environnement -
Air: radon 222 - Partie 2: Méthode de mesure intégrée
pour la détermination de l'énergie alpha potentielle
volumique moyenne de ses descendants à vie courte
(ISO 11665-2:2012)

Ermittlung der Radioaktivität in der Umwelt - Luft:
Radon-222 - Teil 2: Integrierendes Messverfahren für
die Bestimmung des Durchschnittswertes der
potenziellen Alpha-Energiekonzentration der
kurzlebigen Radon-Folgeprodukte (ISO 11665-2:2012)

This European Standard was approved by CEN on 12 June 2015.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels

Contents	Page
European foreword.....	3

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[SIST EN ISO 11665-2:2015](https://standards.iteh.ai/catalog/standards/sist/ea87a7e7-4667-4896-b049-160ca54578f5/sist-en-iso-11665-2-2015)
<https://standards.iteh.ai/catalog/standards/sist/ea87a7e7-4667-4896-b049-160ca54578f5/sist-en-iso-11665-2-2015>

European foreword

The text of ISO 11665-2:2012 has been prepared by Technical Committee ISO/TC 85 “Nuclear energy, nuclear technologies, and radiological protection” of the International Organization for Standardization (ISO) and has been taken over as EN ISO 11665-2:2015 by Technical Committee CEN/TC 430 “Nuclear energy, nuclear technologies, and radiological protection” the secretariat of which is held by AFNOR.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by March 2016, and conflicting national standards shall be withdrawn at the latest by March 2016.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

Endorsement notice

[SIST EN ISO 11665-2:2015](http://standards.iteh.ai/catalog/standards/sist/ca87c777-4667-4896-b049-160ca54578f5/sist-en-iso-11665-2-2015)

The text of ISO 11665-2:2012 has been approved by CEN as EN ISO 11665-2:2015 without any modification.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[SIST EN ISO 11665-2:2015](https://standards.iteh.ai/catalog/standards/sist/ea87a7e7-4667-4896-b049-160ca54578f5/sist-en-iso-11665-2-2015)

<https://standards.iteh.ai/catalog/standards/sist/ea87a7e7-4667-4896-b049-160ca54578f5/sist-en-iso-11665-2-2015>

INTERNATIONAL
STANDARDISO
11665-2First edition
2012-07-15

**Measurement of radioactivity in the
environment — Air: radon-222 —**

Part 2:

**Integrated measurement method for
determining average potential alpha
energy concentration of its short-lived
decay products**

iTeh STANDARD PREVIEW
(standards.iteh.ai)

*Mesurage de la radioactivité dans l'environnement — Air: radon 222 —**Partie 2: Méthode de mesure intégrée pour la détermination de l'énergie
alpha potentielle volumique moyenne de ses descendants à vie courte*

<https://standards.iteh.ai/catalog/standards/sist/ca87a7c7-4607-4876-b049-160ca54578f5/sist-en-iso-11665-2-2015>

Reference number
ISO 11665-2:2012(E)

© ISO 2012

iTeh STANDARD PREVIEW (standards.iteh.ai)

SIST EN ISO 11665-2:2015

<https://standards.iteh.ai/catalog/standards/sist/ea87a7e7-4667-4896-b049-160ca54578f5/sist-en-iso-11665-2-2015>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2012

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Case postale 56 • CH-1211 Geneva 20
Tel. + 41 22 749 01 11
Fax + 41 22 749 09 47
E-mail copyright@iso.org
Web www.iso.org

Published in Switzerland

Contents	Page
Foreword	iv
Introduction	v
1 Scope	1
2 Normative references	1
3 Terms, definitions and symbols	1
3.1 Terms and definitions	1
3.2 Symbols	2
4 Principle of the measurement method	3
5 Equipment	3
5.1 General	3
5.2 Measuring device	3
5.3 Counting system	4
6 Sampling	4
6.1 Sampling objective	4
6.2 Sampling characteristics	4
6.3 Sampling conditions	5
7 Detection method	6
8 Measurement	6
8.1 Procedure	6
8.2 Influence quantities	6
8.3 Calibration	7
9 Expression of results	7
9.1 Average potential alpha energy concentration	7
9.2 Standard uncertainty	8
9.3 Decision threshold and detection limit	8
9.4 Limits of the confidence interval	9
10 Test report	9
Annex A (informative) Example of a method meeting the requirements of this part of ISO 11665	11
Bibliography	13

ISO 11665-2:2012(E)

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 11665-2 was prepared by Technical Committee ISO/TC 85, *Nuclear energy, nuclear technologies, and radiological protection*, Subcommittee SC 2, *Radiological protection*.

ISO 11665 consists of the following parts, under the general title *Measurement of radioactivity in the environment — Air: radon-222*:

- *Part 1: Origins of radon and its short-lived decay products and associated measurement methods*
- *Part 2: Integrated measurement method for determining average potential alpha energy concentration of its short-lived decay products*
- *Part 3: Spot measurement method of the potential alpha energy concentration of its short-lived decay products*
- *Part 4: Integrated measurement method for determining average activity concentration using passive sampling and delayed analysis*
- *Part 5: Continuous measurement method of the activity concentration*
- *Part 6: Spot measurement method of the activity concentration*
- *Part 7: Accumulation method for estimating surface exhalation rate*
- *Part 8: Methodologies for initial and additional investigations in buildings*

The following parts are under preparation:

- *Part 9: Method for determining exhalation rate of dense building materials*
- *Part 10: Determination of diffusion coefficient in waterproof materials using activity concentration measurement*

Introduction

Radon isotopes 222, 220 and 219 are radioactive gases produced by the disintegration of radium isotopes 226, 224 and 223, which are decay products of uranium-238, thorium-232 and uranium-235 respectively, and are all found in the earth's crust. Solid elements, also radioactive, followed by stable lead are produced by radon disintegration^[1].

When disintegrating, radon emits alpha particles and generates solid decay products, which are also radioactive (polonium, bismuth, lead, etc.). The potential effects on human health of radon lie in its solid decay products rather than the gas itself. Whether or not they are attached to atmospheric aerosols, radon decay products can be inhaled and deposited in the bronchopulmonary tree to varying depths according to their size.

Radon is today considered to be the main source of human exposure to natural radiation. The UNSCEAR (2006) report^[2] suggests that, at the worldwide level, radon accounts for around 52 % of global average exposure to natural radiation. The radiological impact of isotope 222 (48 %) is far more significant than isotope 220 (4 %), while isotope 219 is considered negligible. For this reason, references to radon in this part of ISO 11665 refer only to radon-222.

Radon activity concentration can vary by one to multiple orders of magnitude over time and space. Exposure to radon and its decay products varies tremendously from one area to another, as it depends firstly on the amount of radon emitted by the soil and the building materials in each area and, secondly, on the degree of containment and weather conditions in the areas where individuals are exposed.

Variations of a few nanojoules per cubic metre to several thousand nanojoules per cubic metre are observed in the potential alpha energy concentration of short-lived radon decay products.

The potential alpha energy concentration of short-lived radon-222 decay products in the atmosphere can be measured by spot and integrated measurement methods (see ISO 11665-1). This part of ISO 11665 deals with integrated measurement methods. Integrated measuring methods are applicable in assessing human exposure to radiation^[3].

NOTE The origin of radon-222 and its short-lived decay products in the atmospheric environment and other measurement methods are described generally in ISO 11665-1.