
**Soil quality — Estimation of
abundance of selected microbial gene
sequences by quantitative PCR from
DNA directly extracted from soil**

*Qualité du sol — Estimation de l'abondance de séquences de gènes
microbiens par amplification par réaction de polymérisation en
chaîne (PCR) quantitative à partir d'ADN directement extrait du sol*

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 17601:2016

<https://standards.iteh.ai/catalog/standards/sist/837e9eee-02df-483a-be2f-1f21de376ab0/iso-17601-2016>



iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 17601:2016

<https://standards.iteh.ai/catalog/standards/sist/837e9eee-02df-483a-bc2f-1f21de376ab0/iso-17601-2016>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2016, Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Ch. de Blandonnet 8 • CP 401
CH-1214 Vernier, Geneva, Switzerland
Tel. +41 22 749 01 11
Fax +41 22 749 09 47
copyright@iso.org
www.iso.org

Contents

	Page
Foreword	iv
Introduction	v
1 Scope	1
2 Normative references	1
3 Terms and definitions	1
4 Principle	2
5 Test materials	4
5.1 DNA.....	4
5.2 Bacteria.....	4
5.3 Plasmid.....	4
5.4 Enzyme.....	4
5.5 Chemicals.....	4
5.6 Product for bacterial culture medium.....	5
5.7 Buffer and reagents.....	5
6 Apparatus	6
7 Procedure	6
7.1 qPCR standard preparation and calibration of qPCR assay (task 1).....	6
7.1.1 General.....	6
7.1.2 Amplicon design (task 1, step 1).....	6
7.1.3 qPCR standard preparation (task 1, step 2).....	7
7.1.4 Isolate DNA, environmental DNA, artificial DNA.....	7
7.1.5 Calibration of the qPCR (task 1, step 3).....	9
7.2 Preparation of soil DNA template and inhibition test (task 2).....	10
7.2.1 General.....	10
7.2.2 Soil DNA preparation (task 2, step 4).....	10
7.2.3 Inhibition test (task 2, step 5).....	10
7.2.4 Dilution of DNA template.....	12
7.3 qPCR assay (task 3).....	12
7.3.1 General.....	12
7.3.2 qPCR.....	12
7.4 Validation and analysis of qPCR assay (task 4).....	12
7.4.1 General.....	12
7.4.2 Validation of the qPCR assay.....	12
7.4.3 Calculation of the copy number of the gene of interest in the soil DNA extract.....	13
8 Examination of the critical steps of the qPCR assay	14
9 Expression of the results of the qPCR assay	14
10 International ring test	14
11 Test report	14
Annex A (informative) Description of principal steps of TaqMan® qPCR assay	15
Annex B (informative) International ring-test for evaluating qPCR to quantify the abundance of selected microbial gene sequences from DNA directly extracted from soil	17
Bibliography	30

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the WTO principles in the Technical Barriers to Trade (TBT) see the following URL: [Foreword - Supplementary information](#)

The committee responsible for this document is ISO/TC 190, *Soil quality*, Subcommittee SC 4, *Biological methods*.

ISO 17601:2016

<https://standards.iteh.ai/catalog/standards/sist/837e9eee-02df-483a-be2f-1f21de376ab0/iso-17601-2016>

Introduction

DNA (DNAs) is a major component of any living organisms coding for enzymes responsible for their biological activities. The study of DNA sequences from DNA sources extracted from different environmental matrices, by means of numerous molecular approaches, provides molecular markers that can be used to sharply distinguish and identify different organisms (bacteria, *archaea*, and *eucaryotes*).

Up to now, most of the studies aiming to develop microbial quality indicators applicable to complex environment such as soil were biased by the poor culturability of many microorganisms under laboratory conditions and the lack of sensitivity of traditional microbiological methods. The recent development of a large set of molecular biology methods based on amplification of soil-extracted nucleic acids have provided a pertinent alternative to classical culture-based microbiological methods providing unique insight into the composition, richness, and structure of microbial communities.[2] [3] [4] [5] [6] DNA-based approaches are now well established in soil ecology and serve as genotypic markers for determining microbial diversity. The results of molecular analyzes of soil microbial communities and/or populations rely on two main parameters: a) the extraction of DNA representative of the indigenous bacterial community composition and b) PCR bias such as the choice of primers, the concentration of amplified DNA, errors in the PCR, or even the method chosen for analysis.[7] [4] [8] [9]

Numerous studies have investigated new methods to improve extraction, purification, amplification, and quantification of DNA from soils.[10] Recently, ISO 11063 reporting “a method to extract nucleic acids directly from soil samples” derived from Reference [10] is opening a new window for developing standardized molecular approaches to estimate soil quality.[11]

The aim of this International Standard is to describe the procedure used to set up and perform quantitative PCR to quantify the abundance of soil microbial phyla, as well as functional groups from DNA directly extracted from soil samples. The quantification of soil microbial phyla, as well as functional groups by qPCR assays can contribute to the development of routine tools to monitor soil quality. The repeatability and the reproducibility of the procedure of the quantitative PCR were assessed in an international ring test study (see Annex B). The repeatability of this procedure was successfully evaluated for both 16S rRNA genes, as well as genes coding a functional marker of denitrifiers (the nitrite reductase gene *nirK*). The reproducibility of this procedure revealed a laboratory effect which can be overcome by interpreting the results of the quantification of the abundance of the microbial groups by comparison, either by using an external reference (DNA extracted from a control strain) in the assay or by calculating a percentage of variations between treatments to normalize the data.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 17601:2016

<https://standards.iteh.ai/catalog/standards/sist/837e9eee-02df-483a-be2f-1f21de376ab0/iso-17601-2016>

Soil quality — Estimation of abundance of selected microbial gene sequences by quantitative PCR from DNA directly extracted from soil

1 Scope

This International Standard specifies the crucial steps of a quantitative real-time polymerase chain reaction (qPCR) method to measure the abundance of selected microbial gene sequences from soil DNA extract which provides an estimation of selected microbial groups.

It is noteworthy that the number of genes is not necessarily directly linked to the number of organisms that are measured. For example, the number of ribosomal operon is ranging from one copy to 20 copies in different bacterial phyla. Therefore, the number of 16S rRNA sequences quantified from soil DNA extracts does not give an exact estimate of the number of soil bacteria. Furthermore, the number of sequences is not necessarily linked to living microorganisms and can comprise sequences amplified from dead microorganisms.

2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 10381-6, *Soil quality — Sampling — Part 6: Guidance on the collection, handling and storage of soil under aerobic conditions for the assessment of microbiological processes, biomass and diversity in the laboratory*

ISO 11063, *Soil quality — Method to directly extract DNA from soil samples*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

soil DNA

DNA extracted from soil of living and dead biota

EXAMPLE Microorganisms, plants, animals.

3.2

polymerase chain reaction

PCR

method allowing the amplification of a specific DNA sequence using a specific pair of oligonucleotide primers

3.3

quantitative polymerase chain reaction

qPCR

method allowing the quantification in a DNA *template* (3.4) of the number of a specific DNA sequence using a specific pair of oligonucleotide primers

3.4

template

DNA sample used to perform PCR (3.2) to amplify a specific DNA sequence

3.5

amplicon

PCR product obtained by *PCR* (3.2) from a *template* (3.4)

3.6

cloning vector

circular DNA molecule in which the *amplicon* (3.5) is inserted by ligation used to transform competent *Escherichia coli* for cloning the amplicon

3.7

qPCR standard

cloned DNA target used as *template* (3.4) for qPCR reaction to establish the standard curve relating the abundance of target sequence as a function of cycle threshold values (Ct)

3.8

non-template control

NTC

control, usually molecular grade water, that is used as negative control in qPCR assay to check for the absence of contaminant in the qPCR mix

3.9

cycle threshold

Ct

number of qPCR cycles required for the fluorescent signal to cross the threshold (i.e. exceeds background level)

Note 1 to entry: The Ct value is inversely proportional to the abundance of the target sequence.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

4 Principle

ISO 17601:2016

This International Standard describes qPCR assay using fluorescent DNA binding dye as reporter. This qPCR assay has been validated by an international ring test conducted with the SYBR Green, a commonly used fluorescent DNA binding dye which binds all double-stranded DNA and can be detected by measuring the increase in fluorescence throughout the cycle.

The method aims to measure the abundance of selected microbial gene sequences from soil DNA extract. The method comprises four tasks and eight steps as summarized in [Figure 1](#). According to Reference [1], the three critical steps to be validated for each qPCR assay are as shown in [Figure 1](#).




Figure 1 — Main tasks and critical steps to estimate the abundance of selected microbial gene sequences by qPCR assay

This International Standard describes qPCR assay based on the use of fluorescent DNA binding dye which has been validated by an international ring test using SYBR Green®¹⁾ qPCR. In Annex A,

1) SYBR Green is a registered trademark of Molecular Probes. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of the product named. Equivalent products may be used if they can be shown to lead to the same results.

information about TaqMan^{®2)} qPCR assay not tested in the international ring test are given. The first task is made of three steps describing the design of optimal amplicon for qPCR (step one), the preparation of qPCR standards (step two), and the procedure to calibrate the qPCR assay (step three). The second task includes two additional steps describing the procedures to prepare soil DNA samples (step four) and to test for the presence of qPCR inhibitors in soil DNA samples (step five). The third task is constituted of a single step describing the protocol to perform qPCR assay (step six). Finally, the fourth task is made of two steps, one describing the procedure to validate qPCR assays (step 7) to check the quality of qPCR assay and another one describing the different options to calculate the number of sequences of the gene of interest copy from cycle threshold (Ct) obtained from the analysis of qPCR amplification plots (step 8).

5 Test materials

5.1 DNA

5.1.1 DNA, extracted from pure bacterial and fungal isolates using classical extraction procedures or by using commercial kit to extract genomic DNA.

5.1.2 Soil DNA, extracted from aliquots of soil according to ISO 11063.

5.2 Bacteria

5.2.1 *Escherichia coli* strain, usually used for cloning of PCR product.

5.3 Plasmid

5.3.1 Cloning vector, usually used for cloning of PCR product in competent *Escherichia coli*.

5.4 Enzyme

5.4.1 Taq polymerase.

5.4.2 T4 DNA ligase.

5.4.3 T4 gene T32.

5.4.4 Bovine serum albumin (CAS No. 9048-46-8).

5.5 Chemicals

5.5.1 Ampicilline sodium, C₁₆H₁₈N₃NaO₄S (CAS No. 69-52-3).

5.5.2 Boric acid, BH₃O₃ (CAS No. 10043-35-3).

5.5.3 Deoxynucleotide solution, dNTPs.

5.5.4 SYBR Safe[®] DNA gel stain.

2) TaqMan is a trademark of Roche Molecular Systems, Inc. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of the product named. Equivalent products may be used if they can be shown to lead to the same results.

5.5.5 Ethylenediaminetetraacetic acid disodium salt (EDTA), $C_{10}H_{14}N_2O_8Na_2 \cdot 2 H_2O$ (CAS No. 6381-92-6).

5.5.6 Glucose, $C_6H_{12}O_6$ (CAS No. 50-99-7).

5.5.7 Chlorhydric acid, HCl (CAS No. 7647-01-0).

5.5.8 IPTG, Isopropyl-Beta-D-Thiogalactopyranoside, (CAS No. 367-93-1).

5.5.9 Magnesium chloride, $MgCl_2$ (CAS No. 7786-30-3).

5.5.10 Magnesium sulfate, $MgSO_4$ (CAS No. 7487-88-9).

5.5.11 Molecular-biology-grade water, H_2O .

5.5.12 Potassium chloride, KCl (CAS No. 7447-40-7).

5.5.13 Sodium chloride, NaCl (CAS No. 7647-14-5).

5.5.14 Tris[hydroxymethyl]aminomethane, $C_4H_{11}NO_3$ (CAS No. 77-86-1).

5.5.15 X-Gal, 5-bromo-4-chloro-3-indolyl-beta-D-galactopyranoside, (CAS No. 7240-90-6).

5.6 Product for bacterial culture medium

5.6.1 Bacto tryptone^{®3)}, enzymatic digest of casein.

5.6.2 Yeast extract powder (CAS No. 8013-01-2).

5.7 Buffer and reagents

5.7.1 Ampicilline solution, 2 g of ampicilline sodium in 4 ml of 0,22 μm filter sterilized H_2O . Adjust to 20 ml with sterilized H_2O , prepare 1 ml aliquots, and store at $-20\text{ }^\circ C$.

5.7.2 EDTA, 0,5 mol·l⁻¹, 186,10 g of EDTA in 1 000 ml of H_2O adjusting with NaOH (10 mol·l⁻¹) to pH 8,0.

5.7.3 SYBR Safe™ DNA gel stain, dilute 10,000X SYBR Safe™ gel stain in TBE buffer $\times 1$.

5.7.4 IPTG stock solution, 1 g of IPTG in 8 ml of H_2O . After careful mixing, the solution is adjusted to 10 ml and sterilized under security microbiology post. Prepare 1 ml aliquot of IPTG and store at $-20\text{ }^\circ C$.

5.7.5 Solid LB medium, 10 g of Bacto tryptone[®], 5 g of yeast extract, 5 g of sodium chloride, and 15 g of agar in 1 000 ml of H_2O . After autoclaving for 20 min at $120\text{ }^\circ C$, 1 ml of ampicilline stock solution at 100 mg·ml⁻¹ is added to LB medium and plated in Petri dishes (20 ml) under a security microbiology post. 100 μl of IPTG solution are plated on solid LB-ampicilline medium. When IPTG solution is entered in LB-ampicilline medium, 20 μl of X-Gal solution is plated on solid LB-ampicilline medium. Solid LB medium is stored at $4\text{ }^\circ C$ until its use.

3) Bacto tryptone is the trademark of a product supplied by Difco Laboratories. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of the product named. Equivalent products may be used if they can be shown to lead to the same results.

ISO 17601:2016(E)

5.7.6 SOC medium, 20 g of Bacto tryptone®, 5 g of yeast extract, 0,58 g of NaCl, 0,95 g of MgCl₂, 2,46 g of MgSO₄, and 3,60 g of glucose in 1 l H₂O. Sterilize by 20 min autoclaving at 120 °C. Prepare 950 ml aliquots and store at -20 °C.

5.7.7 Tris-HCl, 1 mol·l⁻¹, 121,14 g of Tris in 1 000 ml of H₂O adjusting with 4 mol·l⁻¹HCl to pH 8,0.

TBE buffer × 10, pH 8,0, 108 g of Tris base, 55 g of boric acid, and 40 ml of 0,5 mol·l⁻¹EDTA (pH 8,0) in 1 000 ml of H₂O.

5.7.8 TBE buffer × 1, 100 ml of TBE buffer × 10 in 900 ml of H₂O.

5.7.9 TE buffer × 10, pH 8,0, 100 ml of 1 mol·l⁻¹Tris-HCl pH 8,0, 20 ml of 50 mmol·l⁻¹EDTA pH 8,0 in 880 ml of molecular grade water.

5.7.10 TE buffer × 1, 100 ml of TE buffer × 10 in 900 ml of H₂O.

5.7.11 X-gal solution, 250 mg of X-Gal in 5 ml of dimethylformamide 5 ml. After careful mixing, prepare 0,5 ml aliquot and store at -20 °C.

6 Apparatus

Use standard laboratory equipment including pipettes, a centrifuge, fume hood cabinet, horizontal electrophoresis system and the following.

6.1 Quantitative PCR, allowing the real-time quantification of amplicons from various DNA templates with a theoretical detection limit of one copy of a sequence target per sample analyzed.

6.2 Spectro-photometer, allowing the quantification of double-strand DNA at 260 nm.

6.3 Spectro-fluorimeter, allowing the quantification of double-strand DNA.

NOTE Only one of these two apparatus is required to estimate DNA concentration.

7 Procedure

7.1 qPCR standard preparation and calibration of qPCR assay (task 1)

7.1.1 General

qPCR assay is based on the quantification of the amplicons at the end of each PCR cycle by using a DNA dye which fluoresces when intercalated in the double strand amplicons. The purpose of this task is to describe the definition of appropriate amplicon to settle down a qPCR assay (step one), the preparation of qPCR standard (step two), and the calibration of the qPCR assay (step three).

7.1.2 Amplicon design (task 1, step 1)

7.1.2.1 General

The first step aims at designing oligonucleotide primer pair; it can be designed *in silico* using different programs using the sequence of the microbial gene of interest to be quantified by qPCR from soil DNA extracts. The specificity of the primers shall be checked *in silico* by comparing their sequences to known sequences available in the Genbank database (<http://www.ncbi.nlm.nih.gov/genbank/>). Only primers specific for the gene target shall be considered. The main parameters to be considered to design oligonucleotide primer pair for establishing qPCR assay are listed thereafter.