

SLOVENSKI STANDARD
oSIST prEN ISO 23161:2017
01-januar-2017

Kakovost tal - Določevanje izbranih organokositrovih spojin - Metoda plinske kromatografije (ISO/FDIS 23161:2016)

Soil quality - Determination of selected organotin compounds - Gas-chromatographic method (ISO/FDIS 23161:2016)

Bodenbeschaffenheit - Bestimmung ausgewählter Organozinnverbindungen - Gaschromatographisches Verfahren (ISO/FDIS 23161:2016)

Qualité du sol - Dosage d'une sélection de composés organostanniques - Méthode par chromatographie en phase gazeuse (ISO/FDIS 23161:2016)

<https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-e7a267c0801b/osist-pr-en-iso-23161-2017>

Ta slovenski standard je istoveten z: prEN ISO 23161

ICS:

13.080.10	Kemijske značilnosti tal	Chemical characteristics of soils
71.040.50	Fizikalnokemijske analitske metode	Physicochemical methods of analysis

oSIST prEN ISO 23161:2017

en,fr,de

iTeh STANDARD PREVIEW (standards.iteh.ai)

[oSIST prEN ISO 23161:2017](https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-e7a267c0801b/osist-pren-iso-23161-2017)

<https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-e7a267c0801b/osist-pren-iso-23161-2017>

FINAL
DRAFT

INTERNATIONAL
STANDARD

ISO/FDIS
23161

ISO/TC 190/SC 3

Secretariat: DIN

Voting begins on:
2016-12-02

Voting terminates on:
2017-02-24

Soil quality — Determination of selected organotin compounds — Gas- chromatographic method

*Qualité du sol — Dosage d'une sélection de composés
organostanniques — Méthode par chromatographie en phase gazeuse*

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[oSIST prEN ISO 23161:2017](https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-e7a267c0801b/osist-pren-iso-23161-2017)

[https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-
e7a267c0801b/osist-pren-iso-23161-2017](https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-e7a267c0801b/osist-pren-iso-23161-2017)

RECIPIENTS OF THIS DRAFT ARE INVITED TO SUBMIT, WITH THEIR COMMENTS, NOTIFICATION OF ANY RELEVANT PATENT RIGHTS OF WHICH THEY ARE AWARE AND TO PROVIDE SUPPORTING DOCUMENTATION.

IN ADDITION TO THEIR EVALUATION AS BEING ACCEPTABLE FOR INDUSTRIAL, TECHNOLOGICAL, COMMERCIAL AND USER PURPOSES, DRAFT INTERNATIONAL STANDARDS MAY ON OCCASION HAVE TO BE CONSIDERED IN THE LIGHT OF THEIR POTENTIAL TO BECOME STANDARDS TO WHICH REFERENCE MAY BE MADE IN NATIONAL REGULATIONS.

ISO/CEN PARALLEL PROCESSING



Reference number
ISO/FDIS 23161:2016(E)

© ISO 2016

iTeh STANDARD PREVIEW (standards.iteh.ai)

[oSIST prEN ISO 23161:2017](https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-e7a267c0801b/osist-pren-iso-23161-2017)

<https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-e7a267c0801b/osist-pren-iso-23161-2017>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2016, Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Ch. de Blandonnet 8 • CP 401
CH-1214 Vernier, Geneva, Switzerland
Tel. +41 22 749 01 11
Fax +41 22 749 09 47
copyright@iso.org
www.iso.org

Contents

Page

Foreword	iv
Introduction	v
1 Scope	1
2 Normative references	2
3 Terms and definitions	2
4 Principle	2
5 Reagents	4
5.1 General	4
5.2 Chemicals	4
5.3 Standards	5
5.4 Preparation of reagents and solutions	6
5.5 Clean-up	7
6 Apparatus	8
6.1 Requirements for glassware	8
6.2 Sampling apparatus	8
6.3 Additional apparatus	8
7 Procedure	9
7.1 Sampling and sample pretreatment	9
7.2 Sample extraction	9
7.2.1 General	9
7.2.2 Acidic extraction and derivatization of an aliquot	10
7.2.3 Alkaline treatment and <i>in situ</i> derivatization	10
7.2.4 Separate determination of TTBT in the field-moist sample	10
7.3 Clean-up of the extract	11
7.3.1 General	11
7.3.2 Silica and aluminium oxide clean-up	11
7.4 Determination of dry mass	11
7.5 Measurement	11
7.5.1 Gas chromatographic separation	11
7.5.2 Detection and identification	12
8 Calibration	12
9 Recovery rates of the internal standard compounds	13
10 Quantification	14
11 Expression of results	15
12 Validation	15
13 Test report	15
Annex A (informative) Information about the procedure	16
Annex B (informative) Additional clean-up procedures	18
Annex C (informative) Information about typical instrumental conditions	20
Annex D (informative) Information about GC-MS identification	31
Annex E (informative) Validation data	33
Bibliography	37

ISO/FDIS 23161:2016(E)

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see the following URL: www.iso.org/iso/foreword.html.

The committee responsible for this document is ISO/TC 190, *Soil quality*, Subcommittee SC 3, *Chemical methods and soil characteristics*.

This second edition cancels and replaces the first edition (ISO 23161:2009), of which it constitutes a minor revision.

The changes compared to the previous edition are as follows:

- the Note to [Clause 1](#) and [Table 2](#) have been moved to [Clause 4](#);
- former Note 4 to [Clause 4](#) has been changed to normal text and moved above Note 1;
- former second sentence in [5.5.5](#) has been changed to Note;
- in [7.2.2](#) and [7.2.3](#), a Note has been added;
- the presentation of tables in [Annex E](#) has been improved;
- the Bibliography has been updated.

Introduction

It is absolutely essential that tests conducted in accordance with this document be carried out by suitably qualified staff.

It can be noted whether, and to what extent, particular problems will require the specification of additional boundary conditions.

iTeh STANDARD PREVIEW (standards.iteh.ai)

[oSIST prEN ISO 23161:2017](https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-e7a267c0801b/osist-pren-iso-23161-2017)

<https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-e7a267c0801b/osist-pren-iso-23161-2017>

iTeh STANDARD PREVIEW **(standards.iteh.ai)**

[oSIST prEN ISO 23161:2017](https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-e7a267c0801b/osist-pren-iso-23161-2017)

<https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-e7a267c0801b/osist-pren-iso-23161-2017>

Soil quality — Determination of selected organotin compounds — Gas-chromatographic method

WARNING — Persons using this document should be familiar with normal laboratory practice. This document does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices and to ensure compliance with any national regulatory conditions.

IMPORTANT — It is absolutely essential that tests, conducted in accordance with this document, be carried out by suitably qualified staff.

1 Scope

This document specifies a gas-chromatographic method for the identification and quantification of organotin compounds (OTCs) in soils as specified in [Table 1](#).

This document is also applicable to samples from sediments, sludges and wastes (soil-like materials).

The working range depends on the detection technique used and the amount of sample taken for analysis.

The limit of quantification for each compound is about 10 µg/kg.

Table 1 — Organotin compounds, which can be determined in accordance with this document

$R_n\text{Sn}^{(4-n)+}$	R	n	Name	Acronym
Organotin cations^a				
BuSn^{3+}	Butyl	1	Monobutyltin cation	MBT
$\text{Bu}_2\text{Sn}^{2+}$	Butyl	2	Dibutyltin cation	DBT
Bu_3Sn^+	Butyl	3	Tributyltin cation	TBT
OcSn^{3+}	Octyl	1	Monooctyltin cation	MOT
$\text{Oc}_2\text{Sn}^{2+}$	Octyl	2	Dioctyltin cation	DOT
Ph_3Sn^+	Phenyl	3	Triphenyltin cation	TPhT
Cy_3Sn^+	Cyclohexyl	3	Tricyclohexyltin cation	TCyT
Peralkylated organotin				
Bu_4Sn	Butyl	4	Tetrabutyltin	TTBT
^a Organotin compounds are measured after derivatization.				

Organotin cations can only be determined in accordance with this document after derivatization. The anionic part bound to the organotin cation is mainly dependent on the chemical environment and is not determined using this method. The peralkylated organotin compounds behave in a completely different way from their parent compounds. Tetraalkylated organotin compounds which are already peralkylated, such as tetrabutyltin, are determined directly without derivatization.

The properties such as particle size distribution, water content and organic matter content of the solids to be analysed using this document vary widely. Sample pretreatment is designed adequately with respect to both the properties of the organotin compounds and the matrix to be analysed.

ISO/FDIS 23161:2016(E)

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 3696, *Water for analytical laboratory use — Specification and test methods*

ISO 11465, *Soil quality — Determination of dry matter and water content on a mass basis — Gravimetric method*

ISO 16720, *Soil quality — Pretreatment of samples by freeze-drying for subsequent analysis*

ISO 22892, *Soil quality — Guidelines for the identification of target compounds by gas chromatography and mass spectrometry*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <http://www.electropedia.org/>
- ISO Online browsing platform: available at <http://www.iso.org/obp>

3.1

organotin compound

substance containing 1 to 4 Sn-C bonds

Note 1 to entry: The number of Sn-C bonds is a measure for the degree of substitution.

3.2

organotin cation

part of the *organotin compound* (3.1) that contains all Sn-C bonds and is formally charged

3.3

organotin cation derivatives

non-dissociated tetrasubstituted organotin compounds which are produced by derivatization

3.4

solid

soil, sediment, sludge and waste (soil-like material)

4 Principle

For the ionic and the non-ionic organotin compounds (see [Table 1](#)), a different sample pretreatment and sample preparation are necessary. For the determination of organotin cations, laboratory samples are pretreated by freeze drying and grinding. This procedure enables to achieve homogeneity of the sample to be achieved. The determination of non-ionic TTBT cannot be carried out with freeze-dried materials due to evaporation losses; thus, it shall be determined in the field-moist sample. Organotin cations can only be determined after derivatization, whereas TTBT is already peralkylated and can be determined without derivatization (see the flowchart in [Figure 1](#)).




Figure 1 — Flowchart for the pretreatment and analysis of selected organotin compounds

For the determination of organotin compounds, two alternative extraction methods are given, both followed by *in situ* derivatization with a tetraethylborate compound and simultaneous extraction with hexane:

- a) treatment with acetic acid;
- b) treatment with methanolic potassium hydroxide.

Treatment with potassium hydroxide provides some degree of digestion and is recommended especially when the solid contains high amounts of organic and biological materials.

NOTE 1 If it is necessary to take a large amount of sample, extraction and derivatization can be done in two steps. An aliquot of the extract can be taken for derivatization. This also applies for samples with high levels of contamination by organotin compounds.

ISO/FDIS 23161:2016(E)

NOTE 2 During *in situ* derivatization, the solid phase is still present. This supports the extraction by continuous changing of the polar organotin cations to the non-polar organotin cation derivatives. *In situ* methods can improve the extraction efficiency, particularly for monoalkylated organotin compounds.

NOTE 3 Other extraction techniques can be applied if a comparable extraction efficiency is achieved.

When applying this method to the determination of other organotin compounds not specified in the scope, its suitability has to be proven by proper in-house validation experiments, e.g. methyltin compounds (see Table 2). Methyltin cations are unlikely to evaporate from aqueous solvents, but peralkylated methyltin compounds are volatile and subject to losses (see C.3). Therefore, additional precautions are established.

Table 2 — Methyltin compounds

$R_n\text{Sn}^{(4-n)+}$	R	n	Name	Acronym
MeSn^{3+}	Methyl	1	Monomethyltin cation	MMT
$\text{Me}_2\text{Sn}^{2+}$	Methyl	2	Dimethyltin cation	DMT
Me_3Sn^+	Methyl	3	Trimethyltin cation	TMT

The internal standard mix comprises four compounds representing four alkylation states in order to mimic the behaviour of the target compounds. After alkylation, they cover a wide range of volatility. A recovery of at least 80 % for derivatization/extraction and again 80 % for each clean-up step of the internal standard compounds should be achieved. (For more information, see A.3.) Tetraalkylborate is very reactive and will also alkylate other compounds in the matrix. Those compounds (and also boroxines) may interfere with the target compounds during gas chromatographic determination and influence detection. In order to protect the column and to reduce the interference in chromatography, it will be necessary to apply a pre-cleaning step to most samples. Clean-up with silica or aluminium oxide is the minimum; further clean-up steps (e.g. aluminium oxide/silver nitrate, silica/silver nitrate, pyrogenic copper; see Annex B) may be applied if necessary.

The determination of the tetrasubstituted organotin compounds is carried out after clean-up and concentration steps by separation with capillary gas chromatography and detected with a suitable system [mass spectrometer (MS), (MS/MS), flame photometric detector (FPD), atomic absorption spectrometer (AAS), atomic emission detector (AED), inductively coupled plasma/mass spectrometer ICP/MS]. The concentrations are determined by calibration over the total procedure using aqueous multi-component calibration standard solutions in accordance with 5.4.3.

5 Reagents

5.1 General

Use reagents of highest purity, typically of pesticide grade or better. The reagents may contain impurities of organotin compounds. It is absolutely essential to verify the blanks.

The water shall be free of interferences. Use water in accordance with Grade 3 of ISO 3696.

5.2 Chemicals

5.2.1 Acetic acid, CH_3COOH , glacial.

5.2.2 Sodium hydroxide solution, NaOH , 40 % (m/V).

5.2.3 Sodium acetate, CH_3COONa .

5.2.4 Sodium sulfate, Na_2SO_4 , anhydrous.

5.2.5 Potassium hydroxide, KOH.

5.2.6 Silica gel, grain size 0,085 mm to 0,28 mm (63 mesh to 200 mesh).

5.2.7 Aluminium oxide, Al₂O₃, alkaline.

5.2.8 Tetrahydrofuran, C₄H₈O, free of peroxides, free of water.

5.2.9 Acetone, (CH₃)₂CO.

5.2.10 Hexane, C₆H₁₄.

NOTE Both *n*-hexane and 2-methylpentane (*i*-hexane) have been found to be suitable.

5.2.11 Tetraethylborate compound, e.g sodium tetraethylborate, NaB(C₂H₅)₄.

NOTE The active species during derivatization is the tetraethylborate anion. The choice of the cation is arbitrary. Sodium tetraethylborate was chosen since it is commercially available. In principle, any other tetraethylborate compound can be used for analysis, including complexes formed with tetrahydrofuran (THF). A simple and rapid synthesis of a suitable derivatization agent is described in [A.1](#).

WARNING — Sodium tetraethylborate may contain traces of triethylboron, which may cause instantaneous combustion.

5.2.12 Methanol, CH₃OH.

5.2.13 Dichloromethane, CH₂Cl₂.

5.3 Standards <https://standards.iteh.ai/catalog/standards/sist/b5526690-521c-4c22-9baa-e7a267c0801b/osist-pren-iso-23161-2017>

WARNING — Organotin compounds vary largely regarding toxicological properties towards mammals with respect to the alkylation stage and type of alkyl group. Cautious handling of reagents is mandatory at any time.

[Table 3](#) lists the standards used for calibration of the target compounds (solution A), internal standards (solution B) and injection standard (solution C). Additional information is provided concerning weighing factors for calculation to organotin cations (for 100 % purity of the substances).

Table 3 — Standards and internal standards for calibration of target compounds

No.	Standard	Abbreviation	Formula	CAS-RN ^a	WF ^b	Solution ^c
5.3.1	Monobutyltin trichloride	MBTCl	C ₄ H ₉ SnCl ₃	1118-46-3	0,623	A
5.3.2	Dibutyltin dichloride	DBTCl	(C ₄ H ₉) ₂ SnCl ₂	683-18-1	0,767	A
5.3.3	Tributyltin chloride	TBTCl	(C ₄ H ₉) ₃ SnCl	1461-22-9	0,891	A
5.3.4	Tetrabutyltin	TTBT	(C ₄ H ₉) ₄ Sn	1461-25-2	1,000	A
5.3.5	Monooctyltin trichloride	MOTCl	C ₈ H ₁₇ SnCl ₃	3091-25-6	0,686	A
5.3.6	Diocetyl tin dichloride	DOTCl	(C ₈ H ₁₇) ₂ SnCl ₂	3542-36-7	0,830	A
5.3.7	Triphenyltin chloride	TPhTCl	(C ₆ H ₅) ₃ SnCl	639-58-7	0,908	A

^a Chemical Abstracts Registration Number.

^b WF= Weighing factor = Molar mass of organotin cation/molar mass of organotin compound.

^c A for the multicomponent-standard solution in methanol.

B for the solution of the internal standards in methanol.

C for the solution of the injection standards in hexane.