



SLOVENSKI STANDARD
oSIST prEN ISO 15952:2017
01-junij-2017

Kakovost tal - Vpliv onesnaževal na juvenilne (mladostniške) stadije kopenskih polžev (Helicidae) - Ugotavljanje vplivov na rast zaradi onesnaženja tal (ISO/DIS 15952:2017)

Soil quality - Effects of pollutants on juvenile land snails (Helicidae) - Determination of the effects on growth by soil contamination (ISO/DIS 15952:2017)

Bodenbeschaffenheit - Wirkungen von Schadstoffen auf Jungtiere von Landschnecken - Bestimmung der Wirkungen auf das Wachstum durch Bodenverunreinigung (ISO/DIS 15952:2017)

Qualité du sol - Effets des polluants vis-à-vis des escargots juvéniles (Helicidae) - Détermination des effets sur la croissance par contamination du sol (ISO/DIS 15952:2017)

Ta slovenski standard je istoveten z: prEN ISO 15952

ICS:

13.080.30 Biološke lastnosti tal Biological properties of soils

oSIST prEN ISO 15952:2017 **en,fr,de**

DRAFT INTERNATIONAL STANDARD

ISO/DIS 15952

ISO/TC 190/SC 4

Secretariat: AFNOR

Voting begins on:
2017-04-10Voting terminates on:
2017-07-02

Soil quality — Effects of pollutants on juvenile land snails (Helicidae) — Determination of the effects on growth by soil contamination

Qualité du sol — Effets des polluants vis-à-vis des escargots juvéniles (Helicidae) — Détermination des effets sur la croissance par contamination du sol

ICS: 13.080.30

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[SIST EN ISO 15952:2018](https://standards.iteh.ai/catalog/standards/sist/642d2cc1-1c67-4df9-a54d-aa474577b247/sist-en-iso-15952-2018)

<https://standards.iteh.ai/catalog/standards/sist/642d2cc1-1c67-4df9-a54d-aa474577b247/sist-en-iso-15952-2018>

THIS DOCUMENT IS A DRAFT CIRCULATED FOR COMMENT AND APPROVAL. IT IS THEREFORE SUBJECT TO CHANGE AND MAY NOT BE REFERRED TO AS AN INTERNATIONAL STANDARD UNTIL PUBLISHED AS SUCH.

IN ADDITION TO THEIR EVALUATION AS BEING ACCEPTABLE FOR INDUSTRIAL, TECHNOLOGICAL, COMMERCIAL AND USER PURPOSES, DRAFT INTERNATIONAL STANDARDS MAY ON OCCASION HAVE TO BE CONSIDERED IN THE LIGHT OF THEIR POTENTIAL TO BECOME STANDARDS TO WHICH REFERENCE MAY BE MADE IN NATIONAL REGULATIONS.

RECIPIENTS OF THIS DRAFT ARE INVITED TO SUBMIT, WITH THEIR COMMENTS, NOTIFICATION OF ANY RELEVANT PATENT RIGHTS OF WHICH THEY ARE AWARE AND TO PROVIDE SUPPORTING DOCUMENTATION.

This document is circulated as received from the committee secretariat.

ISO/CEN PARALLEL PROCESSING



Reference number
ISO/DIS 15952:2017(E)

© ISO 2017

iTeh STANDARD PREVIEW (standards.iteh.ai)

SIST EN ISO 15952:2018

<https://standards.iteh.ai/catalog/standards/sist/642d2cc1-1c67-4df9-a54d-aa474577b247/sist-en-iso-15952-2018>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2017, Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Ch. de Blandonnet 8 • CP 401
CH-1214 Vernier, Geneva, Switzerland
Tel. +41 22 749 01 11
Fax +41 22 749 09 47
copyright@iso.org
www.iso.org

Contents

| | Page |
|---|---|
| 1 | Scope 1 |
| 2 | Normative references 1 |
| 3 | Terms and definitions 2 |
| 4 | Principle 3 |
| 5 | Test environment..... 3 |
| 6 | Reagents 3 |
| 7 | Apparatus..... 5 |
| 8 | Storage and preparation of the samples 5 |
| 9 | Procedure 6 |
| 10 | Reference substance 9 |
| 11 | Calculations and expression of results 10 |
| 12 | Validity of test 12 |
| 13 | Test report..... 13 |
| Annex A (normative) Static method..... 14 | |
| Annex B (informative) Breeding technique for snails 15 | |
| Annex C (informative) Example of composition of snail feed 20 | |
| Annex D (informative) Example of table of data 21 | |
| Annex E (informative) Example of results with <i>Helix aspersa aspersa</i> 22 | |
| Annex F (informative) Determination of the effects on growth by food contamination 25 | |
| Annex G (informative) Test performance with other snail species 29 | |

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

International Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare International Standards. Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights.

ISO 15952 was prepared by Technical Committee ISO/TC 190, *Soil quality*, Subcommittee SC 4, *Biological methods*.

This second/third/... edition cancels and replaces the first/second/... edition (ISO 15952:2006), [clause(s) / subclause(s) / table(s) / figure(s) / annex(es)] of which [has / have] been technically revised.

(standards.iteh.ai)

SIST EN ISO 15952:2018

<https://standards.iteh.ai/catalog/standards/sist/642d2cc1-1c67-4df9-a54d-aa474577b247/sist-en-iso-15952-2018>

Introduction

Because of the limited amount of data available concerning toxicity of contaminants on soil organisms, the problems of assessing the ecotoxicity of soils and waste are cause for serious concern at both national and international levels. Currently available tests use soil-fauna organisms restricted to annelid (earthworms and *Enchytraeidae*) and arthropod phyla (insects: Collembola and Coleoptera). Among the latter, two standards assess acute toxicity [earthworms (ISO 11268-1) and coleoptera larvae ^[5]] and three other standards assess sublethal effects of soil contaminants on reproduction (earthworms ^[2], Collembola ^[1], *Enchytraeidae* ^[3]). In the biological cycles of organisms, it appears that growth is, like reproduction, a fundamental ecophysiological parameter to be taken into consideration for the sustainability of species and ecosystems ^[37].

Snails are relevant ecological indicators for assessing the quality of soils ^[15, 17 to 19, 31, 32, 39 to 41], as they are characteristic of the soil surface layer (saprophagous and phytophagous) of which a large part of the biological cycle takes place in the soil (egg-laying, hatching, initial stages of development, hibernation, etc.) ^{[6], [17], [28]}. During the other phases of their cycle, they eat soil and are in contact with the soil via their moist pedal sole (foot) covered with mucus and participate in the permanent exchanges with the soil (water, mineral salts, excrement and finally shell and organic matter when they die) ^{[6], [17], [30]}. In addition, they constitute an important link between plants, fauna and soil microorganisms. They correspond fully to the criteria for a good biological indicator: easy to sample and identify, they are widely distributed; they accumulate contaminants ^[8, 10 to 14, 16, 17, 21, 23, 28, 29, 32 to 47]; their ecological and physiological characteristics are well-known ^{[6], [9], [31]}; and they are now easy to breed under controlled conditions ^{[21], [25], [31]}. Their susceptibility to common contaminants of their environment has been demonstrated ^[10 to 15, 18 to 27, 29, 32 to 37, 36 to 47].

This International Standard describes a method for determining the effects on survival and growth of young snails of substances, preparations, soils or waste materials added to an artificial or a natural soil. The described method is thus applicable to test contaminated soils or to compare different uncontaminated soils. The recommended species is *Helix aspersa aspersa* Müller (also commonly called: common garden snail, brown garden snail, garden snail, land snail, "Petit-Gris"; synonyms: *Cantareus aspersus*, *Cornu aspersum*^[55]). Among land snails (stylommatophoran pulmonate gastropod molluscs of the *Helicidae* family), *Helix aspersa aspersa* Müller is the most ubiquitous. This palearctic species can be acclimated to regions with different types of climate: Mediterranean, oceanic temperate, midcontinental temperate and even tropical. *Helix aspersa aspersa* Müller is of European origin and has been introduced into all parts of the world. They are now on all continents except Antarctica ^[9].

Indeed, in their natural environment, snails integrate the contaminants by contact (with various substrates such as soil, soil leachates, plant litter), by ingestion (of plants and soil), as well as through the respiratory tract ^{[6], [28]}. So, for specific testing purposes (evaluation of the toxicity of a pesticide, for example), another test design, which is focussed on exposure via food uptake, is optionally available (Annex F and Reference [5]).

Soil quality — Effects of pollutants on juvenile land snails (Helicidae) — Determination of the effects on growth by soil contamination

1 Scope

This International Standard specifies a semi-static method for determining the effects of contaminants on growth and survival of young snails, usually *Helix aspersa aspersa* Müller. The animals are exposed via the cutaneous and digestive route using a test substrate (artificial or natural soil according to the objective of the study) to which defined amounts of the following are added:

- substances, mixtures or preparations;
- soils (contaminated or of unknown quality) or waste materials.

This test takes into account the possible changes in the test substance, preparation, soil or waste material because the test mixtures are prepared and renewed every week during the 28-day test period.

A static method may be implemented in addition to the semi-static method (optional). This method is described in Annex A.

This method does not apply to substances for which the air/soil partition coefficient is greater than one, or to substances with vapour pressure exceeding 300 Pa, at 25°C.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 10390, *Soil quality — Determination of pH*

ISO 18400-206, *Soil quality - Sampling — Part 206: Guidance on the collection, handling and storage of soil for the assessment of biological functional and structural endpoints in the laboratory*

ISO 10694, *Soil quality — Determination of organic and total carbon after dry combustion (elementary analysis)*

ISO 11260, *Soil quality — Determination of effective cation exchange capacity and base saturation level using barium chloride solution*

ISO 11268-1, *Soil quality — Effects of pollutants on earthworms — Part 1: Determination of acute toxicity to Eisenia fetida/Eisenia andrei*

ISO 11269-2, *Soil quality — Determination of the effects of pollutants on soil flora — Part 2: Effects of contaminated soil on the emergence and early growth of higher plants*

ISO 11274, *Soil quality — Determination of the water-retention characteristic — Laboratory methods*

ISO 11465, *Soil quality — Determination of dry matter and water content on a mass basis — Gravimetric method*

EN 14735, *Characterization of waste — Preparation of waste samples for ecotoxicity tests*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

- 3.1**
test substrate
artificial soil or natural soil used as control and dilution substrate
- 3.2**
matrix
soil or waste material under test
- 3.3**
test mixture
mixture of the test substance, preparation or matrix with the test substrate
- 3.4**
growth
increase in the biomass, i.e. in the total fresh mass (body and shell) of the organisms and increase in the maximum shell diameter, between the start and completion of the test
- Note 1 to entry: It is expressed in the form of a growth coefficient.
- 3.5**
effect concentration
EC_x
concentration at which a specific effect is detected; x is the percentage (10, 25, 50) of this effect, e.g. growth inhibition
- EXAMPLE EC₅₀ means the concentration estimated to reduce growth at the end of the test to 50 % compared to the control.
- 3.6**
median lethal concentration
LC₅₀
concentration of the substance, of the test preparation initially present, or the concentration of the matrix causing the death of 50 % of the snails submitted to testing
- 3.7**
lowest observed effect concentration
LOEC
lowest tested concentration at which the test substance is observed to have a statistically significant effect ($p < 0,05$) when compared with the control
- Note 2 to entry: All test concentrations above the LOEC have a harmful effect equal to or greater than those observed at the LOEC. When these two conditions cannot be satisfied, a full explanation should be given for how the LOEC (and hence the NOEC) has been selected.
- 3.8**
no observed effect concentration
NOEC
test concentration immediately below the LOEC, which, when compared with the control, has no statistically significant effect ($p > 0,05$) within a given exposure time

Note 3 to entry: The NOEC is the concentration just below the LOEC.

Note 4 to entry: For 3.5, 3.6, 3.7 and 3.8, results are given:

— in dry mass of test substance or preparation per dry mass of the test substrate;

— in mass percentage of the tested matrix in the test mixture (expressed in dry mass).

4 Principle

Juvenile land snails (usually *Helix aspersa aspersa* Müller) are exposed during a period of 28 days to test mixtures containing the test substance, preparation or matrix at different concentrations. The test mixtures are freshly prepared and renewed every 7 days.

According to the objectives, the test mixtures may be prepared with artificial soil (6.3.2) or with a suitable natural soil (6.3.3).

The snails are fed during the test with uncontaminated food.

The effects on growth (fresh mass and shell diameter) and on survival are measured after 28 days of exposure (optionally, effects could be measured every 7 days during 28 days).

The results obtained during testing are compared with those of a control to determine the NOEC or LOEC and to allow the estimation of the concentration which reduces the growth of the snails by 50 % within 28 days with respect to the fresh mass [$EC_{50,m}$ (28 days)] and to the shell diameter [$EC_{50,d}$ (28 days)] or other values of EC_x .

If the concentrations selected result in lethal effects, the results obtained during testing are compared with those of a control and used for estimating the concentration which causes the death of 50 % of the snails [LC_{50} (28 days)].

For particular applications, various parameters (EC_x , NOEC, LOEC, LC_{50}) can be assessed (optional) after exposure periods lower than 28 days (7 days, 14 days or 21 days).

The test is conducted in two stages:

- a preliminary test intended to indicate both the non-observed effect concentration, NOEC, and the complete growth inhibition. The resulting dose-response relationship is important for the proper design of the definitive test;
- a definitive test specifying the concentrations which cause between 10 % and 90 % of growth inhibition. It is not necessary to perform a final test where the preliminary test has not revealed any inhibitory effects at the maximum concentration tested.

5 Test environment

The test shall be carried out at a temperature of (20 ± 2) °C under a day-night photoperiod of 18 h to 6 h. The illumination intensity (artificial light of daylight type), without any natural light in the test containers shall be 50 to 100 lux.

6 Reagents

6.1 Water, of purity at least deionized

6.2 Biological material

Test organisms shall be juvenile snails. The recommended species is *Helix aspersa aspersa* Müller (also known as *Cantareus aspersus* and *Cornu aspersum*) which shall be 3 to 5 weeks old, having a mean fresh mass of $(1 \pm 0,3)$ g and a shell diameter of $(15,5 \pm 1)$ mm.

NOTE The use of some other genus and/or species of *Helicidae* is possible (see examples and conditions in Annex G).

The snails shall be selected from synchronous breeding in order to form a population as homogeneous as possible with respect to size, mass and age. The breeding techniques for snails are described in Annex B. After a nursery

ISO/DIS 15952

period (3 to 5 weeks, see Annex B), the young snails shall be used after at least 1 week of aestivation and no more than 5 months. The aestivation is carried out in round wooden boxes (approximately 12 cm in diameter and 4 cm in height), with the snails under dry conditions, at a temperature of 17 °C to 20 °C.

Two to three days before starting the test, snails shall be woken by spraying water (6.1) into the boxes used for aestivation. The proportion of snails not woken shall be less than 10 %. As soon as they have resumed activity (snails not stuck to the walls of the box and which are beginning to move about), the snails shall be transferred to a box (7.1) that has been moistened with water (6.1). The bottom of this box either can be covered with absorbent paper that has also been moistened, or can contain some test substrate (6.3) moistened to 50 % to 60 % of its water-holding capacity. Between waking and the start of the test (2 to 3 days), the snails shall be fed (6.4).

6.3 Test substrate

6.3.1 General

According to the objectives of the study, either an artificial soil (6.3.2) or a suitable natural soil (6.3.3) is used as test substrate.

NOTE Artificial soil may be used as a control and dilution substrate to assess the effect of a substance or of a preparation, or to compare different soils or waste, or to assess the effects of a contaminated soil.

Natural soil (field soil) may be used as a control and dilution substrate in order to assess, for example, the effect of the incorporation of wastewater treatment plant sludge into the field soil or to test the effect of a contaminated soil (in this case an uncontaminated soil comparable to the soil sample to be tested ought to be used).

6.3.2 Artificial soil

The artificial soil shall have the following composition (as defined by ISO 11268-1).

Table 1 — Composition of artificial soil

| Composition | Percentage expressed in dry mass |
|--|---|
| Sphagnum peat air-dried and finely ground (2 ± 1 mm without any visible plant remains). | 10 % |
| Kaolinite clay, preferably containing not less than 30 % kaolinite. | 20 % |
| Air-dried industrial quartz sand (predominantly fine sand with more than 50 % by mass of particle size 0,05 mm to 0,2 mm). | Approximately 69 % (depending on the amount of CaCO_3 needed). |
| Calcium carbonate (CaCO_3 , pulverised, analytical grade) to bring the pH of the wetted artificial soil to $6,0 \pm 0,5$. | Approximately 0,3 % to 1,0 % |

The artificial soil shall be prepared, at least two days prior to starting the test, by mixing the dry constituents listed above thoroughly in a large-scale laboratory mixer. The amount of calcium carbonate required might vary, depending on the properties of the individual batch (mainly the peat) and should be determined by measuring subsamples immediately before the test.

The mixed artificial soil shall be stored at room temperature for at least two days to equilibrate acidity. To determine pH and the maximum water-holding capacity, the dry artificial soil shall be pre moistened one or two days before starting the test by adding deionized water to obtain half of the required final water content of 50 % to 60 % of the maximum water-holding capacity.

The pH value shall be measured according to ISO 10390. If the measured pH is not within the required range, a sufficient amount of CaCO_3 shall be added or a new batch of artificial soil shall be prepared. The maximum water-holding capacity of the artificial soil shall be determined according to ISO 11274 or to Annex C of ISO 11269-2:2012.

6.3.3 Natural soil

Determine the following parameters on the selected natural soil which shall be sieved through a 4-mm square mesh sieve to remove large fragments:

- pH, according to ISO 10390;
- water-holding capacity, according to ISO 11274 or Annex C of ISO 11269-2:2012;
- water content, according to ISO 11465;
- content of organic matter, according to ISO 10694.

It is also recommended to determine the cation exchange capacity, according to ISO 11260.

6.4 Feed

The feed shall be provided in the form of flour at its natural moisture content (5 % to 10 %).

In order to obtain sufficient growth, it is recommended to carry out the tests with a flour-based feed comprising cereals, forage, mineral salts and vitamins which properly covers the needs of the snails¹. An example of feed composition is given in Annex C.

7 Apparatus

Use ordinary laboratory apparatus and the following.

7.1 Test containers

Disposable mouse boxes made of transparent polystyrene² or any other container having a volume of approximately 1,6 l [advised approximate dimensions: 24 cm (length) × 10,5 cm (width) × 8 cm (height)].

7.2 Feed containers

Petri dishes, approximately 5,5 cm in diameter and approximately 1 cm in height or any other containers of equivalent dimensions.

7.3 Calliper rule, having a precision of 0,1 mm

7.4 Balances

One analytical balance having a precision of at least 1 mg. Two other balances, one having a precision of 0,1 g, another having a precision of 1 g.

8 Storage and preparation of the samples

8.1 Soil to be tested

The soil samples received at the laboratory shall be stored in accordance with ISO 18400-206.

The soil sample submitted for testing shall be sieved through a 4-mm square mesh sieve to remove coarse fragments.

For each soil, the same characteristics than for natural soil (6.3.3) that can be used as control or dilution substrate, shall be determined.