

---

---

**Water quality — Larval development  
test with the harpacticoid copepod  
*Nitocra spinipes***

*Qualité de l'eau — Essai de développement larvaire avec le copépode  
harpacticoïde Nitocra spinipes*

**iTeh STANDARD PREVIEW**  
(standards.iteh.ai)  
Full standard:  
<https://standards.iteh.ai/catalog/standards/sist/0499e07-5299-48c7-a277-1c2f965f514/iso-ts-18220-2016>

**PROOF / ÉPREUVE**

---

---



**iTeh STANDARD PREVIEW**  
(standards.iteh.ai)  
Full standard:  
<https://standards.iteh.ai/catalog/standards/sist/0499eed7-5299-48c7-a277-1c2f965f514/iso-ts-18220-2016>



**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2016, Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Ch. de Blandonnet 8 • CP 401  
CH-1214 Vernier, Geneva, Switzerland  
Tel. +41 22 749 01 11  
Fax +41 22 749 09 47  
copyright@iso.org  
www.iso.org

# Contents

Page

<b>Foreword</b> .....	<b>iv</b>
<b>Introduction</b> .....	<b>v</b>
<b>1 Scope</b> .....	<b>1</b>
<b>2 Terms and definitions</b> .....	<b>1</b>
<b>3 Principle</b> .....	<b>2</b>
<b>4 Reagents</b> .....	<b>2</b>
<b>5 Cultivation</b> .....	<b>3</b>
5.1 Test organism.....	3
5.2 Algae for feeding.....	3
<b>6 Apparatus</b> .....	<b>3</b>
<b>7 Procedure</b> .....	<b>4</b>
7.1 Production of nauplii to be used in test.....	4
7.2 Choice of test concentrations.....	4
7.2.1 Hypothesis testing.....	5
7.2.2 Regression analysis.....	5
7.3 Preparation of solutions to be used in test.....	5
7.3.1 Stock solution.....	5
7.3.2 Test solutions.....	5
7.4 Start of test.....	6
7.5 Incubation/exposure.....	6
7.6 Maintenance.....	6
7.7 Measurements/observations.....	6
7.7.1 Mortality.....	6
7.7.2 Larval development ratio (LDR).....	7
7.7.3 Physical-chemical parameters — oxygen, pH and salinity.....	7
7.7.4 Concentration of the test substance.....	7
<b>8 Validity criteria</b> .....	<b>7</b>
<b>9 Evaluation of results</b> .....	<b>8</b>
9.1 Calculation of results.....	8
9.2 Expression of results.....	8
9.3 Interpretation of results.....	8
<b>10 Reproducibility</b> .....	<b>8</b>
<b>11 Test report</b> .....	<b>8</b>
<b>Annex A (informative) Biology and cultivation of <i>Nitocra spinipes</i></b> .....	<b>10</b>
<b>Annex B (informative) <i>Nitocra spinipes</i> larval development ratio</b> .....	<b>15</b>
<b>Annex C (informative) Calculations</b> .....	<b>17</b>
<b>Bibliography</b> .....	<b>19</b>

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see [www.iso.org/directives](http://www.iso.org/directives)).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see [www.iso.org/patents](http://www.iso.org/patents)).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see the following URL: [www.iso.org/iso/foreword.html](http://www.iso.org/iso/foreword.html).

The committee responsible for this document is ISO/TC 147, *Water quality*, Subcommittee SC 5, *Biological methods*.

## Introduction

Harpacticoid copepods are predominantly benthic, occurring widely in marine, brackish and fresh water ecosystems. They represent important prey items for the benthic larvae of many fish species and larger invertebrates and constitute an ecologically important energy-transfer link between the organic phase of the sediment and higher trophic levels.

The euryhaline brackish water harpacticoid *Nitocra spinipes* (Crustacea) is a common component of the benthic meiofauna in shallow coastal waters around the world (see Reference [6]).

**iTeh STANDARD PREVIEW**  
(standards.iteh.ai)  
Full standard:  
<https://standards.iteh.ai/catalog/standards/sist/0499eed7-5299-48c7-a277-1c2f965f514/iso-ts-18220-2016>

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

Full standard:  
<https://standards.iteh.ai/catalog/standards/sist/0499eed7-5299-48c7-a277-1c2f965f514/iso-ts-18220-2016>

# Water quality — Larval development test with the harpacticoid copepod *Nitocra spinipes*

**WARNING** — Persons using this Technical Specification should be familiar with normal laboratory practice. This Technical Specification does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices and to ensure compliance with any national regulatory conditions.

**IMPORTANT** — It is absolutely essential that tests conducted in accordance with this Technical Specification be carried out by suitably qualified staff.

## 1 Scope

This Technical Specification specifies an early-life stage procedure for determination of the toxic effects of chemicals and water samples on a cold-water brackish water copepod species under semi-static conditions. The biological test variables include survival and development of the early-life stages. The exposure starts with newly hatched (<24 h) nauplii (larvae) and is continued until emergence of (c. 50 %) copepodites (juveniles) in the control.

The benthic living *Nitocra* complements the planktonic *Acartia* species in ISO 16778. These organisms represent different life-history strategies as *Nitocra* is egg-carrying, whereas *Acartia* is a broadcasting calanoid copepod and thus, different sensitivities of specific life stages. *Nitocra* is a fresh to brackish water species, which allows testing low salinity waters and is complementary to *A. tonsa*, which is of marine origin and poorly tolerates low salinities.

## 2 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

### 2.1

#### **nauplii**

larvae

### 2.2

#### **copepodites**

juveniles

### 2.3

#### **larval development ratio**

##### **LDR**

ratio of *copepodites* (2.2) to the total number of surviving early-life stages (nauplii + copepodites) at the end of the test

### 2.4

#### **lowest observed effect concentration**

##### **LOEC**

lowest concentration within the experimental range at which a significant effect is observed

### 2.5

#### **no observed effect concentration**

##### **NOEC**

tested concentration just below the LOEC (2.4)

**2.6**  
**effect concentration**

$EC_x$   
calculated concentration from which an effect of x % is expected

**2.7**  
**mortality**

calculated on dead and missing animals at the end of the test divided by animals at start

**2.8**  
**confidence interval**

$A \times \%$   
interval of values within which the measured or calculated value is likely to be present with a probability of x %

**2.9**  
**salinity**

$S$   
dimensionless value of which, for the purpose of checking water quality, may be regarded as an estimate of the concentration, in grams per kilogram, of dissolved salts in sea water

Note 1 to entry: It is defined algorithmically, in terms of the ratio (K15) of the electrical conductivity of the sample, at 15 °C and 1 atm, to that of defined potassium chloride solution (32,436,6 g/kg of sample) at the same temperature and pressure.

### 3 Principle

The test is an early-life stage test, in which the organisms are exposed to various concentrations of a test substance or water sample under semi-static test conditions from the first naupliar stage (N-1) to the first copepodite stages (C-1, C-2, etc.). Survival and development of early-life stages [larval development ratio (LDR)], are the investigated test variables. The exposure starts with newly hatched (<24 h) nauplii (larvae) and is continued until the emergence of (approximately 50 %) copepodites (juveniles) in the control. The total test duration is about 6 d to 7 d, which is sufficient time to investigate the development from N-1 to 50 % copepodites in the control. The naupliar (larval) and copepodite (juvenile) stages are morphologically distinct, and therefore, the transition from the last naupliar to the first copepodite stage is easily observed.

The outcome of the test is either the no observed effect concentration (NOEC) and the lowest observed effect concentration (LOEC) values or the effect concentrations with a certain degree (x %) of inhibition ( $EC_x$ ) (e.g.  $EC_{50}$  and  $EC_{10}$ ).

### 4 Reagents

#### 4.1 Test organism

The species to be used is the brackish water harpacticoid copepod *Nitocra spinipes* Boeck.

Newly hatched (less than 24 h of age) nauplii should be collected from a healthy stock (i.e. showing no signs of stress, such as high mortality, poor fecundity, etc.). The stock animals shall be maintained in culture conditions (light, temperature, medium and feeding) similar to those to be used in the test (culturing method for *N. spinipes* is described in [Annex A](#)).



**4.2 Water**, deionized or of equivalent purity, to prepare artificial sea water or to dilute natural sea water.

#### 4.2.1 Artificial sea water

An example of artificial sea water suitable for cultivation and testing is included in [Annex A](#). Any artificial sea water with a known composition in which the copepods show suitable long-term survival, normal behaviour, development and fecundity may be used as culture and dilution medium ([4.3](#)).

#### 4.2.2 Natural sea water

Natural sea water shall be collected from an unpolluted location. Any natural sea water with a known composition in which the copepods show suitable long-term survival, normal behaviour, development and fecundity may be used as culture and dilution medium.

Suspended particles shall be <10 mg/l and can be stored cold for approximately 6 months before preparation of culture or dilution medium.

### 4.3 Medium

#### 4.3.1 Culture medium

Culture medium is used for cultivating *Nitocra spinipes* and is prepared from either natural or artificial sea water ([4.2](#)). Natural sea water shall be filtered (30 µm) and heated to 80 °C to kill undesired organisms and then conditioned (24 h) to culture temperature and oxygen saturation. The culture medium can be stored cold for several weeks.

#### 4.3.2 Dilution medium

Dilution medium is used for diluting water samples or dissolving chemicals and is prepared from culture medium that has first been filtered (GF/C-1,2 µm) before use. Salinity of the dilution medium should be the same as the culture medium. Salinities between 3 ‰ and 25 ‰ can be used. The dilution medium shall have a dissolved oxygen concentration above 70 % of the air saturation value and a pH of  $7,5 \pm 1,0$  before being used to prepare the test solutions. If the physical conditions or the salinity of the medium to be used in the test differ more than 5 °C or 10 % from those used for routine culturing, it is good practice to include an adequate cultivating acclimation period at the same salinity ( $\pm 2$  ‰) of 2 weeks to 3 weeks to avoid stressing of the larvae.

## 5 Cultivation

### 5.1 Test organism

See [Annex A](#) for detailed information.

### 5.2 Algae for feeding

See [Annex A](#) for detailed information.

## 6 Apparatus

All equipment, which will come in contact with the test medium, shall be made of glass or chemically inert material.

**6.1 Glass vessels**, approximately 150 ml, diameter 8 cm, and height 4,5 cm, for *Nitocra spinipes* cultivation.

- 6.2 **Test vessels**, approximately 15 ml, diameter 2,5 cm, height 4 cm, with flat bottom.
- 6.3 **pH meter**.
- 6.4 **Oxygen meter**.
- 6.5 **Conductivity meter**.
- 6.6 **Wide pipettes**, for sampling animals, preferably salinized to prevent copepods from adhering to pipette walls.
- 6.7 **Temperature-control cabinet or room**,  $(22 \pm 1) ^\circ\text{C}$ .
- 6.8 **Low-magnifying stereo microscope**.
- 6.9 **Inverted microscope**.
- 6.10 **Apparatus for membrane filtration**.
- 6.11 **Filters**, 1,2  $\mu\text{m}$  and 30  $\mu\text{m}$ .

## 7 Procedure

### 7.1 Production of nauplii to be used in test

Nauplii aged less than 24 h are used for initiating a test. To produce a sufficient number of nauplii, the procedure presented below should be followed.

The day before the test starts, approximately 300 females with well-developed egg sacs are sampled by pipette under a low-magnifying stereomicroscope and randomly transferred (approximately 60 in each) to six glass "hatch" vessels containing 100 ml dilution medium. The isolated female copepods are fed with a suspension of *Rhodomonas salina* to a concentration of  $2,5 \times 10^5$  cells/ml.

### 7.2 Choice of test concentrations

The range of the test concentrations should preferably not include any concentrations that have a significant effect on survival since the main objective of the test is to measure sublethal effects (i.e. development).

Prior knowledge of the toxicity of the test substance, i.e. from an acute test (see Reference [2]) or from range-finding studies, should help in selecting appropriate test concentrations. As a rule of thumb, the highest concentration in the early-life stage test should be set at 10 % to 20 % of the acute 96 h-LC<sub>50</sub> to avoid significant effect on survival.

At least five different concentrations should be tested in a geometric series with a factor between concentrations not exceeding 3,3. Justification should be provided if fewer than five concentrations are used. Substances should not be tested above their solubility limits in dilution medium. A dilution-medium control shall be included, and also, if relevant, a solvent-control containing the same concentration of solvent as the test series should be run additionally.

The number of replicates generally depends on the statistical design (hypothesis testing or regression analysis). When planning the test, it should be taken into consideration if the aim is to achieve a NOEC/LOEC (by use of Chi-square) or an EC<sub>x</sub> value (by use of regression analysis or none parametric alternative; see Reference [4]).