# INTERNATIONAL STANDARD

ISO 18763

First edition 2016-07-01

# Soil quality — Determination of the toxic effects of pollutants on germination and early growth of higher plants

Qualité du sol — Détermination des effets toxiques des polluants sur la germination et la croissance primaire des plantes supérieures

11en Standards

(https://standards.iteh.ai)
Document Preview

ISO 18763:2016

https://standards.iteh.ai/catalog/standards/iso/a7308480-a60e-46ba-a92e-e15f1e85d9c8/iso-18763-2016



# iTeh Standards (https://standards.iteh.ai) Document Preview

ISO 18763:2016

https://standards.iteh.ai/catalog/standards/iso/a7308480-a60e-46ba-a92e-e15fle85d9c8/iso-18763-2016



## COPYRIGHT PROTECTED DOCUMENT

© ISO 2016, Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office Ch. de Blandonnet 8 • CP 401 CH-1214 Vernier, Geneva, Switzerland Tel. +41 22 749 01 11 Fax +41 22 749 09 47 copyright@iso.org www.iso.org

itents	Page
word	iv
duction	<b>v</b>
Scope	1
Normative references	1
Terms and definitions	1
Principle	3
Reagents, test organisms and media	3
Apparatus and materials	4
Treatment and preparation of samples	6
Procedure	6
Measurement	8
Calculation of the percentage inhibition	9
Reference chemical	10
Precision	10
Validity criteria	11
Test report.	11
x A (informative) Application of the phytotoxicity test in transparent test plates on natural and artificial soils and soil materials, with different plant species	12
x B (informative) Assemblage of test plates for the phytotoxicity test	14
x C (informative) International interlaboratory comparisons on the phytotoxicity test	15
ography ISO 18763:2016	20
	word duction Scope Normative references Terms and definitions Principle Reagents, test organisms and media Apparatus and materials Treatment and preparation of samples Procedure Measurement Calculation of the percentage inhibition Reference chemical Precision Validity criteria Test report x A (informative) Application of the phytotoxicity test in transparent test plates on natural and artificial soils and soil materials, with different plant species x B (informative) Assemblage of test plates for the phytotoxicity test x C (informative) International interlaboratory comparisons on the phytotoxicity test

# Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see <a href="www.iso.org/directives">www.iso.org/directives</a>).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see <a href="https://www.iso.org/patents">www.iso.org/patents</a>).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the WTO principles in the Technical Barriers to Trade (TBT) see the following URL: Foreword - Supplementary information

The committee responsible for this document is ISO/TC 190, *Soil quality*, Subcommittee SC 4, *Biological methods*.

ISO 18763-2016

https://standards.iteh.ai/catalog/standards/iso/a7308480-a60e-46ba-a92e-e15fle85d9c8/iso-18763-2016

# Introduction

Ecotoxicological testing of test soils or waste materials to be disposed on soil are required to assess the potential environmental risk resulting from soil pollution or the disposal of wastes such as sewage sludge on farmland. There is also a need to monitor the quality of soil after reclamation of industrial sites. Therefore, a very practical and rapid germination and growth test has been developed based on seed germination and seedling growth in controlled environmental conditions.

The assay, which does not require any pretreatment of the seeds, is performed in "transparent test plates", incubated vertically, to allow the roots and the shoots of the germinated seeds to be seen. After 72 h exposure, a picture of the transparent test plates is taken and can be analysed "by image analysis" for multiple endpoints, such as percentage of seed germination and of length of roots and shoots. To account for the plant species variability in sensitivity, the assays are performed with the seeds of three plant species: one monocotyl (*Sorghum saccharatum*) and two dicotyls (*Lepidium sativum* and *Sinapis alba*).

A major advantage of this test is that after the shooting and storing of the pictures of the test plates, the measurements by image analysis can be postponed to any appropriate timing.

Reference or standard soils can be used as negative controls, such as, for example, the ISO standard artificial soil according to ISO 11269-1 and ISO 11269-2.

Commercially available seeds, with a shelf life longer than one year, allow the use of this test at any time of the year.

Two International interlaboratory comparisons demonstrated that the test provides good results.

A substantial number of studies report data on the application of this test on various types of soils and soil materials with several types of plant species.

# **Document Preview**

ISO 18763:2016

https://standards.iteh.ai/catalog/standards/iso/a7308480-a60e-46ha-a92e-e15fle85d9c8/iso-18763-2016

# iTeh Standards (https://standards.iteh.ai) Document Preview

ISO 18763:2016

https://standards.iteh.ai/catalog/standards/iso/a7308480-a60e-46ba-a92e-e15fle85d9c8/iso-18763-2016

# Soil quality — Determination of the toxic effects of pollutants on germination and early growth of higher plants

# 1 Scope

This International Standard describes a technique for determining the effects of soil and soil-related materials on the seed germination and early growth of higher plants. These endpoints are useful indicators for the assessment of the quality of a soil as a habitat for organisms. This International Standard is applicable to all soils in which soil organisms are active and may be used to evaluate:

- the effects on plants due to toxicity of solid or liquid chemicals contaminating soil or materials (compost, sludge, waste) and chemicals added to soil;
- the changes in the soil effect on plants after restoration measures.

# 2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 11269-1, Soil quality — Determination of the effects of pollutants on soil flora — Part 1: Method for the measurement of inhibition of root growth

ISO 11269-2, Soil quality — Determination of the effects of pollutants on soil flora — Part 2: Effects of contaminated soil on the emergence and early growth of higher plants

ISO/TS 20281, Water quality — Guidance on statistical interpretation of ecotoxicity data

https://standards.iteh.ai/catalog/standards/iso/a7308480-a60e-46ba-a92e-e15fle85d9c8/iso-18763-2016

## 3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

# 3.1

#### artificial soil

mixture of sand, kaolinite, peat and calcium carbonate prepared according to ISO 11269-1 and ISO 11269-2

#### 3.2

#### control soil

reference or standard soil used as a control and as a medium for preparing dilution series with test soils or a reference substance

# 3.3

#### reference soil

uncontaminated site-specific soil (e.g. collected in the vicinity of a contaminated site) with similar properties (nutrient concentrations, pH, organic carbon content and texture) as the test soil

#### 3.4

#### standard soil

field-collected soil or artificial soil whose main properties (e.g. pH, texture, organic matter content) are within a known range

Note 1 to entry: The properties of standard soils can differ from the test soil.

# ISO 18763:2016(E)

EXAMPLE Euro-soils[1], artificial soil[2], LUFA soil.1)

#### 3.5

#### test soil

either a natural or an artificial clean soil that is spiked with the test substance or a contaminated natural soil (a site soil)<sup>[5]</sup>

#### 3.6

#### seeding emergence

appearance of a visible seedling above the surface of the cover material

[SOURCE: ISO 17126:2005, 3.1, modified]

#### 3.7

# germination

appearance of a root of at least 1 mm of length

#### 3.8

# pure water

grade of water, produced, for example, by single distillation, by de-ionization, by ultra-filtration or by reverse osmosis[5]

#### 3.9

# root length

length of the root from seed to root tip

# 3.10

# shoot length

length of the part that grows upward, from seed to tip 2 10 S.11eh.21

#### 3.11

#### water saturation

maximum water content that a soil can retain against gravity under undisturbed soil conditions, conventionally stated as water content two days to three days after full saturation with water

[SOURCE: ISO 11074:2015, 2.1.5 field capacity, modified] -a60e-46ba-a92e-e15fle85d9c8/iso-18763-2016

#### 3.12

#### water saturated soil

soil which has reached its maximum water content

#### 3.13

## water-holding capacity

mass of water that evaporates from soil saturated with water when the soil is dried to constant mass at 105 °C, divided by the dry mass of the soil[ $\[ \]$ ]

## 3.14

## negative control

any well-characterized material or substance that, when tested by a specific procedure, demonstrates the suitability of the procedure to yield a reproducible, appropriately negative, non-reactive or minimal response in the test system

[SOURCE: ISO 10993-10:2010, 3.12, modified]

# 3.15

#### effect percentage

percentage decrease of the seed germination and the growth of the plant roots and/or shoots in the test soil in comparison to the control soil

<sup>1)</sup> Euro-soils, artificial soil and LUFA soil are examples of suitable products available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of these products.

# 4 Principle

This method compares the seed germination and early growth of monocotyledonous and dicotyledonous plants in a test soil and/or a series of mixtures with a control soil. This method may also be used for the testing of compost, sludge or waste.

Seeds of one monocotyledonous plant, such as *Sorghum saccharatum* (L.) Moench, and two dicotyledonous plants, such as *Lepidium sativum* L. and *Sinapis alba* L., are exposed to the test material under controlled conditions. After  $(72 \pm 1)$  h, the number of germinated seeds is recorded and the length of the roots of the test plants is measured in the test soil and in the control soil.

If different seed species are used, the length of the incubation period may be adjusted, depending on the time of germination of the seeds and the growth speed of the roots.

The test makes use of unique flat and shallow transparent test plates (6.3) composed of two compartments, the lower one of which contains hydrated soil.

Seeds of the selected test plants are positioned at equal distance near the middle ridge of the test plate (6.3) on a black filter paper (6.5) placed on top of the hydrated soil.

After closing the test plates (6.3) with their transparent cover, the test plates are placed vertically in a holder (6.4) and incubated at (25  $\pm$  1) °C for (72  $\pm$  1) h.

At the end of the incubation period, the length of each root (and shoot, if wished) can be measured directly with a ruler and recorded.

Alternatively, a "digital" picture is taken of the test plates (6.3) with the germinated plants (either with a digital camera, a webcam camera or a flatbed paper scanner) for storage in a computer file. The subsequent root length measurements are performed by image analysis. The analyses on germination and root growth can then be made immediately or postponed to any appropriate timing.

NOTE The same procedure can be applied to also measure the shoot height, if wished. Calculation of the shoot/root length ratio is a possible additional effect parameter.

#### ISO 18763:2016

# 5 // Reagents, test organisms and media - a60e-46ba-a92e-e15f1e85d9e8/iso-18763-2016

#### **5.1** Water.

Pure water having a conductivity below 10 μS/cm.

# 5.2 Test organisms.

The test organisms are seeds of one monocotyledonous plant, such as *Sorghum saccharatum* (L.) Moench, and two dicotyledonous plants, such as *Lepidium sativum* L. and *Sinapis alba* L.

Investigations have been performed not only with the three plant species indicated in <u>5.2</u> but also with other monocotyl and dicotyl plant species. A synthesis on these published studies is given in <u>Annex A</u>.

Seeds coated with insecticides and/or fungicides should be avoided.

#### 5.3 Control soil.

Either reference or standard soils can be used as the control soil, if unhindered growth of the test plants in these soils can be expected.

When comparing the root elongation in soils of known and unknown quality, the control soil and soil under test should be of the same textural class, and be as similar as practicable in all respects other than the presence of the chemical or contaminant being investigated. Indeed, significant differences in soil characteristics other than the presence of contaminant may lead to differences in root lengths and may induce false positive test results.

Alternatively, artificial soil according to ISO 11269-1 and ISO 11269-2 may be used. The substrate called "artificial soil" has the following composition:

	Percentage expressed on dry-mass basis
Sphagnum peat finely ground and with no visible plant remains	10 %
Kaolinite clay containing not less than 30 % kaolinite	20 %
Industrial quartz sand (dominant fine sand with more than 50 $\%$ of particle size 0,05 mm to 0,2 mm)	69 %

NOTE As indicated in ISO 11269-1, 5 % peat have proven to be sufficient for maintaining the desired structure of the artificial soil (with a corresponding increase of the sand percentage to 75 %). A lower percentage of kaolinite clay (10 % instead of 20 %) is furthermore very close to the clay content of LUFA 2.2 and hence more representative for a natural soil. The following composition of the artificial soil is therefore recommended: peat 5 %, kaolinite clay 10 % and sand 85 %.

Approximately 0,3 % to 1,0 % calcium carbonate (CaCO3, pulverized, analytical grade) is necessary to get a pH of 6,0  $\pm$  0,5.

# 6 Apparatus and materials

- **6.1 Incubator or temperature controlled room**, suitable for maintaining the specified conditions at  $(25 \pm 1)$  °C.
- **6.2 Digital camera, webcam camera or flatbed paper scanner**, to shoot pictures of the test plates with the germinated seeds, for storage in a computer file.
- **6.3 Test plates**, transparent plates in polyvinylchloride (PVC).

The test plates are composed of a bottom part separated by a middle ridge into an upper part and a lower part and a flat cover. Test plates can be "handmade" with the aid of transparent PVC sheet and small rectangular sticks as described in <u>Annex B</u>.

Alternatively, commercially available<sup>2)</sup> test plates can be used. The lower part of these test plates is intended to hold approximately 90 cm<sup>3</sup> of test soil. The latter test plates have on both parts on their side small rectangular cavities for closing the plates tightly by a unique click system. The test plates shall be provided with a label to record the specifics of each test plate (type of soil, type of seed, number of the replicate).

-

<sup>2)</sup> The test plates supplied by MicroBioTests Inc. Mariakerke-Gent, Belgium are an example of a suitable product available commercially. This information is given for the convenience of users of this International Standard and does not constitute an endorsement by ISO of this product.




Figure 1 — Test plate with cavities on the side for tight closing of the plate

- **6.4 Test plate holders**, cardboard holders for vertical incubation of six test plates (<u>6.3</u>) each.
- **6.5 Black filter papers**, rectangular high purity black filter papers (e.g. 85 g/m², 0,17 mm thickness, 45 s filtration speed) fitting the lower part of the test plate, to be placed on top of the soil in the lower compartment of the test plates (6.3).
- **6.6 Microsieve cylinder**, small plastic cylinder provided at the bottom with a nylon gauze, to be used for determination of the water to be added to the test soil.
- **6.7 Wide mouth micropipette**, plastic micropipette to be used with the microsieve cylinder (<u>6.6</u>) for determination of the water to be added to the test soil.
- **6.8 Sieve**, sieve of 2 mm mesh for sieving the test soil prior to use for the tests.
- **6.9 Thin spatula**, hand tool with a thin blade used to mix the test soil with water.
- **6.10** Flat spatula, hand tool with a broad, flat blade that is used to spread and flatten the test soil into the lower compartment of the test plates (6.3).
- **6.11 Tweezers**, small pincer-like tool for handling the seeds.