

Second edition
2017-06

Corrected version
2018-06

**Microbiology of the food chain —
Horizontal method for the
detection and enumeration of
Enterobacteriaceae —**

**Part 2:
Colony-count technique**

iTeh STANDARD PREVIEW
(standards.iteh.ai)

*Microbiologie de la chaîne alimentaire — Méthode horizontale
pour la recherche et le dénombrement des Enterobacteriaceae —*

Partie 2: Technique par comptage des colonies

<https://standards.iteh.ai/catalog/standards/sist/c8ce5547-a3ed-4128-a2e3-d9acca126a8e/iso-21528-2-2017>



Reference number
ISO 21528-2:2017(E)

© ISO 2017

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 21528-2:2017

<https://standards.iteh.ai/catalog/standards/sist/c8ce5547-a3ed-4128-a2e3-d9acca126a8e/iso-21528-2-2017>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2017

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Fax: +41 22 749 09 47
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

Contents

| | Page |
|--|-----------|
| Foreword | iv |
| Introduction | v |
| 1 Scope | 1 |
| 2 Normative references | 1 |
| 3 Terms and definitions | 1 |
| 4 Principle | 2 |
| 4.1 Preparation of initial suspension and decimal dilutions..... | 2 |
| 4.2 Isolation and selection for confirmation..... | 2 |
| 4.3 Confirmation..... | 2 |
| 4.4 Calculation..... | 2 |
| 5 Diluent, culture media and reagent | 2 |
| 6 Equipment and consumables | 2 |
| 7 Sampling | 3 |
| 8 Preparation of test sample | 3 |
| 9 Procedure | 3 |
| 9.1 General..... | 3 |
| 9.2 Test portion, initial suspension and dilutions..... | 4 |
| 9.3 Inoculation and incubation..... | 4 |
| 9.4 Counting and selection of colonies for confirmation..... | 4 |
| 9.5 Subculturing selected colonies..... | 4 |
| 9.6 Biochemical confirmation tests..... | 5 |
| 9.6.1 Oxidase reaction..... | 5 |
| 9.6.2 Fermentation test..... | 5 |
| 9.6.3 Interpretation of biochemical tests..... | 5 |
| 10 Expression of results | 5 |
| 11 Performance characteristics of the method | 5 |
| 11.1 Interlaboratory study..... | 5 |
| 11.2 Repeatability limit..... | 5 |
| 11.3 Reproducibility limit..... | 6 |
| 12 Test report | 6 |
| 13 Quality assurance | 7 |
| Annex A (normative) Culture media and reagents | 8 |
| Annex B (informative) Method validation studies and performance characteristics | 12 |
| Bibliography | 15 |

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the WTO principles in the Technical Barriers to Trade (TBT) see the following URL: www.iso.org/iso/foreword.html.

This document was prepared by the European Committee for Standardization (CEN) Technical Committee CEN/TC 275, *Food analysis — Horizontal methods*, in collaboration with ISO Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 9, *Microbiology* in accordance with the agreement on technical cooperation between ISO and CEN (Vienna Agreement).

This second edition cancels and replaces the first edition (ISO 21528-2:2004), which has been technically revised with the following main changes:

- the confirmation step has been changed by replacing glucose agar by glucose OF medium;
- precision data based on the results of an interlaboratory study using the method according to this revised edition has been included in an informative annex.

A list of all the parts in the ISO 21528 series can be found on the ISO website.

This corrected version of ISO 21528-2:2017 incorporates the following corrections:

- in [6.4](#), a water bath capable of being maintained between 44 °C to 47 °C has been added to the list of equipment;
- in [9.3.2](#), the temperature has been reduced from “47 °C to 50 °C” to “44 °C to 47 °C”.

Introduction

This document is intended to provide general guidance for the examination of products not dealt with by existing International Standards and to be taken into account by organizations preparing microbiological test methods for application to foods or animal feeding stuffs. Because of the large variety of products within this field of application, these guidelines may not be appropriate in every detail for certain products, and for some other products, it may be necessary to use different methods. Nevertheless, it is hoped that in all cases, every attempt will be made to apply the guidelines provided as far as possible and that deviations from them will only be made if absolutely necessary for technical reasons.

The main changes, listed in the foreword, introduced in this document compared to ISO 21528-2:2004, are considered as minor changes (see ISO 17468).

The harmonization of test methods cannot be immediate, and for certain groups of products, International Standards and/or national standards may already exist that do not comply with this horizontal method. It is hoped that when such standards are reviewed, they will be changed to comply with this document so that eventually the only remaining departures from this horizontal method will be those necessary for well-established technical reasons.

iTeh STANDARD PREVIEW (standards.iteh.ai)

[ISO 21528-2:2017](https://standards.iteh.ai/catalog/standards/sist/c8ce5547-a3ed-4128-a2e3-d9acca126a8e/iso-21528-2-2017)

<https://standards.iteh.ai/catalog/standards/sist/c8ce5547-a3ed-4128-a2e3-d9acca126a8e/iso-21528-2-2017>

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 21528-2:2017

<https://standards.iteh.ai/catalog/standards/sist/c8ce5547-a3ed-4128-a2e3-d9acca126a8e/iso-21528-2-2017>

Microbiology of the food chain — Horizontal method for the detection and enumeration of *Enterobacteriaceae* —

Part 2: Colony-count technique

WARNING — In order to safeguard the health of laboratory personnel, it is essential that tests for detecting *Enterobacteriaceae* are only undertaken in properly equipped laboratories, under the control of a skilled microbiologist, and that great care is taken in the disposal of all incubated materials. Persons using this document should be familiar with normal laboratory practice. This document does not purport to address all of the safety aspects, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices.

1 Scope

This document specifies a method for the enumeration of *Enterobacteriaceae*. It is applicable to

- products intended for human consumption and the feeding of animals, and
- environmental samples in the area of primary production, food production and food handling.

This technique is intended to be used when the number of colonies sought is expected to be more than 100 per millilitre or per gram of the test sample.

The most probable number (MPN) technique, as included in ISO 21528-1, is generally used when the number sought is expected to be below 100 per millilitre or per gram of test sample.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 6887 (all parts), *Microbiology of the food chain — Preparation of test samples, initial suspension and decimal dilutions for microbiological examination*

ISO 7218, *Microbiology of food and animal feeding stuffs — General requirements and guidance for microbiological examinations*

ISO 11133, *Microbiology of food, animal feed and water — Preparation, production, storage and performance testing of culture media*

ISO 18593, *Microbiology of food and animal feeding stuffs — Horizontal methods for sampling techniques from surfaces using contact plates and swabs*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <http://www.electropedia.org/>
- ISO Online browsing platform: available at <http://www.iso.org/obp>

3.1
Enterobacteriaceae
microorganism that forms characteristic colonies on violet red bile glucose agar and that ferment glucose and show a negative oxidase reaction when the tests are carried out in accordance with the methods specified in this document

3.2
count of *Enterobacteriaceae*
number of *Enterobacteriaceae* found per millilitre or per gram of the test sample or per unit area swabbed, when the test is carried out according to the method specified in this document

4 Principle

4.1 Preparation of initial suspension and decimal dilutions

An initial suspension and decimal dilutions are prepared from the test sample.

4.2 Isolation and selection for confirmation

Violet red bile glucose (VRBG) agar is inoculated with a specified quantity of the test sample if the product is liquid, or of the initial suspension in the case of other products. An overlay of the same medium is added. Other plates are prepared under the same conditions, using decimal dilutions of the test sample or of the initial suspension.

The dishes are incubated at 37 °C (or 30 °C) for 24 h.

NOTE The incubation temperature of 37 °C for enrichment and isolation/enumeration on the plating medium is generally used when the detection and enumeration of *Enterobacteriaceae* is for a hygiene indicator. Alternatively, a temperature of 30 °C can be chosen when the detection or enumeration of *Enterobacteriaceae* is conducted for technological purposes and includes psychrotrophic *Enterobacteriaceae*. In this document, 37 °C will be used throughout the text.

4.3 Confirmation

Colonies of presumptive *Enterobacteriaceae* are subcultured onto non-selective medium, and confirmed by means of tests for the fermentation of glucose and the presence of oxidase.

4.4 Calculation

The number of *Enterobacteriaceae* per millilitre or gram of the test sample is calculated from the number of confirmed typical colonies per dish.

5 Diluent, culture media and reagent

For current laboratory practice, see ISO 7218.

Composition of culture media and reagents and their preparation are described in [Annex A](#).

For performance testing of culture media, see ISO 11133 and [Annex A](#).

6 Equipment and consumables

Disposable equipment is an acceptable alternative to reusable glassware if it has suitable specifications. Usual microbiological laboratory equipment (see ISO 7218) and, in particular, the following.

6.1 Apparatus for dry sterilization (oven) or **wet sterilization** (autoclave), as specified in ISO 7218.

6.2 Incubator, capable of operating at $37\text{ °C} \pm 1\text{ °C}$ (or $30\text{ °C} \pm 1\text{ °C}$).

6.3 Drying cabinet (ventilated by convection) or **incubator**, capable of operating between 25 °C and 50 °C .

6.4 Water baths, one capable of being maintained between 47 °C to 50 °C and another capable of being maintained between 44 °C to 47 °C .

6.5 Test tubes or **flasks**, of appropriate capacity.

6.6 Petri dishes, made of glass or plastic, with a diameter of approximately 90 mm and (optional) large size (diameter approximately 140 mm).

6.7 Loops (of diameter approximately 3 mm) and **wires**, made of platinum/iridium or nickel/chromium, and/or **glass rods**, or equivalent sterile disposable loops or inoculating needles.

6.8 Graduated pipettes or **automatic pipettes**, of nominal capacities 10 ml, 1 ml and 0,1 ml.

6.9 pH-meter, accurate to within $\pm 0,1$ pH unit at 25 °C .

6.10 Homogenizer, as specified in ISO 7218.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

7 Sampling

Sampling is not part of the method specified in this document. See the specific International Standard dealing with the product concerned. If there is no specific International Standard dealing with the sampling of the product concerned, it is recommended that the parties concerned come to an agreement on this subject.

Recommended sampling techniques are given in:

- ISO/TS 17728 for food and animal feed;
- ISO 13307 for primary production stage;
- ISO 17604 for carcasses;
- ISO 18593 for environmental samples.

It is important that the laboratory receives a sample that is representative and the sample should not have been damaged or changed during transport or storage.

8 Preparation of test sample

Prepare the test sample in accordance with the specific International Standard appropriate to the product concerned. If there is no specific International Standard available, it is recommended that the parties concerned come to an agreement on this subject.

9 Procedure

9.1 General

See ISO 7218.

9.2 Test portion, initial suspension and dilutions

See ISO 6887 (all parts).

Prepare a single decimal dilution series from the test sample if the product is liquid, or from the initial suspension in the case of other products.

9.3 Inoculation and incubation

9.3.1 Take one sterile Petri dish (6.6). Using a sterile pipette (6.8), transfer to the dish 1 ml of the test sample if the product is liquid, or 1 ml of the initial suspension in case of other products.

Repeat the procedure described with the further dilutions, if necessary, using a fresh sterile pipette for each dilution.

If only the initial suspension is used, then inoculate two plates of this dilution (ISO 7218).

9.3.2 Add into each Petri dish approximately 15 ml of the violet red bile glucose (VRBG) agar (A.2) which has been prepared then cooled to 44 °C to 47 °C in the water bath (6.4). The time elapsing between the inoculation of the Petri dishes and the moment when the medium is poured into the dishes shall not exceed 15 min.

Carefully mix the inoculum with the medium by horizontal movements and allow the medium to solidify, with the Petri dishes standing on a cool surface.

9.3.3 After complete solidification of the mixture, add a covering layer of approximately 5 ml to 10 ml of the violet red bile glucose (VRBG) agar (A.2), prepared then cooled as described in 9.3.2, to prevent spreading growth and to achieve semi-anaerobic conditions. Allow to solidify as described above.

9.3.4 Invert the prepared dishes and incubate them at 37 °C (6.2) for 24 h ± 2 h.

9.4 Counting and selection of colonies for confirmation

Characteristic colonies are pink to red or purple (with or without precipitation haloes).

Select the dishes (see 9.3.4) containing less than 150 characteristic colonies; count these colonies. Then choose at random five such colonies from each dish for subculturing (see 9.5) for the biochemical confirmation tests (see 9.6). If there are less than five colonies on the plate, take all presumptive colonies present.

Spreading colonies shall be considered as single colonies. If less than one-quarter of the dish is overgrown by spreading, count the colonies on the unaffected part of the dish and calculate by extrapolation the theoretical number of colonies for the entire plate. If more than one-quarter is overgrown by spreading colonies, discard the count.

Certain *Enterobacteriaceae* may cause decolouration of their colonies or of the medium. Therefore, when no characteristic colonies are present, choose five whitish colonies for confirmation.

9.5 Subculturing selected colonies

Streak the selected colonies (9.4) onto the surface of pre-dried non-selective agar medium (A.3), in a manner which will allow well-isolated colonies to develop.

Incubate these plates at 37 °C for 24 h ± 2 h.

Select a well-isolated colony from each of the incubated plates for the biochemical confirmation tests (see 9.6).

9.6 Biochemical confirmation tests

9.6.1 Oxidase reaction

Using a platinum/iridium loop, wire or glass rod (6.7), take a portion of each well-isolated colony (see 9.5) and streak onto a filter paper moistened with the oxidase reagent (A.5) or onto a commercially available disc or stick. A nickel/chromium loop or wire shall not be used.

Consider the test to be negative when the colour of the filter paper does not turn dark blue purple within 10 s.

Consult the manufacturer's instructions for ready-to-use discs or sticks.

9.6.2 Fermentation test

Using a wire (6.7), stab the same colonies selected in 9.5 that gave a negative oxidase test into tubes containing Glucose OF medium (A.4). Overlay the surface of the medium with minimal 1 cm of sterile mineral oil (A.6).

Incubate these tubes at 37 °C for 24 h ± 2 h.

If a yellow colour develops throughout the content of the tube, regard the reaction as being positive.

9.6.3 Interpretation of biochemical tests

Colonies that are oxidase-negative and glucose-positive are confirmed as *Enterobacteriaceae*.

10 Expression of results

See ISO 7218. Calculate and report the results as the number of *Enterobacteriaceae* in cfu per gram, millilitre, per square centimetre or per sampling device.

11 Performance characteristics of the method

11.1 Interlaboratory study

Results of the interlaboratory study to determine the precision of the method are summarized in Annex B. Repeatability and reproducibility limits were determined using five types of food contaminated at various levels. The values derived from the interlaboratory study may not be applicable to concentration ranges and food types other than those given in Annex B.

NOTE In this document, the word "type" is combined with "food" to improve the readability of this document. However, the word "food" is interchangeable with "feed" and the other areas of the food chain as mentioned in the scope of this document.

11.2 Repeatability limit

The absolute difference between two independent single (\log_{10} -transformed) test results (number of cfu per gram or per millilitre) or the ratio of the higher to the lower of the two test results on the normal scale, obtained using the same method on identical test material in the same laboratory by the same operator using the same apparatus within the shortest feasible time interval, will, in not more than 5 % of cases, exceed the repeatability limit r .

As a general indication of repeatability limit (r), the following values (derived from the mean of the variance estimates for all levels per matrix tested in the interlaboratory study (see Annex B) may be used when testing samples:

$r = 0,37$ (expressed as a difference between \log_{10} -transformed test results); or