

---

---

**Microbiology of the food chain —  
Horizontal method for the  
enumeration of beta-glucuronidase-  
positive *Escherichia coli* —**

Part 1:

**Colony-count technique at 44 °C using  
membranes and 5-bromo-4-chloro-3-  
indolyl beta-D-glucuronide**

ISO 16649-1:2018  
https://standards.iteh.ai/catalog/standards/sist/7254767c-bacc-4197-8b00-751ce39e2dce/iso-16649-1-2018

*Microbiologie de la chaîne alimentaire — Méthode horizontale pour  
le dénombrement des *Escherichia coli* bêta-glucuronidase positive —  
Partie 1: Technique de comptage des colonies à 44 °C au moyen de  
membranes et de 5-bromo-4-chloro-3-indolyl bêta-D glucuronide*



**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

ISO 16649-1:2018

<https://standards.iteh.ai/catalog/standards/sist/7254767c-0ace-4f97-8b09-751ce39e2dee/iso-16649-1-2018>



**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2018

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
CP 401 • Ch. de Blandonnet 8  
CH-1214 Vernier, Geneva  
Phone: +41 22 749 01 11  
Fax: +41 22 749 09 47  
Email: [copyright@iso.org](mailto:copyright@iso.org)  
Website: [www.iso.org](http://www.iso.org)

Published in Switzerland

# Contents

	Page
Foreword.....	iv
Introduction.....	v
<b>1 Scope.....</b>	<b>1</b>
<b>2 Normative references.....</b>	<b>1</b>
<b>3 Terms and definitions.....</b>	<b>1</b>
<b>4 Principle.....</b>	<b>2</b>
4.1 Test portion, initial suspension, dilutions and resuscitation step.....	2
4.2 Culture on selective medium.....	2
4.3 Calculation.....	2
<b>5 Culture media and reagents.....</b>	<b>2</b>
<b>6 Equipment and consumables.....</b>	<b>2</b>
<b>7 Sampling.....</b>	<b>3</b>
<b>8 Preparation of test sample.....</b>	<b>3</b>
<b>9 Procedure.....</b>	<b>3</b>
9.1 General.....	3
9.2 Test portion, initial suspension and dilutions.....	4
9.3 Resuscitation.....	4
9.4 Transfer to selective medium and incubation.....	4
9.5 Counting the colony-forming units (cfu).....	4
9.6 Calculation.....	4
<b>10 Expression of results.....</b>	<b>4</b>
<b>11 Performance characteristics of the method.....</b>	<b>5</b>
<b>12 Test report.....</b>	<b>5</b>
<b>13 Quality assurance.....</b>	<b>5</b>
<b>Annex A (normative) Composition and preparation of culture media and reagents.....</b>	<b>6</b>
<b>Bibliography.....</b>	<b>10</b>

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see [www.iso.org/directives](http://www.iso.org/directives)).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see [www.iso.org/patents](http://www.iso.org/patents)).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see the following URL: [www.iso.org/iso/foreword.html](http://www.iso.org/iso/foreword.html). (standards.itech.ai)

This document was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 9, *Microbiology*.

<https://standards.itech.ai/catalog/standards/sist/7254767c-0ace-4f97-8b09-175f39e2d431/iso-16649-1:2018>

This second edition cancels and replaces the first edition (ISO 16649-1:2001), which has been technically revised with the following main changes:

- samples from the environment and the primary production stage have been added to the Scope;
- the minimum length of incubation (20 h) for tryptone-bile X-glucuronide agar (TBX) during culture on selective medium has been adopted;
- performance testing for the quality assurance of the culture media and the membrane for transfer has been added;
- to improve safety for the user, the solvent dimethyl sulphoxide (DMSO) is no longer recommended to dissolve the chromogenic substrate (BCIG);
- the composition of the minerals-modified glutamate agar (MMGA) has been corrected (aspartic acid 0,024 g and arginine 0,02 g) to the values in the original formulation<sup>[12]</sup>.

A list of all parts in the ISO 16649 series can be found on the ISO website.

## Introduction

Because of the large variety of food and feed products, this horizontal method may not be appropriate in every detail for certain products. In this case, different methods which are specific to these products may be used if absolutely necessary for justified technical reasons. Nevertheless, every attempt should be made to apply this horizontal method as far as possible.

The main changes, listed in the Foreword, introduced in this document compared to ISO 16649-1:2001 are considered as minor (see ISO 17468)<sup>[5]</sup>.

When this document is next reviewed, account will be taken of all information then available regarding the extent to which this horizontal method has been followed and the reasons for deviations from this method in the case of particular products.

The harmonization of test methods cannot be immediate and, for certain groups of products, International Standards and/or national standards may already exist that do not comply with this horizontal method. It is hoped that when such standards are reviewed they will be changed to comply with this document so that eventually the only remaining departures from this horizontal method will be those necessary for well-established technical reasons.

There are three horizontal methods (ISO 16649-1, ISO 16649-2 and ISO 16649-3) for the enumeration of  $\beta$ -glucuronidase-positive *Escherichia coli*<sup>[3][4]</sup>.

The user may choose either ISO 16649-1, ISO 16649-2 or ISO 16649-3. All parts are for general application. However, ISO 16649-1 or ISO 16649-3, which both include a resuscitation step, should be used in preference for foodstuffs likely to contain sub-lethally injured cells as a result of properties associated with the food or processing conditions.

[ISO 16649-1:2018](https://standards.iteh.ai/catalog/standards/sist/7254767c-0ace-4f97-8b09-751ce39e2dee/iso-16649-1-2018)

<https://standards.iteh.ai/catalog/standards/sist/7254767c-0ace-4f97-8b09-751ce39e2dee/iso-16649-1-2018>

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

ISO 16649-1:2018

<https://standards.iteh.ai/catalog/standards/sist/7254767c-0ace-4f97-8b09-751ce39e2dee/iso-16649-1-2018>

# Microbiology of the food chain — Horizontal method for the enumeration of beta-glucuronidase-positive *Escherichia coli* —

Part 1:

## Colony-count technique at 44 °C using membranes and 5-bromo-4-chloro-3-indolyl beta-D-glucuronide

### 1 Scope

This document specifies a horizontal method for the enumeration of  $\beta$ -glucuronidase-positive *Escherichia coli* by colony-count technique after resuscitation using membranes and incubation at 44 °C on a solid medium containing a chromogenic ingredient for detection of the enzyme  $\beta$ -glucuronidase<sup>[9]</sup><sup>[10]</sup><sup>[13]</sup><sup>[14]</sup><sup>[17]</sup><sup>[18]</sup><sup>[19]</sup><sup>[20]</sup>. It is applicable to

- products intended for human consumption,
- products intended for feeding animals,
- environmental samples in the area of food production and food handling, and
- samples from the primary production stage such as animal faeces, dust, and swabs.

**WARNING** — Some strains of *Escherichia coli* may grow poorly or not at all in media incubated at 44 °C. This includes strains of *E. coli* O157:H7 and O157:H-. Additionally, some strains of *Escherichia coli*, notably those belonging to serotype O157:H7, are mostly  $\beta$ -glucuronidase negative<sup>[11]</sup>. Consequently, some strains of *E. coli*, including pathogenic ones, will not be detected by this method.  $\beta$ -glucuronidase activity may also be exhibited at 44 °C by certain other members of the *Enterobacteriaceae*, notably *Shigella*<sup>[15]</sup> and *Salmonella*<sup>[16]</sup>.

### 2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 6887 (all parts), *Microbiology of the food chain — Preparation of test samples, initial suspension and decimal dilutions for microbiological examination*

ISO 7218, *Microbiology of food and animal feeding stuffs — General requirements and guidance for microbiological examinations*

ISO 11133, *Microbiology of food, animal feed and water — Preparation, production, storage and performance testing of culture media*

### 3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <http://www.electropedia.org/>

— ISO Online browsing platform: available at <https://www.iso.org/obp>

**3.1**  
 **$\beta$ -glucuronidase-positive *Escherichia coli***  
bacteria which at 44 °C forms typical, blue or blue-green colonies on tryptone-bile X-glucuronide agar (TBX) under the conditions specified in this document

**3.2**  
**enumeration of  $\beta$ -glucuronidase-positive *Escherichia coli***  
determination of the number of colony-forming units (cfu) of  $\beta$ -glucuronidase-positive *Escherichia coli* (3.1), per gram of sample, per millilitre, per square centimetre or per sampling device when the analysis is carried out in accordance with this document

## 4 Principle

### 4.1 Test portion, initial suspension, dilutions and resuscitation step

A specified quantity of the test sample, initial suspension or decimal dilutions, is inoculated onto membranes overlaid on minerals-modified glutamate agar (MMGA), then incubated at 37 °C for 4 h<sup>[13]</sup><sup>[14]</sup>.

### 4.2 Culture on selective medium

For isolation, the membranes from the resuscitation stage on the MMGA are transferred to tryptone-bile X-glucuronide agar (TBX), then incubated at 44 °C for 20 h to 24 h.

### 4.3 Calculation

The number of colony-forming units (cfu) of  $\beta$ -glucuronidase-positive *Escherichia coli* per gram or per millilitre of sample is calculated from the number of typical blue or blue-green colonies per plate.

## 5 Culture media and reagents

For current laboratory practice, see ISO 7218 and ISO 11133.

Composition of culture media and reagents and their preparation are described in [Annex A](#).

For performance testing of culture media and membrane for transfer, see ISO 11133 and [Annex A](#).

## 6 Equipment and consumables

Disposable equipment is an acceptable alternative to reusable glassware if it has suitable specifications.

Usual microbiological laboratory equipment (see ISO 7218) and, in particular, the following.

**6.1 Apparatus for dry sterilization (oven) or wet sterilization (autoclave).** As specified in ISO 7218.

**6.2 Drying cabinet or ventilated oven,** capable of being maintained between 25 °C and 50 °C, or a laminar airflow cabinet.

**6.3 Incubator,** capable of operating at 37 °C  $\pm$  1 °C.

**6.4 Incubator,** capable of operating at 44 °C  $\pm$  1 °C.

**6.5 Water bath,** capable of operating at 47 °C to 50 °C.



- 6.6 Refrigerator** (for storage of prepared media), capable of operating at  $5\text{ °C} \pm 3\text{ °C}$ .
- 6.7 Blunt-ended forceps**, sterile, of approximately 12 cm length.
- 6.8 Sterile and non-inhibitory membranes**, made of cellulose acetate or mixed esters of cellulose, with  $0,45\text{ }\mu\text{m}$  to  $1,2\text{ }\mu\text{m}$  pore size and 85 mm diameter.
- 6.9 pH-meter**, capable of being read to the nearest 0,01 pH unit at  $25\text{ °C}$ , enabling measurements to be made which are accurate to  $\pm 0,1$  pH unit.
- 6.10 Sterile graduated pipettes or automatic pipettes**, of nominal capacity 1 ml.
- 6.11 Sterile petri dishes** with a diameter of approximately 90 mm diameter.
- 6.12 Sterile spreaders**, made of glass or plastic, for example, hockey sticks made from a rod of approximately  $3,5\text{ mm}$  diameter and 20 cm length, bent at right angles about 3 cm from one end and with the cut ends made smooth by heating.

## 7 Sampling

Sampling is not part of the method specified in this document. See the specific International Standard dealing with the product concerned. If there is no specific International Standard dealing with the sampling of the product concerned, it is recommended that the parties concerned come to an agreement on this subject.

Recommended sampling methods are given in the following documents:

- ISO/TS 17728<sup>[7]</sup> for food and animal feed;  
<https://standards.iteh.ai/catalog/standards/sist/7254767c-0ace-4f97-8b09-751ce39e2dee/iso-16649-1-2018>
- ISO 707<sup>[1]</sup> for milk and milk products;
- ISO 13307<sup>[2]</sup> for sampling at primary production stage;
- ISO 17604<sup>[6]</sup> for sampling of carcasses;
- ISO 18593<sup>[8]</sup> for sampling of surfaces.

It is important that the laboratory receives a sample that is representative and has not been damaged or changed during transport or storage.

## 8 Preparation of test sample

Prepare the test sample from the laboratory sample in accordance with the specific International Standard dealing with the product concerned: see ISO 6887 (all parts). If there is no specific International Standard, it is recommended that the parties concerned come to an agreement on this subject.

## 9 Procedure

### 9.1 General

For general aspects refer to ISO 7218.