
INTERNATIONAL STANDARD



1897 / VII

INTERNATIONAL ORGANIZATION FOR STANDARDIZATION • МЕЖДУНАРОДНАЯ ОРГАНИЗАЦИЯ ПО СТАНДАРТИЗАЦИИ • ORGANISATION INTERNATIONALE DE NORMALISATION

**Phenol, *o*-cresol, *m*-cresol, *p*-cresol, cresylic acid and xylenols
for industrial use — Methods of test —
Part VII : Measurement of colour (Cresylic acid and xylenols
only)**

iTeh STANDARD PREVIEW

*Phénol, *o*-crésol, *m*-crésol, *p*-crésol, acide crésylique et xylénols à usage industriel — Méthodes d'essai —
Partie VII : Mesurage de la coloration (Acide crésylique et xylénols uniquement)*

<https://standards.iteh.ai/catalog/standards/sist/ac9e4b2a-0fb7-4100-a192-c16011ddaca6/iso-1897-7-1977>

First edition — 1977-04-15

<https://standards.iteh.ai/catalog/standards/sist/ac9e4b2a-0fb7-4100-a192-c16011ddaca6/iso-1897-7-1977>

UDC 661 : 547.562/563 : 535.661.4 : 620.1

Ref. No. ISO 1897/VII-1977 (E)

Descriptors : phenols, phenol, cresols, xyleneol, chemical analysis, determination of content, water, volumetric analysis, Karl Fischer reagent.

Price based on 2 pages

FOREWORD

ISO (the International Organization for Standardization) is a worldwide federation of national standards institutes (ISO member bodies). The work of developing International Standards is carried out through ISO technical committees. Every member body interested in a subject for which a technical committee has been set up has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work.

Draft International Standards adopted by the technical committees are circulated to the member bodies for approval before their acceptance as International Standards by the ISO Council.

Prior to 1972, the results of the work of the technical committees were published as ISO Recommendations; these documents are in the process of being transformed into International Standards. As part of this process, Technical Committee ISO/TC 47, *Chemistry*, has reviewed ISO Recommendation R 1909-1971 and found it technically suitable for transformation. Number 1909 however, has been changed to 1897/VII. International Standard ISO 1897/VII therefore replaces ISO Recommendation R 1909-1971, to which it is technically identical.

ISO Recommendation R 1909 had been approved by the member bodies of the following countries :

Australia	India	South Africa, Rep. of
Belgium	Israel	Spain
Chile	Italy	Switzerland
Czechoslovakia	Japan	Thailand
Egypt, Arab Rep. of	Netherlands	Turkey
France	New Zealand	United Kingdom
Germany	Poland	U.S.S.R.
Greece	Portugal	
Hungary	Romania	

No member body had expressed disapproval of the Recommendation.

The member body of the following country disapproved the transformation of the Recommendation into an International Standard :

Netherlands

Phenol, *o*-cresol, *m*-cresol, *p*-cresol, cresylic acid and xylenols for industrial use – Methods of test – Part VII : Measurement of colour (Cresylic acid and xylenols only)

1 SCOPE AND FIELD OF APPLICATION

This part of ISO 1897 specifies a method for the measurement of colour of cresylic acid of high *o*-cresol content, cresylic acid of high *m*-cresol content and xylenols, for industrial use.

This document should be read in conjunction with part I (see the annex).

2 PRINCIPLE

Comparison of the colour of a test portion against that of standard colour matching solutions.

NOTE – The colour of cresols present in the test portion is liable to darken on keeping and on exposure to light.

3 REAGENTS

During the test, use only reagents of recognized analytical grade, and only distilled water or water of equivalent purity.

3.1 Cobalt sulphate heptahydrate ($\text{CoSO}_4 \cdot 7\text{H}_2\text{O}$).

3.2 Copper(II) sulphate pentahydrate ($\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$).

3.3 Potassium dichromate ($\text{K}_2\text{Cr}_2\text{O}_7$).

3.4 Potassium hexacyanoferrate(III) [$\text{K}_3\text{Fe}(\text{CN})_6$].

4 APPARATUS

Ordinary laboratory apparatus, and

4.1 Two matched Nessler cylinders, capacity 50 ml.

4.2 Opaque opal glass sheet.

5 PREPARATION OF STANDARD COLORIMETRIC SOLUTIONS

Prepare the following standard colorimetric solutions, dissolving in water the indicated quantities of reagents and diluting to 1 000 ml in one-mark volumetric flasks.

The tint of the even-numbered colours is red and that of the odd-numbered colours is yellow.

The standard colorimetric solution No.7 should be freshly prepared on the day of the test. The other solutions

keep well and may be used up to 1 month from the date of preparation.

Colour No.	Mass of reagent g	Reagent
1	0,90 0,015	Cobalt sulphate (3.1) Potassium dichromate (3.3)
2	6,0 0,015	Cobalt sulphate (3.1) Potassium dichromate (3.3)
3	4,0 0,075 1,0	Cobalt sulphate (3.1) Potassium dichromate (3.3) Copper(II) sulphate (3.2)
4	22,5 0,06	Cobalt sulphate (3.1) Potassium dichromate (3.3)
5	10,0 0,18 2,5	Cobalt sulphate (3.1) Potassium dichromate (3.3) Copper(II) sulphate (3.2)
6	70,0 0,5	Cobalt sulphate (3.1) Potassium dichromate (3.3)
7	320,0	Potassium hexacyanoferrate(III) (3.4)
8		Any cresylic acid darker than colour No. 7

6 PROCEDURE

6.1 Test portion

Pour 50 ml of the test sample into one of the Nessler cylinders (4.1).

6.2 Comparison

Pour 50 ml of the standard colorimetric solution (see clause 5) agreed between the parties, into the second Nessler cylinder.

Compare the colour of the two liquids in the Nessler cylinders, held vertically 75 mm above the surface of the opaque opal glass sheet (4.2) reflecting diffused daylight.

NOTE – It is usual to specify two standard colorimetric solutions, one from the even-numbered series and one from the odd-numbered series.

7 EXPRESSION OF RESULTS

Report the colour of the test portion as being not darker than, equal to, or darker than those of the standard colorimetric solutions agreed between the parties.

ANNEX

ISO PUBLICATIONS RELATING TO (A) PHENOL, (B) *o*-CRESOL, (C) *m*-CRESOL, (D) *p*-CRESOL, (E) CRESYLIC ACID, AND (F) XYLENOLS, FOR INDUSTRIAL USE

Applicability						
A ¹⁾	B ²⁾	C	D ²⁾	E	F	ISO 1897/I – General.
A	B	C	D	E	F	ISO 1897/II – Determination of water – Dean and Stark method.
A	B	C	D	E	F	ISO 1897/III – Determination of neutral oils and pyridine bases.
A	B	C	D			ISO 1897/IV – Visual test for impurities insoluble in sodium hydroxide solution.
A						ISO 1897/V – Visual test for impurities insoluble in water.
				E	F	ISO 1897/VI – Test for absence of hydrogen sulphide.
				E	F	ISO 1897/VII – Measurement of colour.
				E	F	ISO 1897/VIII – Determination of <i>o</i> -cresol content.
				E		ISO 1897/IX – Determination of <i>m</i> -cresol content.
A	B	C	D			ISO/R 1900 – Determination of residue on evaporation.
A	B	C	D			ISO/R 1901 – Determination of crystallizing point.
A ³⁾						ISO 1904 – Determination of phenols content – Bromination method.
				E	F	ISO/R 1906 – Determination of distillation range.
				E	F	ISO/R 1907 – Determination of residue on distillation.
A	B	C	D			ISO 2208 – Determination of crystallizing point after drying with a molecular sieve.

1) In the case of phenol, the determination of density at 20 °C specified in ISO 1897/I is applicable only to liquefied phenol.

2) The determination of density at 20 °C specified in ISO 1897/I is not applicable to these products.

3) Applicable only to liquefied phenol.