

SLOVENSKI STANDARD SIST-TS CEN/TS 17303:2019

01-maj-2019

Živila - Črtno kodiranje DNK rib in ribjih proizvodov z uporabo segmentov genov, ki nosijo zapis za mitohondrijski citokrom b in citokrom c oksidaze l

Foodstuffs - DNA barcoding of fish and fish products using defined mitochondrial cytochrome b and cytochrome c oxidase I gene segments

Lebensmittel - DNA-Barcoding von Fisch und Fischprodukten anhand definierter mitochondrialer Cytochrom b- und Cytochrom c-Oxidase I-Genabschnitte

Produits alimentaires - Codes-barres d'ADN de poissons et de produits à base de poissons à l'aide de segments de gènes mitochondriaux du cytochrome b et cyctochrome c oxydase/lstandards.iteh.ai/catalog/standards/sist/ad380310-ca08-48cf-a25e-

c95e681d2583/sist-ts-cen-ts-17303-2019

Ta slovenski standard je istoveten z: CEN/TS 17303:2019

ICS:

67.120.30 Ribe in ribji proizvodi Fish and fishery products

SIST-TS CEN/TS 17303:2019

en,fr,de

iTeh STANDARD PREVIEW (standards.iteh.ai)

SIST-TS CEN/TS 17303:2019 https://standards.iteh.ai/catalog/standards/sist/ad380310-ca08-48cf-a25ec95e681d2583/sist-ts-cen-ts-17303-2019

SIST-TS CEN/TS 17303:2019

TECHNICAL SPECIFICATION SPÉCIFICATION TECHNIQUE TECHNISCHE SPEZIFIKATION

CEN/TS 17303

March 2019

ICS 67.120.30

English Version

Foodstuffs - DNA barcoding of fish and fish products using defined mitochondrial cytochrome b and cytochrome c oxidase I gene segments

Produits alimentaires - Codes-barres d'ADN de poissons et de produits à base de poissons à l'aide de segments de gènes mitochondriaux du cytochrome b et cyctochrome c oxydase I Lebensmittel - DNA-Barcoding von Fisch und Fischprodukten anhand definierter mitochondrialer Cytochrom-b- und Cytochrom-c-Oxidase-I-Genabschnitte

This Technical Specification (CEN/TS) was approved by CEN on 14 January 2019 for provisional application.

The period of validity of this CEN/TS is limited initially to three years. After two years the members of CEN will be requested to submit their comments, particularly on the question whether the CEN/TS can be converted into a European Standard.

CEN members are required to announce the existence of this CEN/TS in the same way as for an EN and to make the CEN/TS available promptly at national level in an appropriate form. It is permissible to keep conflicting national standards in force (in parallel to the CEN/TS) until the final decision about the possible conversion of the CEN/TS into an EN is reached.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom. c95e681d2583/sist-ts-cen-ts-17303-2019



EUROPEAN COMMITTEE FOR STANDARDIZATION COMITÉ EUROPÉEN DE NORMALISATION EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Rue de la Science 23, B-1040 Brussels

Ref. No. CEN/TS 17303:2019 E

SIST-TS CEN/TS 17303:2019

CEN/TS 17303:2019 (E)

Contents

| Europ | opean foreword | | |
|--------|--|----|--|
| Introd | uction | 4 | |
| 1 | Scope | 5 | |
| 2 | Normative references | 5 | |
| 3 | Terms and definitions | 5 | |
| 4 | Principle | 7 | |
| 5 | Reagents and materials | | |
| 5.1 | General | | |
| 5.2 | PCR reagents | 8 | |
| 6 | Apparatus | 9 | |
| 7 | Procedure | 9 | |
| 7.1 | Sample preparation | | |
| 7.2 | DNA extraction | | |
| 7.3 | PCR | 9 | |
| 7.3.1 | | | |
| 7.3.2 | PCR setup | 10 | |
| 7.3.3 | | | |
| 7.3.4 | PCR controls | 11 | |
| 8 | Evaluation | 12 | |
| 8.1 | Evaluation of PCR products c95e681d2583/sist-ts-cen-ts-17303-2019 | | |
| 8.2 | Evaluation of the PCR results | | |
| 8.3 | Sequencing of PCR products | | |
| 8.4 | Evaluation of sequence data | | |
| 8.5 | Comparison of the sequence with public databases | | |
| 8.5.1 | General | | |
| 8.5.2 | Sequence comparison of <i>cytb</i> and/or <i>cox1</i> DNA sequences with GenBank | | |
| 8.5.3 | Sequence comparison of <i>cox1</i> DNA sequences with BOLD | | |
| 9 | Interpretation of database query results | 15 | |
| 10 | Validation status and performance criteria | 15 | |
| | Collaborative study for the identification of fish species based on <i>cytb</i> sequence | | |
| 10.1 | analysis | | |
| 10.2 | Collaborative study for the identification of fish species based on <i>cox1</i> sequence | 15 | |
| 10.2 | analysis | 16 | |
| | | | |
| 11 | Test report | 18 | |
| Annex | A (informative) Practical laboratory experiences with the amplificability of cytb or | 10 | |
| | cox1 segments from tested fish species | 19 | |
| Biblio | graphy | 24 | |

European foreword

This document (CEN/TS 17303:2019) has been prepared by Technical Committee CEN/TC 275 "Food analysis - Horizontal methods", the secretariat of which is held by DIN.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN shall not be held responsible for identifying any or all such patent rights.

According to the CEN/CENELEC Internal Regulations, the national standards organisations of the following countries are bound to announce this Technical Specification: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

iTeh STANDARD PREVIEW (standards.iteh.ai)

SIST-TS CEN/TS 17303:2019 https://standards.iteh.ai/catalog/standards/sist/ad380310-ca08-48cf-a25ec95e681d2583/sist-ts-cen-ts-17303-2019

CEN/TS 17303:2019 (E)

Introduction

Food safety is a key aspect in terms of consumer protection. In the last three decades, globalization has taken place in the trade of food. Fish trade channels are becoming steadily longer and more complicated so that sophisticated traceability tools are needed to ensure food safety. Correct food labelling is a prerequisite to ensure safe fish products and fair trade as well as to minimize illegal, unreported and unregulated (IUU) fishing. In particular, the fact that fish is increasingly being processed in export countries makes the identification of species by morphological characteristics impossible.

The development of harmonized and standardized protocols for the authentication of fish products is necessary to establish reliable methods for the detection of potential food fraud.

iTeh STANDARD PREVIEW (standards.iteh.ai)

<u>SIST-TS CEN/TS 17303:2019</u> https://standards.iteh.ai/catalog/standards/sist/ad380310-ca08-48cf-a25ec95e681d2583/sist-ts-cen-ts-17303-2019

1 Scope

This document describes a procedure for the identification of single fish and fish fillets to the level of genus or species.

The identification of fish species is carried out by PCR amplification of either a segment of the mitochondrial cytochrome b gene (*cytb*) [1] or the cytochrome c oxidase I gene (*cox1, syn COI*) [2], [3] or both, followed by sequencing of the PCR products and subsequent sequence comparison with entries in databases [4], [5]. The methodology allows the identification of a large number of commercially important fish species.

The decision whether the *cytb* or *cox1* gene segment or both are used for fish identification depends on the declared fish species, the applicability of the PCR method for the fish species and the availability of comparative sequences in the public databases.

This method has been successfully validated on raw fish fillets, however, laboratory experience is available that it can also be applied to processed, e.g. cold smoked, hot smoked, salted, frozen, cooked, fried, deep-fried samples.

This document is usually unsuitable for the analysis of highly processed foods, e.g. tins of fish, with highly degraded DNA where the fragment lengths are not sufficient for amplification of the targets. Furthermore, it is not applicable for complex fish products containing mixtures of two or more fish species.

2 Normative references TANDARD PREVIEW

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN ISO 24276, Foodstuffs — Methods of analysis for the detection of genetically modified organisms and derived products — General requirements and definitions (ISO 24276)

ISO 16577, Molecular biomarker analysis — Terms and definitions

3 Terms and definitions

For the purposes of this document, the terms and definitions given in ISO 16577 and the following apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at http://www.electropedia.org/
- ISO Online browsing platform: available at http://www.iso.org/obp

3.1

alignment

process or result of matching up the nucleotide residues of two or more biological sequences to achieve maximal levels of identity

[SOURCE: BLAST Glossary]

3.2 BLAST (The Basic Local Alignment Search Tool) [4]

sequence comparison algorithm optimized for speed used to search sequence databases for optimal local alignments to a query

Note 1 to entry: It directly approximates alignments that optimize a measure of local similarity, the maximum signal pair (MSP) score or high scoring signal pair (HSP) score.

3.3 BOLD

(Barcode of Life Data Systems) [5]

informatics workbench aiding the acquisition, storage, analysis, and publication of DNA barcode records

Note 1 to entry: By assembling molecular, morphological, and distributional data, it bridges a traditional bioinformatics chasm. BOLD is freely available to any researcher with interests in DNA barcoding. By providing specialized services, it aids the assembly of records that meet the standards needed to gain BARCODE designation in the global sequence databases. Because of its web-based delivery and flexible data security model, it is also well positioned to support projects that involve broad research alliances.

[SOURCE: BOLDSYSTEMS About Us]

3.4

FASTA format

iTeh STANDARD PREVIEW

text-based format for representing either nucleotide sequences or amino acid sequences, which begins with a single-line description, followed by lines of sequence data

Note 1 to entry: The description line (defline) is distinguished from the sequence data by a greater-than (">") symbol at the beginning. https://standards.iteh.ai/catalog/standards/sist/ad380310-ca08-48cf-a25e-c95e681d2583/sist-ts-cen-ts-17303-2019

[SOURCE: BLAST topics, modified]

EXAMPLE An example sequence in FASTA format is shown below:

> Sample_04_cytb

3.5

FishBase

global biodiversity online platform on finfishes providing a wide range of information on all species currently known in the world

3.6

GenBank

comprehensive public database of e.g. genetic sequences [6]

Note 1 to entry: GenBank is part of the <u>International Nucleotide Sequence Database Collaboration</u>, which comprises the DNA DataBank of Japan (DDBJ), the European Nucleotide Archive (ENA), and GenBank at National Center for Biotechnology Information (NCBI). These three organizations exchange data on a daily basis.

CEN/TS 17303:2019 (E)

3.7

identity

extent to which two (nucleotide or amino acid) sequences have the same residues at the same positions in an alignment, often expressed as a percentage

[Source: BLAST Glossary]

Note 1 to entry: In the database BOLD, the term similarity is used instead of identity.

3.8

introgressed DNA

DNA sequence (allele) from one taxonomic entity (species) incorporated in the gene pool of another, divergent entity (species)

Note 1 to entry: Introgression has usually happened via hybridization and backcrossing of individuals belonging to different species.

3.9

NCBI (National Center for Biotechnology Information)

institution which houses molecular biology databases (e.g. GenBank) and provides the BLAST suite

3.10

nucleotide collection (nr/nt)

non-redundant database consisting of GenBank sequences, in which identical sequences have been merged into one entry

3.11

(standards.iteh.ai)

query <u>SIST-TS CEN/TS 17303.2019</u> sequence (or other type of search term) to which all of the entries in a data base are to be compared

[SOURCE: BLAST Glossary].

c95e681d2583/sist-ts-cen-ts-17303-2019

3.12

query coverage

percentage of query covered by alignment to the data base sequence

[SOURCE: BLAST help]

4 Principle

DNA is extracted from fish and fish products applying a suitable method. Segments of approximately 460 base pairs of the *cytb* gene and/or approximately 650 base pairs of the *cox1* gene are amplified by PCR. In the further course, the nucleotide sequence of the PCR product is determined by a suitable DNA sequencing method (e.g. Sanger sequencing). The sequence is evaluated by comparison to sequence entries in databases, thus allowing the assignment to a fish species or genus according to the degree of identity with stored sequences.

5 Reagents and materials

5.1 General

During the analysis, unless otherwise stated, use only reagents of recognized molecular biology grade and distilled or demineralized water or water of equivalent purity, according to EN ISO 24276. Regarding laboratory organization, see EN ISO 24276.

5.2 PCR reagents

- 5.2.1 Thermostable DNA polymerase (for hot start PCR)¹
- 5.2.2 PCR reaction buffer (including MgCl₂ or with separate MgCl₂ solution)¹
- 5.2.3 Deoxynucleoside triphosphate mix (dATP, dCTP, dGTP and dTTP)²
- 5.2.4 Oligonucleotides (see Tables 1 and 2)

Table 1 — Oligonucleotides for amplification of the *cytb* gene region [1]

| Name | DNA Sequence of oligonucleotide |
|----------|--|
| L14735 | 5'-AAA AAC CAC CGT TGT TAT TCA ACT A-3' |
| H15149ad | 5'-GCI CCT CAR AAT GAY ATT TGT CCT CA-3' |

Table 2 — Oligonucleotides for amplification of the *cox1* gene region [2], [3]

| Name | DNA Sequence of oligonucleotide | | |
|--|--|--|--|
| VF2_t1 | 5'- TGT AAA ACG ACG GCC AGT CAA CCA ACC ACA AAG ACA TTG GCA C-3' | | |
| FishF2_t1 | 5'- TGT AAA ACG ACG GCC AGT CGA CTA ATC ATA AAG ATA TCG GCA C-3' | | |
| FishR2_t1 | 5'- CAG GAA ACA GCT ATG AC A CTT CAG GGT GAC CGA AGA ATC AGA A-3' | | |
| FR1d_t1 | 5'-CAG GAA ACA GCT ATG ACA CCT CAG GGT GTC CGA ARA AYC ARA A-3' | | |
| M13 tails of the primers are highlighted in bold and italic. | | | |

5.2.5 Agarose

https://standards.iteh.ai/catalog/standards/sist/ad380310-ca08-48cf-a25e-

5.2.6 Suitable DNA length standard for assessing the amplification product length

5.2.7 Sequencing primers [3] (see Table 3)

Table 3 — Sequencing primers for *cox1* PCR products

| Name | DNA Sequence of oligonucleotide |
|------------|---------------------------------|
| M13F (-21) | 5'-TGT AAA ACG ACG GCC AGT-3' |
| M13R (-27) | 5'-CAG GAA ACA GCT ATG AC-3' |

¹ During the collaborative study the Maxima® Hot Start PCR Master Mix (2 x) of Fermentas GmbH (ready to use PCR buffer solution including thermostable DNA polymerase) was used for the *cytb* amplification and the BIOTAQ DNA polymerase of Bioline with 10 x reaction buffer and separate MgCl₂ solution for the *cox1* amplification. In addition to the recommended BIOTAQ DNA polymerase other mastermixes and polymerases were successfully used in the collaborative study.

² Deoxynucleotide triphosphates can also be part of a commercial PCR master mix.

CEN/TS 17303:2019 (E)

6 Apparatus

Apart from the usual laboratory equipment, the following equipment is required:

- 6.1 UV-spectrophotometer or fluorometer, to determine the concentration of DNA
- 6.2 Thermocycler
- 6.3 Gel electrophoresis device
- 6.4 Gel documentation system
- 6.5 DNA sequencer

7 Procedure

7.1 Sample preparation

It should be ensured that the test portion used for DNA extraction is representative for the laboratory sample. In composed samples (e.g. ready-to-use meals), single pure fish pieces have to be separated and analysed. With the analysis of samples composed of several pieces (e.g. bags with different fillets), test portions for every putative fish species are taken and analysed separately. To minimize the risk of amplifying adhering contaminants, test sample material shall not be taken from the surface of the laboratory sample. For further information regarding sample preparation, see ISO 20813³.

(standards.iteh.ai)

7.2 DNA extraction

Concerning the extraction of DNA from the test sample, the general instructions and measures described in EN ISO 21571 should be followed, see ISO 208133 It is recommended to choose one of the DNA extraction methods described in EN ISO 21571:2005, Annex A⁴. Alternatively, commercial kits can be used for the extraction and purification of DNA.

7.3 PCR

7.3.1 General

The primers used for the amplification of the section from the mitochondrial *cytb* gene are universal primers. The primer L14735 binds to a section in the neighbouring highly conserved tRNA-Glu gene. The primer H15149ad is a universal primer with fish-specific adaptations [1]. It is currently known that with some exceptions the primer pair L14735/H15149ad did not react with samples labelled as Barramundi (*Lates calcarifer*) or Nile perch (*Lates niloticus*) [7].

The primers used for the amplification of the section from the mitochondrial *cox1* gene were designed to amplify a segment from the 5' region of the *cox1* gene of fish [2]. The primer pair has been tested against a very broad taxonomic range of fish species, and has only failed in a small minority of cases (< 5 % of species tested) [3].

³ Under development; currently at approval stage as ISO/FDIS 20813.

⁴ EN ISO 21571:2005 is currently impacted by EN ISO 21571:2005/A1:2013.