

---

---

**Microbiology — Cosmetics —  
Guidelines for the application of ISO  
standards on Cosmetic Microbiology**

*Microbiologie — Cosmétique — Lignes directrices pour l'application  
des normes ISO relatives à la microbiologie cosmétique*

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

[ISO/TR 19838:2016](https://standards.iteh.ai/catalog/standards/sist/d790fe4c-c70e-4861-8a77-8cbe665ee45e/iso-tr-19838-2016)

[https://standards.iteh.ai/catalog/standards/sist/d790fe4c-c70e-4861-8a77-  
8cbe665ee45e/iso-tr-19838-2016](https://standards.iteh.ai/catalog/standards/sist/d790fe4c-c70e-4861-8a77-8cbe665ee45e/iso-tr-19838-2016)



**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

ISO/TR 19838:2016

<https://standards.iteh.ai/catalog/standards/sist/d790fe4c-c70e-4861-8a77-8cbe665ee45e/iso-tr-19838-2016>



**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2016, Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Ch. de Blandonnet 8 • CP 401  
CH-1214 Vernier, Geneva, Switzerland  
Tel. +41 22 749 01 11  
Fax +41 22 749 09 47  
copyright@iso.org  
www.iso.org

# Contents

	Page
<b>Foreword</b> .....	<b>iv</b>
<b>Introduction</b> .....	<b>v</b>
<b>1 Scope</b> .....	<b>1</b>
<b>2 Terms and definitions</b> .....	<b>1</b>
<b>3 Microbial content</b> .....	<b>2</b>
3.1 General requirements.....	2
3.2 Enumeration of mesophilic microorganisms (bacteria, yeasts and moulds).....	3
3.3 Detection of specified microorganisms.....	3
3.4 Detection of specified and non-specified microorganisms.....	4
<b>4 Antimicrobial preservation</b> .....	<b>5</b>
4.1 General requirements.....	5
4.2 Evaluation of the preservation of a cosmetic formulation.....	5
4.3 Evaluation of the preservation of a cosmetic product.....	6
<b>5 Examples of microbial content results</b> .....	<b>6</b>
5.1 General.....	6
5.2 Eye make-up remover.....	6
5.3 Mascara.....	6
5.4 Face cream.....	7
5.5 Shampoo.....	8
<b>6 Examples of interpretation of preservation efficacy test results</b> .....	<b>8</b>
<b>Annex A (informative) Relationship between the ISO cosmetic microbiology standards</b> .....	<b>11</b>
<b>Annex B (informative) Relationship between the ISO standards dealing with the microbial content of cosmetics</b> .....	<b>12</b>
<b>Annex C (informative) Evaluation of the microbial content of a cosmetic product</b> .....	<b>13</b>
<b>Annex D (informative) Interpretation of test results</b> .....	<b>14</b>
<b>Bibliography</b> .....	<b>15</b>

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see [www.iso.org/directives](http://www.iso.org/directives)).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see [www.iso.org/patents](http://www.iso.org/patents)).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the WTO principles in the Technical Barriers to Trade (TBT) see the following URL: [Foreword - Supplementary information](#)

The committee responsible for this document is ISO/TC 217, *Cosmetics*.

ISO/TR 19838:2016

<https://standards.iteh.ai/catalog/standards/sist/d790fe4c-c70e-4861-8a77-8cbe665ee45e/iso-tr-19838-2016>

## Introduction

Every cosmetic manufacturer has a dual responsibility relative to the microbiological quality of its products.

- The first is to ensure that the product, as purchased, is free from the numbers and types of microorganisms that could affect product quality and consumer health. This is generally ensured by applying cosmetic good manufacturing practice (GMP) (see ISO 22716) during the manufacturing and packaging operations and, if necessary, by using **microbial content tests** on finished products.
- The second is to ensure that microorganisms introduced during normal product use will not adversely affect the quality or safety of the product. This is generally ensured by conducting **preservation efficacy tests** (or **challenge tests**) during the development stage of the new product.

In order to ensure product quality and safety for consumers, it is advisable that an appropriate microbiological risk analysis be performed to determine the types of cosmetic products to which this Technical Report would be applicable.

- Products considered to present a low microbiological risk are described in ISO 29621. These products identified as “hostile” and produced in compliance with GMP pose a very low overall risk to the user. Therefore, products that comply with the characteristics outlined in ISO 29621 do not require microbiological testing including both challenge test and end product testing.
- For those products which are not considered “hostile”, the microbiological quality has to be assessed by conducting tests with appropriate methods. ISO TC 217 provides a comprehensive set of standards to assess the antimicrobial preservation of cosmetic products and the microbiological quality of finished products (methods and limits). Manufacturers can decide not to test if they can demonstrate that their products comply with those requirements specified in ISO 17516 and/or ISO 11930.

[ISO/TR 19838:2016](https://standards.iteh.ai/catalog/standards/sist/d790fe4c-c70e-4861-8a77-8cbe665ee45e/iso-tr-19838-2016)

<https://standards.iteh.ai/catalog/standards/sist/d790fe4c-c70e-4861-8a77-8cbe665ee45e/iso-tr-19838-2016>

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

ISO/TR 19838:2016

<https://standards.iteh.ai/catalog/standards/sist/d790fe4c-c70e-4861-8a77-8cbe665ee45e/iso-tr-19838-2016>

# Microbiology — Cosmetics — Guidelines for the application of ISO standards on Cosmetic Microbiology

## 1 Scope

This Technical Report gives general guidelines to explain the use of ISO cosmetic microbiological standards depending on the objective (in-market control, product development, etc.) and the product to be tested.

This Technical Report can be used to fulfil the requirements of the ISO standard on microbiological limits (ISO 17516).

## 2 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

### 2.1

#### **cosmetic formulation**

preparation of raw materials with a qualitatively and quantitatively defined composition

### 2.2

#### **cosmetic product**

finished cosmetic product that has undergone all stages of production, including packaging in its final container for shipment

### 2.3

#### **sample**

one or more representative elements selected from a set to obtain information about that set

### 2.4.1

#### **microbial content**

<quantitative> estimated number of viable aerobic mesophilic microorganisms (bacteria, yeasts and moulds) within a cosmetic

### 2.4.2

#### **microbial content**

<qualitative> detectable specified or non-specified microorganisms within a cosmetic sample

### 2.5

#### **preservation of a cosmetic formulation**

set of means used to avoid microbial proliferation in a cosmetic formulation

EXAMPLE Preservatives, multifunctional compounds, hostile raw materials, extreme pH, low water activity values, etc.

### 2.6

#### **antimicrobial protection of a cosmetic product**

ability of a cosmetic product to overcome microbial contamination that might present a potential risk to the user

Note 1 to entry: The overall antimicrobial protection includes preservation of the formulation, the specific manufacturing process and protective packaging.

**2.7 microbiologically low-risk products**  
products whose environment denies microorganisms the physical and chemical requirements for growth and/or survival (hostile products)

Note 1 to entry: This category of low-risk products applies to microbiological contamination which may occur during manufacturing and/or use by the consumer.

Note 2 to entry: A product whose packaging prevents the ingress of microorganisms is considered a microbiological low-risk product during its use.

Note 3 to entry: The inclusion of preservatives or other antimicrobial compounds in a formulation by itself would not necessarily constitute a low-risk product.

**2.8 microbiological Risk Assessment (low risk products)**  
evaluation of product characteristics to determine if that product may be subject to microbial contamination

Note 1 to entry: These characteristics include the composition of the product, the production conditions, packaging and a combination of these factors (see ISO 29621).

### 3 Microbial content

#### 3.1 General requirements

There are eight International Standards dealing with the microbial content of cosmetic samples (see [Annex A](#)).

- ISO 16212
- ISO 17516
- ISO 18415
- ISO 18416
- ISO 21149
- ISO 21150
- ISO 22717
- ISO 22718

ITeH STANDARD PREVIEW  
(standards.iteh.ai)  
<https://standards.iteh.ai/catalog/standards/sist/d790fe4c-c70e-4861-8a77-8cbe665ee45e/iso-tr-19838-2016>

Additionally, ISO 21148 gives general instructions for carrying out microbiological examinations of cosmetic products.

Because of the large variety of cosmetic products within this field of application, the methods described in these International Standards may not be appropriate for some products in every detail (e.g. certain water immiscible products).

Other methods (e.g. automated) may be substituted for the tests presented here provided that their equivalence has been demonstrated or the method has been otherwise verified.

The possible inhibition of microbial growth by the sample should be neutralized to allow the detection of viable microorganisms. In all cases and whatever the methodology, the neutralization of the antimicrobial properties of the product should be checked and demonstrated.

The study of the neutralization of the antimicrobial activity should be performed when the tests have to be carried out on new products and whenever there is a change in the experimental conditions of the test.



The given International Standards will allow the following:

- the enumeration and detection of mesophilic microorganisms which may grow under aerobic conditions;
- the determination of the absence of specified microorganisms that are of interest for cosmetic products;

and, therefore, to estimate if the product under test complies with the requirements of the International Standard on microbiological limits (ISO 17516).

The test methods are described in the individual standards. The choice of a specific method, or combination of methods, will depend on the purpose for performing the test. It is up to the user to decide which approach is best for each application.

Depending on the expected level of contamination of the sample to be tested, two different approaches can be used (see [Annex A](#)).

- Quantitative tests (enumeration) are to be used when there is no information on the microbiological quality of the sample or if it is expected to be contaminated.
- Qualitative tests (detection) can be used if the sample is presumably free from microbial contamination (e.g. based on product history) and can be very useful from an economical and production time standpoint.

In the event where microorganisms are detected, the presence of specified microorganisms should be checked according to [3.3](#) or [3.4](#).

### 3.2 Enumeration of mesophilic microorganisms (bacteria, yeasts and moulds)

For bacteria, the enumeration method described in ISO 21149 involves enumeration of colonies on a non-selective agar medium. Enumeration of colonies can be performed by a plate count method or a filtration method using a specified culture medium incubated under defined conditions.

The results are expressed as follows:

“[number of] aerobic mesophilic bacteria per gram or per millilitre of product, expressed as cfu/g or cfu/ml”.

For yeasts and moulds, the method described in ISO 16212 involves enumeration of colonies by a plate count method or a filtration method using a specified culture medium with antibiotic incubated under defined conditions.

An alternative condition using the culture medium without antibiotic is proposed in ISO 16212.

The results are expressed as follows:

“[number of] yeast and mould per gram or per millilitre of product expressed, as cfu/g or cfu/ml”.

### 3.3 Detection of specified microorganisms

The detection of skin pathogens such as *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Candida albicans* may be relevant. The detection of other kinds of microorganisms might be of interest since these microorganisms (including indicators of fecal contamination e.g. *Escherichia coli*) suggest hygienic failure during the manufacturing process.

The methods are described in ISO 18416 (*C. albicans*), ISO 21150 (*E. coli*) ISO 22717 (*P. aeruginosa*), and ISO 22718 (*S. aureus*).

The main steps of the methods are the following:

- an enrichment using a non-selective broth medium incubated under defined conditions to increase the number of microorganisms while avoiding the risk of inhibition by the selective ingredients present in selective/differential growth media;
- then, if growth is detected, an isolation on a selective medium followed by identification tests.

If the identification of the colonies confirms the presence of the specified microorganism (*Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Candida albicans* or *Escherichia coli*) the result is expressed as follows:

“Presence of the specified microorganism (*name of the species*) in the sample *S*.”

If no growth after enrichment is observed and/or if the identification of the colonies does not confirm the presence of the specified microorganism (*Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Candida albicans* or *Escherichia coli*), the result is expressed as follows:

“Absence of the specified microorganism (*name of the species*) in the sample, *S*.”

### 3.4 Detection of specified and non-specified microorganisms

Two International Standards describe how to detect specified and non-specified microorganisms.

- The principle of ISO 18415 is to perform an enrichment by using a non-selective broth incubated under defined conditions to increase the number of microorganisms without the risk of inhibition by the selective ingredients that are present in selective/differential growth media followed by isolation on a non-selective medium incubated under defined conditions and identification conducted according to need by using appropriate incubation conditions and suitable identification tests.

For each species of specified microorganism, and if the identification of the colonies confirms the presence of this species, the result is expressed as follows:

“Presence of (*name of the species*) in the sample, *S*”  
<https://standards.iteh.ai/catalog/standards/sist/d790fe4c-c70e-4861-8a77-6c0663c47c96/iso-tr-19838-2016>

If growth is observed after enrichment and if the colonies are isolated and recognized as non-specified microorganisms, the result is expressed as follows:

“Presence of (*name of the species* or main morphological characteristics) in the sample, *S*, and absence of specified microorganisms”.

If no growth after enrichment is observed, the result is expressed as follows:

“Absence of aerobic mesophilic bacteria and yeast (specified microorganisms included) in the sample, *S*”.

- ISO 21149 describes a method for the detection of aerobic mesophilic bacteria.

The enrichment method described in ISO 21149 consists of incubation under defined conditions of a defined quantity of the sample (*S*) in a non-selective broth containing suitable neutralizers and/or dispersing agents followed by a transfer of a defined quantity of the previous suspension on non-selective solid agar medium under defined conditions. The difference between ISO 18415 and ISO 21149 is that the neutralization of the antimicrobial activity in ISO 21149 is only checked for bacteria, while ISO 18415 refers to both bacteria and yeast.

Therefore, if no growth after enrichment is observed, the results are expressed as follows:

“Absence of aerobic mesophilic bacteria (specified bacteria included) in the sample *S*”.

If growth is observed after enrichment, the results are expressed as:

“Presence of aerobic mesophilic bacteria in the sample *S*”.

NOTE Enrichment methods are not appropriate (temperature, culture media) to detect moulds. Therefore, even if no growth after enrichment is observed, it is necessary to look for moulds using appropriate culture conditions (see ISO 16212).

## 4 Antimicrobial preservation

### 4.1 General requirements

The antimicrobial protection of a cosmetic product (cosmetic formulation in its final container) can come from many sources, such as the following:

- chemical preservation;
- inherent characteristics of the formulation;
- package design;
- manufacturing process

When evaluating the overall antimicrobial protection of a cosmetic product, these different sources should be taken into account in a microbiological risk assessment.

ISO 11930 describes a procedure for the interpretation of data generated by the preservation efficacy test (if appropriate) and by the microbiological risk assessment

### 4.2 Evaluation of the preservation of a cosmetic formulation

A preservation efficacy test or challenge test is commonly used to evaluate the preservation of a cosmetic formulation. This test is not required for those cosmetic products for which the microbiological risk has been determined to be low (see ISO 29621).

This test is primarily designed for water-soluble or water-miscible cosmetic products and can require adaptation, for example, to test products in which water is the internal phase. The test described in ISO 11930 involves, for each test microorganism, placing the formulation in contact with a calibrated inoculum, and then measuring the changes in the microorganism count at set time intervals for a set period and at a set temperature.

As for the microbial content tests, the possible inhibition of microbial growth by the sample should be neutralized to allow the detection of viable microorganism. In all cases and whatever the methodology, the neutralization of the antimicrobial properties of the product should be checked and demonstrated.

The microorganism counts are converted in log reduction values and compared with 2 sets of criteria expressed as minimum log reduction.

- Criteria A, whereby the formulation is protected against microbial proliferation that may present a potential risk for the user and no additional factors are considered.
- Criteria B, whereby the level of protection is acceptable if the risk analysis demonstrates the existence of control factors not related to the formulation indicating that the microbiological risk is tolerable for the cosmetic product.

The inherent variability in microbial counts should be taken into consideration when comparing the obtained values and the preset criteria A or B. A deviation of 0,5 log units from the preset criteria is considered acceptable.