

---

---

**Microbiology of the food chain —  
Horizontal method for the detection  
and enumeration of *Clostridium*  
spp. —**

Part 1:

**Enumeration of sulfite-reducing  
*Clostridium* spp. by colony-count  
technique**

*Microbiologie de la chaîne alimentaire — Méthode horizontale pour  
la recherche et le dénombrement de *Clostridium* spp. —*

*Partie 1: Dénombrement des bactéries *Clostridium* spp. sulfito-  
réductrices par la technique de comptage des colonies*



iTeh STANDARD PREVIEW  
(standards.iteh.ai)

[ISO 15213-1:2023](https://standards.iteh.ai/catalog/standards/sist/1132d699-4297-4fd0-8e1a-5932712cc27b/iso-15213-1-2023)

<https://standards.iteh.ai/catalog/standards/sist/1132d699-4297-4fd0-8e1a-5932712cc27b/iso-15213-1-2023>



**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2023

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
CP 401 • Ch. de Blandonnet 8  
CH-1214 Vernier, Geneva  
Phone: +41 22 749 01 11  
Email: [copyright@iso.org](mailto:copyright@iso.org)  
Website: [www.iso.org](http://www.iso.org)

Published in Switzerland

# Contents

	Page
Foreword.....	iv
Introduction.....	vi
<b>1 Scope.....</b>	<b>1</b>
<b>2 Normative references.....</b>	<b>2</b>
<b>3 Terms and definitions.....</b>	<b>2</b>
<b>4 Principle.....</b>	<b>2</b>
4.1 General.....	2
4.2 Preparation of dilutions.....	3
4.3 Enumeration.....	3
4.4 Confirmation.....	3
<b>5 Culture media and reagents.....</b>	<b>3</b>
<b>6 Equipment and consumables.....</b>	<b>3</b>
<b>7 Sampling.....</b>	<b>4</b>
<b>8 Preparation of test sample.....</b>	<b>4</b>
<b>9 Procedure.....</b>	<b>5</b>
9.1 General.....	5
9.2 Test portion, initial suspension and dilutions.....	5
9.3 Heat treatment to select spores.....	5
9.4 Inoculation and incubation.....	5
9.5 Enumeration of typical colonies.....	6
9.6 Confirmation of sulfite-reducing <i>Clostridium</i> spp.....	6
<b>10 Expression of results.....</b>	<b>7</b>
<b>11 Validation of the method.....</b>	<b>7</b>
11.1 Interlaboratory study.....	7
11.2 Performance characteristics.....	7
<b>12 Test report.....</b>	<b>8</b>
<b>13 Quality assurance.....</b>	<b>8</b>
<b>Annex A (normative) Flow diagram of the procedure.....</b>	<b>9</b>
<b>Annex B (normative) Culture media and reagents.....</b>	<b>11</b>
<b>Annex C (informative) Performance characteristics of the method.....</b>	<b>14</b>
<b>Annex D (informative) Special protocol for the enumeration of sulfite-reducing <i>Clostridium</i> spp. in feed.....</b>	<b>17</b>
<b>Bibliography.....</b>	<b>21</b>

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see [www.iso.org/directives](http://www.iso.org/directives)).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see [www.iso.org/patents](http://www.iso.org/patents)).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see [www.iso.org/iso/foreword.html](http://www.iso.org/iso/foreword.html).

This document was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 9, *Microbiology*, in collaboration with the European Committee for Standardization (CEN) Technical Committee CEN/TC 463, *Microbiology of the food chain*, in accordance with the Agreement with technical cooperation between ISO and CEN (Vienna Agreement).

This first edition of ISO 15213-1 cancels and replaces ISO 15213:2003, which has been technically revised.

The main changes are as follows:

- the Scope has been expanded to include samples from the primary production stage;
- the scope of the method has been changed from “sulfite-reducing bacteria” to “sulfite-reducing *Clostridium* spp.”: therefore, typical colonies on the iron sulfite agar plates are confirmed;
- the concentration of sulfite in the iron sulfite agar has been reduced from 1,0 g/l to 0,5 g/l;
- the heat treatment of 10 min at 80 °C has been made optional, in the case of high background flora or for the enumeration of only spores of sulfite-reducing *Clostridium* spp. present in the sample;
- the option for using tubes for inoculation has been removed;
- the option for incubating the samples at 50 °C for the enumeration of thermophilic sulfite-reducing bacteria has been removed;
- a description of how the confirmation of typical colonies has to be performed has been added;
- the flow diagram in [Annex A](#) giving a short description of the procedure has been revised;
- in [Annex C](#), the performance characteristics have been added;
- [Annex D](#) has been added to provide a special protocol for the enumeration of sulfite-reducing *Clostridium* spp. in feed.

A list of all parts in the ISO 15213 series can be found on the ISO website.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at [www.iso.org/members.html](http://www.iso.org/members.html).

**iTeh STANDARD PREVIEW**  
**(standards.iteh.ai)**

[ISO 15213-1:2023](https://standards.iteh.ai/catalog/standards/sist/1132d699-4297-4fd0-8e1a-5932712cc27b/iso-15213-1-2023)

<https://standards.iteh.ai/catalog/standards/sist/1132d699-4297-4fd0-8e1a-5932712cc27b/iso-15213-1-2023>

## Introduction

Sulfite-reducing *Clostridium* spp. are obligate anaerobic, Gram-positive, spore-forming, rod-shaped bacteria. The most important species which belong to this group are *Clostridium (C.) perfringens*, *C. bifermentans*, *C. sporogenes* and *C. botulinum*. Some species can cause foodborne illness. As ubiquitous bacteria they are predominantly found in nature. The *Clostridium* species inhabit soils and the intestinal tract of animals and humans.

Sulfite-reducing *Clostridium* spp., including *C. perfringens*, are widely used as microbial indicators of clostridial contamination in the manufacturing of foods (e.g. meat production). These have the capacity to produce heat-resistant spores. Outside the dairy industry, the use of sulfite-reducing *Clostridium* spp. as a microbial indicator is limited to a relatively small number of foods. Its current application in non-dairy foods is either an indication of faecal contamination (especially *C. perfringens*, see also ISO 15213-2 and ISO/TS 15213-3) and/or as an indicator of sanitation/process control related to potential growth and survival of anaerobic spore-forming bacteria.

This document describes the horizontal method for the enumeration of sulfite-reducing *Clostridium* spp. in food, feed, environmental samples and samples from the primary production stage. The method for the enumeration of *C. perfringens* is described in ISO 15213-2. The method for the detection of *C. perfringens* is described in ISO/TS 15213-3. These three parts are published as a series of International Standards because the methods are closely linked to each other. These methods are often conducted in association with each other in a laboratory, and the media and their performance characteristics can be similar.

The main technical changes listed in the Foreword, introduced in this document compared with ISO 15213:2003, are considered as major (see ISO 17468).

These changes have a major impact on the performance characteristics of the method.

[ISO 15213-1:2023](https://standards.iteh.ai/catalog/standards/sist/1132d699-4297-4fd0-8e1a-5932712cc27b/iso-15213-1-2023)

<https://standards.iteh.ai/catalog/standards/sist/1132d699-4297-4fd0-8e1a-5932712cc27b/iso-15213-1-2023>

# Microbiology of the food chain — Horizontal method for the detection and enumeration of *Clostridium* spp. —

## Part 1:

## Enumeration of sulfite-reducing *Clostridium* spp. by colony-count technique

**WARNING** — In order to safeguard the health of laboratory personnel, it is essential that tests for the enumeration of sulfite-reducing *Clostridium* spp. are only undertaken in properly equipped laboratories, under the control of a skilled microbiologist, and that great care is taken in the disposal of all incubated materials. Persons using this document should be familiar with normal laboratory practice. This document does not purport to address all of the safety aspects, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices.

### 1 Scope

This document specifies the enumeration of sulfite-reducing *Clostridium* spp. by the colony-count technique.

This document is applicable to:

- products intended for human consumption;
- products for feeding animals;
- environmental samples in the area of food and feed production and handling;
- samples from the primary production stage.

**NOTE** This method has been validated in an interlaboratory study for the following food categories:

- ready-to-eat, ready-to-reheat meat products;
- eggs and egg products (derivates);
- processed fruits and vegetables;
- infant formula and infant cereals;
- multi-component foods or meal components.

It has also been validated for the following other categories:

- pet food and animal feed;
- environmental samples (food or feed production).

As this method has been validated for at least five food categories, this method is applicable for a broad range of food. For detailed information on the validation, see [Clause 11](#) and [Annex C](#). Since the method is not commonly used for samples in the primary production stage, this category was not included in the interlaboratory study. Therefore, no performance characteristics were obtained for this category.

This horizontal method was originally developed for the examination of all samples belonging to the food chain. Based on the information available at the time of publication of this document, this method is considered to be fully suited to the examination of all samples belonging to the food chain. However, because of the large variety of products in the food chain, it is possible that this horizontal method is not

appropriate in every detail for all products. Nevertheless, it is expected that the required modifications are minimized so that they do not result in a significant deviation from this horizontal method.

This technique is suitable for, but not limited to, the enumeration of microorganisms in test samples with a minimum of 10 colonies counted on a plate. This corresponds to a level of contamination that is expected to be higher than 10 cfu/ml for liquid samples or higher than 100 cfu/g for solid samples.

## 2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 6887 (all parts), *Microbiology of the food chain — Preparation of test samples, initial suspension and decimal dilutions for microbiological examination*

ISO 7218, *Microbiology of the food chain — General requirements and guidance for microbiological examinations*

ISO 11133, *Microbiology of food, animal feed and water — Preparation, production, storage and performance testing of culture media*

ISO 19036:2019, *Microbiology of the food chain — Estimation of measurement uncertainty for quantitative determinations*

## 3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminology databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <https://www.electropedia.org/>

### 3.1 sulfite-reducing *Clostridium* spp.

genus of microorganisms of the family of *Clostridiaceae*, usually capable of growth in/on iron sulfite agar (ISA) under anaerobic conditions, forming typical or less typical colonies, and displaying certain characteristics with biochemical confirmation tests

Note 1 to entry: The biochemical confirmation tests are described in [9.6](#).

### 3.2 enumeration of sulfite-reducing *Clostridium* spp.

determination of the number of colony-forming units (cfu) of *sulfite-reducing Clostridium* spp. ([3.1](#)) bacteria per gram, per millilitre, per square centimetre or per sampling device when a specified test is conducted

Note 1 to entry: Specified tests are given in [Clause 9](#).

## 4 Principle

### 4.1 General

A specified quantity of the liquid test sample, or of an initial suspension in the case of other products, is dispensed into an empty Petri dish and mixed well with a specified molten agar culture medium to form a poured plate. Other plates are prepared under the same conditions using decimal dilutions of the test sample. After solidification of the agar culture medium, an overlay is used to prevent the



development of spreading colonies on the surface of the medium. If it is the intention to count only spores, a heat treatment of 10 min at 80 °C needs to be performed before plating.

When the number of cfu is expected to be at or near the limit of determination, the use of duplicate plates is preferable. If duplicate plates are used, the minimum for the sum of colonies on both plates should be 10 colonies. In this case, the level of contamination is expected to be higher than 5 cfu/ml for liquid samples or higher than 50 cfu/g for solid samples.

A pour-plate technique with overlay is especially suited for the enumeration of products expected to contain spreading colonies that can obscure colonies of the target microorganisms.

The enumeration of sulfite-reducing *Clostridium* spp. requires four successive stages as specified in [Annex A](#).

## 4.2 Preparation of dilutions

For the preparation of decimal dilutions from the test portion, follow the procedure as specified in the ISO 6887 series.

## 4.3 Enumeration

The plates are incubated under anaerobic conditions at 37 °C for 48 h. After incubation, the number of typical colonies, which show black or grey to yellow-brown staining, are counted. The colour of the colonies and the surrounding zone changes due to the formation of iron(II)sulfide as a result of the reaction between sulfide ions and trivalent iron [Fe(III)] present in the medium.

## 4.4 Confirmation

Typical colonies are picked for confirmation.

NOTE When no confirmation is performed, the results can be reported as “anaerobic sulfite-reducing bacteria”.

## 5 Culture media and reagents

Follow current laboratory practices in accordance with ISO 7218. The composition of culture media and reagents and their preparation are specified in [Annex B](#). For performance testing of culture media, follow the procedures in accordance with ISO 11133 and [Annex B](#).

## 6 Equipment and consumables

Disposable equipment is an acceptable alternative to reusable glassware if it has suitable specifications. The usual microbiological laboratory equipment (see ISO 7218) and, in particular, the following shall be used.

**6.1 Appropriate apparatus for achieving an anaerobic atmosphere**, a jar that can be hermetically sealed or any other appropriate equipment which enables anaerobic atmosphere conditions to be maintained for the total incubation time of the culture medium. Other systems of equivalent performance, such as anaerobic cabinets, may be used. Follow the manufacturer’s instructions for installation and maintenance.

The composition of the atmosphere required can be achieved by means of the addition of a gas mixture (e.g. from a gas cylinder) after evacuation of air from the jar, by displacement of the atmosphere in a cabinet or by any other appropriate means (such as commercially available gas packs). In general, anaerobic incubation requires an atmosphere of less than 1 % volume fraction oxygen, 9 % volume fraction to 13 % volume fraction carbon dioxide.

**6.2 Apparatus for dry sterilization (oven) or wet sterilization (autoclave).**

**6.3 Incubator**, capable of operating at  $37\text{ °C} \pm 1\text{ °C}$ .

**6.4 pH-meter**, having an accuracy of calibration of  $\pm 0,1$  pH unit at  $25\text{ °C}$ .

**6.5 Refrigerator**, capable of operating at  $5\text{ °C} \pm 3\text{ °C}$ .

**6.6 Sterile bottles, flasks or tubes**, of appropriate capacity. Bottles, flasks or tubes with non-toxic metallic or plastic screwcaps may be used.

**6.7 Sterile graduated pipettes or automatic pipettes**, of nominal capacities of 10 ml and 1 ml.

**6.8 Sterile loops**, of approximately 1  $\mu\text{l}$  volume, or inoculation needle or wire.

**6.9 Sterile Petri dishes**, with a diameter of approximately 90 mm and (optional) large size (diameter approximately 140 mm).

**6.10 Thermostatically controlled water bath**, capable of operating at  $44\text{ °C}$  to  $47\text{ °C}$  and  $80\text{ °C} \pm 2\text{ °C}$ .

## 7 Sampling

Sampling is not part of the method specified in this document. Follow the specific International Standard dealing with the product concerned. If there is no specific International Standard dealing with the sampling of the product concerned, it is recommended that the parties concerned come to an agreement on this subject.

Recommended sampling techniques are given in the following documents:

- ISO/TS 17728 for food and animal feed;
- ISO 707 for milk and milk products;
- ISO 6887-3 for raw molluscs, tunicates and echinoderms from primary production areas;
- ISO 13307 for primary production stage;
- ISO 17604 for carcasses;
- ISO 18593 for surfaces.

It is important that the laboratory receives a sample that is representative of the product under consideration. The sample should not have been damaged or changed during transport or storage.

## 8 Preparation of test sample

Prepare the test sample from the laboratory sample in accordance with the specific International Standard dealing with the product concerned. Follow the procedures as specified in the ISO 6887 series. If there is no specific International Standard available, it is recommended that the parties concerned come to an agreement on this subject.

## 9 Procedure

### 9.1 General

A flow diagram of the procedure is given in [Annex A](#).

### 9.2 Test portion, initial suspension and dilutions

Follow the procedures in accordance with the ISO 6887 series and the specific International Standard dealing with the product concerned.

Prepare a single decimal dilution series from the test portion if the product is liquid, or from the initial suspension in the case of other products.

A special protocol for preparing the initial suspension of feed samples is given in [Annex D](#).

### 9.3 Heat treatment to select spores

If it is the intention to count only spores, heat the decimal dilution series to 80 °C in a water bath ([6.10](#)) for 10 min ± 1 min. Heat treatment shall be given within 15 min after preparation of the initial suspension to avoid germination of spores. If the tube is not placed in the water bath within 15 min, it should be placed immediately in melting ice for a maximum of 2 h.

The temperature during heat treatment should be monitored by placing an appropriate thermometer in a reference bottle of the same size as the sample bottle and containing the same volume of water at the same initial temperature as the sample being treated ([6.6](#)). The tubes should not be hermetically sealed during the heat treatment. The time taken to reach 80 °C shall not exceed 5 min and can be minimized by ensuring the water level to be at least 4 cm above the level of the sample and equipping the water bath with a circulating-water pump to maximize heat exchange.

Start the time of heating (10 min) when the temperature of the reference sample has reached 80 °C. After heat treatment, the samples should be cooled immediately until approximately 20 °C.

Heat treatment should also reduce the competitive flora in some matrices containing a high level of background flora (e.g. liquid whey, feed silage).

### 9.4 Inoculation and incubation

**9.4.1** Take two sterile Petri dishes with a diameter of approximately 90 mm ([6.9](#)). Transfer to each dish, by means of a sterile pipette ([6.7](#)), 1 ml of the test sample if liquid, or 1 ml of the initial suspension ( $10^{-1}$  dilution) in the case of other products. If plates from more than one dilution are prepared, this may be reduced to one dish (see ISO 7218).

When, for certain products, it is necessary to estimate low numbers of sulfite-reducing *Clostridium* spp., the limit of enumeration may be lowered by a factor of 10 by examining 10 ml of the initial suspension in three large (140 mm) Petri dishes ([6.9](#)).

**9.4.2** Take one other sterile Petri dish ([6.9](#)). Use another sterile pipette ([6.7](#)) to dispense 1 ml of the  $10^{-1}$  dilution (liquid product) or 1 ml of the  $10^{-2}$  dilution (other products).

**9.4.3** If necessary, repeat the procedure with further dilutions, using a new sterile pipette ([6.7](#)) for each decimal dilution.

**9.4.4** If appropriate and possible, select only the critical dilution steps (at least two consecutive decimal dilutions) for the inoculation of the Petri dishes ([6.9](#)) that will give colony counts of between 10 and 150 colonies per plate (on 90 mm Petri dishes) or between 10 and 360 colonies per plate (on 140 mm Petri dishes).