
**Soil quality — Measurement of
enzyme activity patterns in soil
samples using colorimetric substrates
in micro-well plates**

*Qualité du sol — Mesure de l'activité enzymatique dans des
échantillons de sol en utilisant des substrats colorimétriques*

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[ISO 20130:2018](https://standards.iteh.ai/catalog/standards/sist/917b2d5d-4862-446a-a3d3-b6918ed5a4ec/iso-20130-2018)

[https://standards.iteh.ai/catalog/standards/sist/917b2d5d-4862-446a-a3d3-
b6918ed5a4ec/iso-20130-2018](https://standards.iteh.ai/catalog/standards/sist/917b2d5d-4862-446a-a3d3-b6918ed5a4ec/iso-20130-2018)



iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 20130:2018

<https://standards.iteh.ai/catalog/standards/sist/917b2d5d-4862-446a-a3d3-b6918ed5a4ec/iso-20130-2018>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2018

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Fax: +41 22 749 09 47
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

Contents

Page

Foreword.....	iv
Introduction.....	v
1 Scope.....	1
2 Normative references.....	1
3 Terms and definitions, symbols and abbreviated terms.....	1
3.1 Terms and definitions.....	1
3.2 Symbols and abbreviated terms.....	1
4 Principle.....	2
5 Reactives.....	2
5.1 Buffers and reagents.....	2
5.2 Substrates and standards.....	5
5.2.1 Preparation of standard solutions.....	5
5.2.2 Preparation of substrate solutions.....	5
6 Apparatus and materials.....	7
7 Procedure.....	8
7.1 Establishment of calibration curves.....	8
7.1.1 General.....	8
7.1.2 Solution of PNP.....	8
7.1.3 Solution of β -naphthylamine.....	8
7.1.4 Solution of ammonium chloride.....	8
7.2 Sampling.....	9
7.2.1 Sample preparation.....	9
7.2.2 Addition of substrate.....	10
7.2.3 Absorbance measurements.....	11
7.2.4 Measurements of urease activities.....	12
8 Calculation of results.....	12
9 Expression of results.....	12
10 Validity criteria.....	12
11 Interlaboratory validation.....	13
12 Test report.....	13
Annex A (informative) Validation of the method and intralaboratory tests for evaluating soil enzymatic activities with colorimetric method.....	14
Annex B (informative) International ring test for evaluating soil enzymatic activities with colorimetric method.....	19
Bibliography.....	28

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see the following URL: www.iso.org/iso/foreword.html. (standards.iteh.ai)

This document was prepared by Technical Committee ISO/TC 190, *Soil quality*, Subcommittee SC 4, *Biological characterization*.

ISO 20130:2018
<https://standards.iteh.ai/catalog/standards/sist/917b2d5d-4862-446a-a3d3-b6918ed5a4ec/iso-20130-2018>

Introduction

Microorganisms are responsible for many key processes in the cycle of elements. Enzymes are responsible for the degradation of organic molecules and their mineralization. The main postulate is the microbial origin of soil enzymes, even if plant root exudates include enzymes. Extracellular enzymes in soil play key roles in the biodegradation of organic macromolecules. The simultaneous monitoring of several enzyme activities important in the biodegradation of organic compounds and mineralization of carbon, nitrogen, phosphorus and sulfur in soil may reveal harmful effects caused by chemicals and other anthropogenic impacts. However, the measurements carried out under selected laboratory conditions using artificial substrates cannot be a substitute for the actual rate of enzymatic processes in soil *in situ*.

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 20130:2018

<https://standards.iteh.ai/catalog/standards/sist/917b2d5d-4862-446a-a3d3-b6918ed5a4ec/iso-20130-2018>

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 20130:2018

<https://standards.iteh.ai/catalog/standards/sist/917b2d5d-4862-446a-a3d3-b6918ed5a4ec/iso-20130-2018>

Soil quality — Measurement of enzyme activity patterns in soil samples using colorimetric substrates in micro-well plates

1 Scope

This document specifies a method for the measurement of several hydrolase activities (arylamidase, arylsulfatase, β -galactosidase, α -glucosidase, β -glucosidase, N-acetyl-glucosaminidase, acid, alkaline and global phosphatases, urease) simultaneously (or not) in soil samples, using colorimetric substrates. Enzyme activities of soil vary seasonally and depend on soil chemical, physical and biological characteristics. This method can be applied either to detect harmful effects on soil enzyme activities derived from toxic substances or other anthropogenic agents in contaminated soils against a control soil, or to test chemicals.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 18400-206, *Soil quality — Sampling — Part 206: Collection, handling and storage of soil under aerobic conditions for the assessment of microbiological processes, biomass and diversity in the laboratory*

3 Terms and definitions, symbols and abbreviated terms

3.1 Terms and definitions

No terms and definitions are listed in this document.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <http://www.electropedia.org/>
- ISO Online browsing platform: available at <https://www.iso.org/obp>

3.2 Symbols and abbreviated terms

ARN	Arylamidase
ARS	Arylsulfatase
E.C.	Enzyme code number by the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology (NC-IUBMB)
NAG	N-acetyl-glucosaminidase
PAC	acid phosphatase
PAK	alkaline phosphatase
PHOS	phosphatase

URE	urease
β -GAL	β -galactosidase
α -GLU	α -glucosidase
β -GLU	β -glucosidase

4 Principle

This document describes a method for the simultaneous measurement of several enzymes in soil samples (see [Table 1](#)). It is based on the use of soil samples solutions and colorimetric substrates, which are incubated during specific times at $25\text{ °C} \pm 2\text{ °C}$ or $37\text{ °C} \pm 2\text{ °C}$ in multi-well plates. After the incubation, reactions are stopped, plates are then centrifuged and supernatants transferred into new plates. The intensities of the coloration are measured with absorbance with a 96 wells microplate spectrophotometer UV/visible.

Table 1 — Enzymatic activity measurements with colorimetric method

Enzyme	Abbreviation	N°	Soil cycle	Macromolecule degraded
Arylamidase	ARN	E.C. 3.4.11.2	Nitrogen	
Arylsulfatase	ARS	E.C. 3.1.6.1	Sulfur	Mineralization of organic sulfur
β -Galactosidase	β GAL	E.C. 3.2.1.22	Carbon	Hemicellulose
α -Glucosidase	α GLU	E.C. 3.2.1.20	Carbon	Starch and glycogen
β -Glucosidase	β GLU	E.C. 3.2.1.21	Carbon	Cellulose
N-acetyl-glucosaminidase	NAG	E.C. 3.2.1.52	Carbon	Chitin and other β -1,4-linked glucosamine polymers
Phosphatase	PHOS	E.C. 3.1.4.1	Phosphorus	Phosphate esters
Acid phosphatase	PAC	E.C. 3.1.4.1	Phosphorus	Phosphate esters
Alkaline phosphatase	PAK	E.C. 3.1.4.1	Phosphorus	Phosphate esters
Urease	URE	E.C. 3.5.1.5	Nitrogen	Urea

An interlaboratory trial was carried out for the validation of the standard; summary of the international ring test is given in [Table 8](#), and the whole data of the interlaboratory validation are described in [Annex B](#).

5 Reactives

5.1 Buffers and reagents

5.1.1 General

The choice is made to use deionized water as medium to evaluate native soil enzyme activities at soil pH and also to allow the analysis of multiple enzymes using the same soil suspension. The soil (in g)/water (in ml) ratio (4:25) is optimized to maximize reaction, sensitivity and facilitate pipetting technique. The use of the same soil solution for analysing multiple enzymes also makes data more comparable. Arylamidase is measured with Tris buffer 50 mmol/l, pH 7,5 and acid and alkaline phosphatases are involved with the use of Tris-HCl 50 mmol/l at pH 5,5 and Tris base 50 mmol/l at pH 11, respectively.

NOTE The volume can be adapted according to needs.

5.1.2 Tris hydrochloride 50 mmol/l pH 5,5 \pm 0,1.

— Tris(hydroxymethyl)aminomethane hydrochloride (CAS N°: 1185-53-1 – Mw:157,6): 7,88 g;

- deionized water: 1 000 ml;
- hydrochloric acid (HCl) (CAS N°7647-01-0) 1 mol/l.

Dissolve 7,88 g of Tris(hydroxymethyl)aminomethane hydrochloride into 800 ml deionized water and adjust to pH 5,5 with hydrochloric acid (1 mol/l). Fill in to 1 000 ml. The storage duration shall not exceed one month at $4\text{ °C} \pm 2\text{ °C}$ in glass or polypropylene bottle.

5.1.3 Tris base 50 mmol/l pH $11 \pm 0,1$.

- Tris(hydroxymethyl)aminomethane (CAS N°: 77-86-1 - Mw:121,14): 6,06 g;
- deionized water: 1 000 ml;
- sodium hydroxide (CAS N° 1310-73-2) (1 mol/l).

Dissolve 6,06 g of Tris(hydroxymethyl)aminomethane into 800 ml deionized water and adjust to pH 11 with sodium hydroxide (1 mol/l). Fill in to 1 000 ml. The storage duration shall not exceed one month at $4\text{ °C} \pm 2\text{ °C}$.

5.1.4 Tris base 50 mmol/l pH $7,5 \pm 0,1$.

- Tris(hydroxymethyl)aminomethane (CAS N°: 77-86-1 - Mw:121,14): 6,06 g;
- deionized water: 1 000 ml;
- hydrochloric acid (HCl) (CAS N°7647-01-0) 1 mol/l.

Dissolve 6,06 g of Tris(hydroxymethyl)aminomethane into 800 ml deionized water and adjust to pH 7,5 with hydrochloric acid (1 mol/l). Fill in to 1 000 ml. The storage duration shall not exceed one month at $4\text{ °C} \pm 2\text{ °C}$.

iTech STANDARD PREVIEW
(standards.iteh.ai)
ISO 20130:2018
<https://standards.iteh.ai/catalog/standards/sist/917b2d5d-4862-446a-a3d3-12918016a4ec/iso-20130-2018>

5.1.5 Tris base 100 mmol/l pH $12 \pm 0,1$.

- Tris(hydroxymethyl)aminomethane (CAS N°: 77-86-1 - Mw:121,14): 12,11 g;
- deionized water: 1 000 ml;
- sodium hydroxide (CAS N° 1310-73-2) (5 mol/l).

Dissolve 12,11 g of Tris(hydroxymethyl)aminomethane into 800 ml deionized water and adjust to pH 12 with sodium hydroxide (5 M). Fill in to 1 000 ml. The storage duration shall not exceed one month at $4\text{ °C} \pm 2\text{ °C}$.

5.1.6 Calcium chloride dihydrate 0,5 mol/l.

- calcium chloride dihydrate (CAS N°: 10035-04-8 - Mw:147,01): 14,7 g;
- Deionized water: 200 ml.

Dissolve 14,7 g of calcium chloride dihydrate in 200 ml of deionized water. The storage duration shall not exceed one month at $4\text{ °C} \pm 2\text{ °C}$.

5.1.7 Salicylate reagent.

- sodium salicylate 270 mmol/l (CAS N°: 54-21-7 - Mw:160,1): 865 mg;
- tri sodium citrate 145 mmol/l (CAS N°: 6132-04-3 - Mw:294,1): 853 mg;
- di sodium tartrate 60 mmol/l (CAS N°: 6106-24-7 - Mw:230,08): 276 mg;
- sodium nitroferricyanide 2 mmol/l (CAS N°: 13755-38-9 - Mw:297,95): 12 mg;

- deionized water: 20 ml.

Salicylate reagent is prepared with the 4 compounds listed above just before analysis; dissolve 865 mg of sodium salicylate, 853 mg of tri sodium citrate, 276 mg of di sodium tartrate and 12 mg of sodium nitroferricyanide in 20 ml of deionized water.

5.1.8 Cyanurate reagent.

- tri sodium citrate 580 mmol/l (CAS N°: 6132-04-3 - Mw:294,1): 3,4 g;
- di sodium tartrate 90 mmol/l (CAS N°: 6106-24-7 - Mw:230,08): 414 mg;
- lithium hydroxide 280 mmol/l (CAS N° : 1310-65-2 - Mw:23,95): 134 mg;
- dichloroisocyanurate 10 mmol/l (CAS N° : 51580-86-0 - Mw:255,98): 51 mg;
- deionized water: 20 ml.

Cyanurate reagent is prepared with the 4 compounds listed above just before analysis; dissolve 3,4 g of tri sodium citrate, 414 mg of di sodium tartrate, 134 mg of lithium hydroxide and 51 mg of dichloroisocyanurate in 20 ml of deionized water.

5.1.9 Ethanol, 96 %.

- Ethanol 96 % (CAS N° 41340-36-7).

5.1.10 Acidified ethanol (0,26 mol/l HCl).

- Hydrochloric acid ACS reagent, 37 % (CAS N°7647-01-0) 4,32 ml;
- Ethanol 96 % (CAS N° 41340-36-7).

Dilute 4,32 mL of concentrated HCl into 200 ml ethanol 96%. The storage duration shall not exceed one month at 4 °C ± 2 °C.

5.1.11 p-dimethylaminocinnamaldehyde (DMCA) (3,5 mmol/l).

- DMCA (CAS N°: 6203-18-5 - Mw:175,23): 0,12 g;
- Ethanol 96 % (CAS N° 41340-36-7).

Dissolve 0,12 g of DMCA into 200 ml ethanol 96 %. The storage duration shall not exceed one week at -20 °C ± 2 °C.

Table 2 — Buffer utilization for enzymatic activity measurement

	ARS; α -GLU; β -GLU; β -GAL; NAG; PHOS;	ARN	URE	PAC	PAK
Soil solution	deionized water	Tris base 50 mmol/l, pH 7,5	deionized water	Tris HCl 50 mmol/l, pH 5,5	Trisbase 50 mmol/l, pH 11
Stop/ revelation	Tris 100 mmol/l pH12	Ethanol 96 % Acidified ethanol	salicylate reagent	Tris 100 mmol/l pH 12	
	CaCl ₂ 0,5 mol/l	DMCA	cyanurate reagent	CaCl ₂ 0,5 mol/l	

5.2 Substrates and standards

5.2.1 Preparation of standard solutions

5.2.1.1 Para-nitrophenol (CAS N°: 100-02-7 - PNP) at 3,6 mmol/l.

- para-nitrophenol (PNP) (Mw:139,11): 10 mg;
- deionized water: 20 ml.

PNP as a powder should be stored at $-20\text{ °C} \pm 2\text{ °C}$ and protected from light. Weigh PNP carefully and dissolve it into deionized water. Working concentration is 0,36 mM (i.e. dilution of the concentrated solution 1/10). The storage of the concentrated and working concentrations shall not exceed two years at $-20\text{ °C} \pm 2\text{ °C}$. Solutions could be aliquoted for one use or maximum 3 freeze/defreeze cycles.

NOTE Paranitrophenol can cause damage to organs through prolonged or repeated exposure if swallowed (H373) and harmful if swallowed, in contact with skin or if inhaled (H302, H312, and H332). Appropriated ventilation and protections need to be used.

5.2.1.2 Ammonium chloride (NH₄Cl) at 62 mmol/l.

- ammonium chloride (CAS N°: 12125-02-9 - Mw:53,49): 66,4 mg;
- deionized water: 20 ml.

Ammonium chloride as a powder can be stored at room temperature and protected from light. Weigh ammonium chloride carefully and dissolve it into water. The concentrated solution should be stored at $-20\text{ °C} \pm 2\text{ °C}$ for two years. The storage of the concentrated solution shall not exceed two years at $-20\text{ °C} \pm 2\text{ °C}$. Working concentration is 0,62 mmol/l (dilution 1/100). The storage of the working concentration shall not exceed two years at $-20\text{ °C} \pm 2\text{ °C}$.

<https://standards.iteh.ai/catalog/standards/sist/917b2d5d-4862-446a-a3d3-118ed5a4ec/iso-20130-2018>

5.2.1.3 β -naphthylamine, 1 mmol/l.

- β -naphthylamine (CAS N°: 91-59-8 - Mw:143,19): 35,8 mg;
- ethanol 96 %: 20 ml;
- deionized water.

Dissolve 35,8 mg of β -naphthylamine into 50 ml ethanol 96 % (0,071 %). This 5 mmol/l stock solution shall be stored at -20 °C for one year. Working concentration is 1 mmol/l (dilution 1/5 in water) and shall be stored at $-20\text{ °C} \pm 2\text{ °C}$ for one year.

NOTE β -Naphthylamine has acute toxicity (oral, dermal, inhalation), category 4, respiratory sensitization, category 1, and hazardous to the aquatic environment. Eyeshields¹⁾, full-face particle respirator type N100 (US)²⁾, appropriated ventilation, specific gloves³⁾, glasses and protective screen need to be used.

5.2.2 Preparation of substrate solutions

Commercially available colorimetric substrates are delivered as powders that should be stored, according to the specifications, frozen at $-20\text{ °C} \pm 2\text{ °C}$ or cooled at $4\text{ °C} \pm 2\text{ °C}$ or stored at room

1) <https://www.sigmaaldrich.com/labware/labware-products.html?TablePage=20009868>

2) <http://www.sigmaaldrich.com/catalog/search?interface=Substance&term=msaadvantageseries1000fullfacerespirator1234598765+OR+msaultratwinfullfacerespirator1234598765&focus=product&mode=boolean>

3) <http://www.sigmaaldrich.com/labware/labware-products.html?TablePage=9577418>

temperature (RT). All the substrates should be prepared in advance according to the requirements of [Table 3](#).

NOTE The volume needs to be sufficient for reliable weighing and measurement of substrates. It depends also on the number of plates needed. Examples are given in [Table 3](#).

Table 3 — Colorimetric substrates available commercially for enzymatic activity measurements and their preparation for measures

Enzyme	Substrate ^a (Storage temperature)	Molar mass g/mol	Concentration mol/l	Examples of preparation	Storage of solutions Temperature, duration
Arylamidase	L-leucine β -naphthylamide hydrochloride (-20 °C \pm 2 °C) CAS N°: 893-36-7	292,8	0,008	Dissolve 0,2342 g in 50 ml of deionized water	-20 °C \pm 2 °C 1 year
Arylsulfatase	Potassium 4-nitrophenyl sulfate (-20 °C \pm 2 °C) CAS N°: 6217-68-1	257,26	0,025	Dissolve 0,322 g in 50 ml of deionized water	-20 °C \pm 2 °C 1 year
β -Galactosidase	p-nitrophenyl β -D-ga- lactopyranoside (-20 °C \pm 2 °C) CAS N°: 3150-24-1	301,25	0,05	Dissolve 0,753 g in 50 ml of deionized water	-20 °C \pm 2 °C 1 year
α -Glucosidase	4-nitrophenyl α -D-glu- copyranoside (-20 °C \pm 2 °C) CAS N°: 3767-28-0	301,25	0,025	Dissolve 0,375 g in 50 ml of deionized water	-20 °C \pm 2 °C 1 year
β -Glucosidase	4-nitrophenyl β -D-glu- copyranoside (-20 °C \pm 2 °C) CAS N°: 2492-87-7	301,25	0,05	Dissolve 0,753 g in 50 ml of deionized water	-20 °C \pm 2 °C 1 year

^a Sigma-Aldrich is an example of producer of colorimetric substrates. This information is given for the convenience of users of this international standard and does not constitute an endorsement by ISO of these products. Equivalent products may be used if they can be shown to lead to the same results.

NOTE All Substrates need to be manipulated with appropriated ventilation, specific gloves, and protective screen.