
**Cosmetics — Microbiology —
Detection of *Staphylococcus aureus***

Cosmétiques — Microbiologie — Détection de Staphylococcus aureus

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[ISO 22718:2015](https://standards.iteh.ai/catalog/standards/sist/0181b09f-a964-47b1-9e3b-fl e167dbb4e8/iso-22718-2015)

<https://standards.iteh.ai/catalog/standards/sist/0181b09f-a964-47b1-9e3b-fl e167dbb4e8/iso-22718-2015>



iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 22718:2015

<https://standards.iteh.ai/catalog/standards/sist/0181b09f-a964-47b1-9e3b-fl e167dbb4e8/iso-22718-2015>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2015, Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Ch. de Blandonnet 8 • CP 401
CH-1214 Vernier, Geneva, Switzerland
Tel. +41 22 749 01 11
Fax +41 22 749 09 47
copyright@iso.org
www.iso.org

Contents

	Page
Foreword.....	iv
Introduction.....	v
1 Scope	1
2 Normative references	1
3 Terms and definitions	1
4 Principle	2
5 Diluents and culture media	2
5.1 General.....	2
5.2 Diluent for the bacterial suspension (tryptone sodium chloride solution).....	3
5.2.1 General.....	3
5.2.2 Composition.....	3
5.2.3 Preparation.....	3
5.3 Culture media.....	3
5.3.1 General.....	3
5.3.2 Agar medium for the suitability test (see Clause 11) [soybean-casein digest agar medium (SCDA) or tryptic soy agar (TSA)].....	3
5.3.3 Enrichment broth.....	4
5.3.4 Selective agar medium for isolation of <i>Staphylococcus aureus</i>	5
6 Apparatus and glassware	6
7 Strains of microorganisms	6
8 Handling of cosmetic products and laboratory samples	6
9 Procedure	7
9.1 General recommendation.....	7
9.2 Preparation of the initial suspension in the enrichment broth.....	7
9.2.1 General.....	7
9.2.2 Water-miscible products.....	7
9.2.3 Water-immiscible products.....	7
9.2.4 Filterable products.....	7
9.3 Incubation of the inoculated enrichment broth.....	7
9.4 Detection and identification of <i>Staphylococcus aureus</i>	7
9.4.1 Isolation.....	7
9.4.2 Identification of <i>Staphylococcus aureus</i>	8
10 Expression of the results (detection of <i>Staphylococcus aureus</i>)	8
11 Neutralization of the antimicrobial properties of the product	9
11.1 General.....	9
11.2 Preparation of inoculum.....	9
11.3 Suitability of the detection method.....	9
11.3.1 Procedure.....	9
11.3.2 Interpretation of suitability test results.....	9
12 Test report	10
Annex A (informative) Other media	11
Annex B (informative) Neutralizers of antimicrobial activity of preservatives and rinsing liquids	14
Bibliography	15

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the WTO principles in the Technical Barriers to Trade (TBT) see the following URL: [Foreword - Supplementary information](#)

The committee responsible for this document is ISO/TC 217, *Cosmetics*.

This second edition cancels and replaces the first edition (ISO 22718:2006), of which it constitutes a minor revision.

<https://standards.iteh.ai/catalog/standards/sist/0181b09f-a964-47b1-9e3b-fl e167dbb4e8/iso-22718-2015>

Introduction

Microbiological examinations of cosmetic products are carried out according to an appropriate microbiological risk analysis in order to ensure their quality and safety for consumers.

Microbiological risk analysis depends on several parameters such as the following:

- potential alteration of cosmetic products;
- pathogenicity of microorganisms;
- site of application of the cosmetic product (hair, skin, eyes, mucous membranes);
- type of users (adults, children under 3 years).

For cosmetics and other topical products, the detection of skin pathogens such as *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Candida albicans* may be relevant because they can cause skin or eye infections. The detection of other kinds of microorganism might be of interest since these microorganisms (including indicators of faecal contamination e.g. *Escherichia coli*) suggest hygienic failure during the manufacturing process.

iTeh STANDARD PREVIEW (standards.iteh.ai)

[ISO 22718:2015](https://standards.iteh.ai/catalog/standards/sist/0181b09f-a964-47b1-9e3b-fl e167dbb4e8/iso-22718-2015)

<https://standards.iteh.ai/catalog/standards/sist/0181b09f-a964-47b1-9e3b-fl e167dbb4e8/iso-22718-2015>

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 22718:2015

<https://standards.iteh.ai/catalog/standards/sist/0181b09f-a964-47b1-9e3b-fl e167dbb4e8/iso-22718-2015>

Cosmetics — Microbiology — Detection of *Staphylococcus aureus*

1 Scope

This International Standard gives general guidelines for the detection and identification of the specified microorganism *Staphylococcus aureus* in cosmetic products. Microorganisms considered as specified in this International Standard might differ from country to country according to national practices or regulations.

In order to ensure product quality and safety for consumers, it is advisable to perform an appropriate microbiological risk analysis to determine the types of cosmetic product to which this International Standard is applicable. Products considered to present a low microbiological (see ISO 29621) risk include those with low water activity, hydro-alcoholic products, extreme pH values, etc.

The method described in this International Standard is based on the detection of *Staphylococcus aureus* in a non-selective liquid medium (enrichment broth), followed by isolation on a selective agar medium. Other methods may be appropriate dependent on the level of detection required.

NOTE For the detection of *Staphylococcus aureus*, subcultures can be performed on non-selective culture media followed by suitable identification steps (e.g. using identification kits).

Because of the large variety of cosmetic products within this field of application, this method may not be appropriate for some products in every detail (e.g. certain water immiscible products). Other International Standards (ISO 18415) may be appropriate. Other methods (e.g. automated) may be substituted for the tests presented here provided that their equivalence has been demonstrated or the method has been otherwise shown to be suitable.

2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 21148:2005, *Cosmetics — Microbiology — General instructions for microbiological examination*

EN 12353, *Chemical disinfectants and antiseptics — Preservation of test organisms used for the determination of bactericidal (including Legionella), mycobactericidal, sporicidal, fungicidal and virucidal (including bacteriophages) activity*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

3.1

product

portion of an identified cosmetic product received in the laboratory for testing

3.2

sample

portion of the product (at least 1 g or 1 ml) that is used in the test to prepare the initial suspension

**3.3
initial suspension**

suspension (or solution) of the sample in a defined volume of an appropriate enrichment broth

**3.4
sample dilution**

dilution of the initial suspension

**3.5
specified microorganism**

aerobic mesophilic bacterium or yeast that is undesirable in a cosmetic product and is recognized as a skin pathogen species that may be harmful for human health or as an indication of hygienic failure in the manufacturing process

**3.6
*Staphylococcus aureus***

gram-positive cocci, mainly aggregated in grape-like clusters, smooth colonies generally pigmented in yellow

Note 1 to entry: The main characteristics for identification are: growth on specific selective medium, catalase positive, coagulase positive.

Note 2 to entry: *Staphylococcus aureus* is an opportunistic pathogen for humans that can also be present on the skin of healthy people without causing disorder for them. It is undesirable in cosmetic products due to its potential pathogenicity.

**3.7
enrichment broth**

non-selective liquid medium containing suitable neutralizers and/or dispersing agents and demonstrated to be suitable for the product under test

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO 22718:2015

<https://standards.iteh.ai/catalog/standards/sist/0181b09f-a964-47b1-9e3b-fl e167dbb4e8/iso-22718-2015>

4 Principle

The first step of the procedure is to perform an enrichment by using a non-selective broth medium to increase the number of microorganisms without the risk of inhibition by the selective ingredients that are present in selective/differential growth media.

The second step of the test (isolation) is performed on a selective medium followed by identification tests.

The possible inhibition of microbial growth by the sample shall be neutralized to allow the detection of viable microorganisms^[1]. In all cases and whatever the methodology, the neutralization of the antimicrobial properties of the product shall be checked and demonstrated (see [Clause 11](#)).

5 Diluents and culture media

5.1 General

General instructions are given in ISO 21148. When water is mentioned in this International Standard, use distilled water or purified water as specified in ISO 21148.

The enrichment broth is used to disperse the sample and to increase the initial microbial population. It may contain neutralizers if the specimen to be tested has antimicrobial properties. The efficacy of the neutralization shall be demonstrated (see [Clause 11](#)). Information relative to suitable neutralizers is given in [Annex B](#).

The enrichment broth ([5.3.3.1](#)), or any of the ones listed in [Annex A](#), is suitable for checking the presence of *Staphylococcus aureus* in accordance with this International Standard provided that it has been demonstrated to be suitable in accordance with [Clause 11](#).

Other diluents and culture media may be used if it has been demonstrated that they are suitable for use.

5.2 Diluent for the bacterial suspension (tryptone sodium chloride solution)

5.2.1 General

The diluent is used for the preparation of bacterial suspension used for the suitability test procedure (see [Clause 11](#)).

5.2.2 Composition

— tryptone, pancreatic digest of casein	1,0 g
— sodium chloride	8,5 g
— water	1 000 ml

5.2.3 Preparation

Dissolve the components in water by mixing while heating. Dispense into suitable containers. Sterilize in the autoclave at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to $7,0 \pm 0,2$ when measured at room temperature.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

5.3 Culture media

5.3.1 General

Culture media may be prepared using the descriptions provided below or from dehydrated culture media according to the instructions from the manufacturer. The instructions provided by the supplier of the media should be followed.

NOTE Ready-to-use media can be used when their composition and/or growth yields are comparable to those of the formulae given herein.

5.3.2 Agar medium for the suitability test (see [Clause 11](#)) [soybean-casein digest agar medium (SCDA) or tryptic soy agar (TSA)]

5.3.2.1 Composition

— pancreatic digest of casein	15,0 g
— papaic digest of soybean meal	5,0 g
— sodium chloride	5,0 g
— agar	15,0 g
— water	1 000 ml

5.3.2.2 Preparation

Dissolve the components or the dehydrated complete medium in the water by mixing while heating. Dispense the medium into suitable containers. Sterilize in the autoclave at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to $7,3 \pm 0,2$ when measured at room temperature.

5.3.3 Enrichment broth

5.3.3.1 Eugon LT 100 broth

5.3.3.1.1 General

This medium contains ingredients which neutralize inhibitory substances present in the sample: lecithin and polysorbate 80, and dispersing agent: octoxynol 9.

5.3.3.1.2 Composition

— pancreatic digest of casein	15,0 g
— papaic digest of soybean meal	5,0 g
— L-cystine	0,7 g
— sodium chloride	4,0 g
— sodium sulfite	0,2 g
— glucose	5,5 g
— egg lecithin	1,0 g
— polysorbate 80	5,0 g
— octoxynol 9	1,0 g
— water	1 000 ml

iTeh STANDARD PREVIEW
(standards.iteh.ai)

<https://standards.iteh.ai/catalog/standards/sist/0181b09f-a964-47b1-9e3b-fl e167dbb4e8/iso-22718-2015>

5.3.3.1.3 Preparation

Dissolve the components, polysorbate 80, octoxynol 9 and egg lecithin successively into boiling water until their complete dissolution. Dissolve the other components by mixing while heating. Dispense the medium into suitable containers. Sterilize in the autoclave at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to $7,0 \pm 0,2$ when measured at room temperature.

5.3.3.2 Other enrichment broths

Other enrichment broths may be used as appropriate (see [Annex A](#)).

5.3.4 Selective agar medium for isolation of *Staphylococcus aureus*

5.3.4.1 Baird Parker agar medium

5.3.4.1.1 Base medium

5.3.4.1.1.1 Composition

— pancreatic digest of casein	10,0 g
— yeast extract	1,0 g
— meat extract	5,0 g
— sodium pyruvate	10,0 g
— L-glycine	12,0 g
— lithium chloride	5,0 g
— agar	12 g to 22 g
— water	to a final volume of 950 ml

5.3.4.1.1.2 Preparation

Dissolve the components or the complete dehydrated base in the water by boiling. Transfer the medium in quantities of 100 ml to flasks or bottles of appropriate capacity. Sterilize the medium in the autoclave at 121 °C for 15 min.

After sterilization and cooling down, the pH shall be equivalent to $7,2 \pm 0,2$ when measured at room temperature.

5.3.4.1.2 Potassium tellurite solution

5.3.4.1.2.1 Composition

— potassium tellurite (K_2TeO_3)	1,0 g
— water	100 ml

5.3.4.1.2.2 Preparation

Dissolve the potassium tellurite completely in the water with minimal heating.

Sterilize by filtration using 0,22 µm pore size membranes. The solution may be stored at the maximum for one month at $3\text{ °C} \pm 2\text{ °C}$. Discard the solution if a white precipitate forms.

The solid should be readily soluble. If a white insoluble material is present in the water, the powder should be discarded.

5.3.4.1.3 Egg yolk emulsion (concentration approximately 20 % or according to the manufacturer's instructions)

If a commercial preparation is not available, prepare the medium as follows.

Use fresh hens' eggs, the shells being intact. Clean the eggs with a brush using a liquid detergent. Rinse them under running water, then disinfect the shells either by immersing them in 70 % (volume fraction) ethanol for 30 s and allow them to dry in the air, or by spraying them with alcohol followed by flame