



SLOVENSKI STANDARD
oSIST prEN 17719:2023
01-maj-2023

Rastlinski biostimulanti - Ugotavljanje števila na anaerobnih mikrotitrskih ploščah

Plant biostimulants - Determination of the anaerobic plate count

Pflanzen-Biostimulanzien - Bestimmung der anaeroben Keimzahl

Biostimulants des végétaux - Détermination du dénombrement sur plaque des germes anaérobies

Ta slovenski standard je istoveten z: prEN 17719

ICS:

65.080 Gnojila Fertilizers

oSIST prEN 17719:2023 **en,fr,de**

EUROPEAN STANDARD
NORME EUROPÉENNE
EUROPÄISCHE NORM

DRAFT
prEN 17719

April 2023

ICS 65.080

Will supersede CEN/TS 17719:2022

English Version

Plant biostimulants - Determination of the anaerobic plate count

Biostimulants des végétaux - Détermination du dénombrement sur plaque des germes anaérobies

Pflanzen-Biostimulanzien - Bestimmung der anaeroben Keimzahl

This draft European Standard is submitted to CEN members for enquiry. It has been drawn up by the Technical Committee CEN/TC 455.

If this draft becomes a European Standard, CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration.

This draft European Standard was established by CEN in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Republic of North Macedonia, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Türkiye and United Kingdom.

Recipients of this draft are invited to submit, with their comments, notification of any relevant patent rights of which they are aware and to provide supporting documentation.

Warning : This document is not a European Standard. It is distributed for review and comments. It is subject to change without notice and shall not be referred to as a European Standard.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Rue de la Science 23, B-1040 Brussels

Contents	Page
European foreword	4
Introduction	5
1 Scope	7
2 Normative references	7
3 Terms and definitions	7
4 Principle	8
4.1 General	8
4.2 Brief description	8
5 Culture media and reagents	8
6 Equipment and consumables	8
7 Preparation of test sample	9
7.1 General	9
7.2 Liquid – water-based formulations	9
7.3 Liquid – oil-based (emulsifiable concentrate - EC) formulations	9
7.4 Solid – wettable powder (WP) formulations	9
7.5 Solid – water dispersible granules (WDG) formulations	9
7.6 Solid – pellets, granules, microgranules (slow release) formulations	10
7.7 Solid substrates	10
8 Procedure	10
8.1 Test portion, initial suspension and dilutions	10
8.2 Inoculation and incubation	10
8.3 Critical control point	10
9 Calculation	10
10 Expression of results	11
11 Performance characteristics of the method	11
11.1 Interlaboratory studies	11
11.2 Sensitivity	11
11.3 Specificity	11
12 Test report	11
13 Quality assurance	11
Annex A (informative) Guidelines for calculating and reporting the Anaerobic Plate Counts (ANPC)	12
A.1 General	12
A.2 Example	12
Annex B (normative) Composition and preparation of culture media and reagents	13
B.1 Anaerobe Agar (ANA)	13

Annex C (informative) Results of the interlaboratory study for the determination of the anaerobic plate count	15
C.1 Results of the interlaboratory study	15
C.2 Statistical analysis.....	15
Annex ZA (informative) Relationship of this European Standard and the essential requirements of Regulation (EU) 2019/1009 making available on the market of EU fertilising products aimed to be covered.....	17
Bibliography	18

iTeh STANDARD PREVIEW (standards.iteh.ai)

[oSIST prEN 17719:2023](https://standards.iteh.ai/catalog/standards/sist/905146b9-702f-4f6e-a2a4-f98567cd7a50/osist-pren-17719-2023)

<https://standards.iteh.ai/catalog/standards/sist/905146b9-702f-4f6e-a2a4-f98567cd7a50/osist-pren-17719-2023>

prEN 17719:2023 (E)

European foreword

This document (prEN 17719:2023) has been prepared by Technical Committee CEN/TC 455 “Plant Biostimulants”, the secretariat of which is held by AFNOR.

This document is currently submitted to the CEN enquiry.

This document will supersede CEN/TS 17719:2022.

This document has been prepared under a Standardization Request given to CEN by the European Commission and the European Free Trade Association, and supports essential requirements of EU Directive(s) / Regulation(s).

For relationship with EU Directive(s) / Regulation(s), see informative Annex ZA, which is an integral part of this document.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[oSIST prEN 17719:2023](https://standards.iteh.ai/catalog/standards/sist/905146b9-702f-4f6e-a2a4-f98567cd7a50/osist-pren-17719-2023)

<https://standards.iteh.ai/catalog/standards/sist/905146b9-702f-4f6e-a2a4-f98567cd7a50/osist-pren-17719-2023>

Introduction

This document was prepared by the experts of CEN/TC 455 “Plant Biostimulants”. The European Committee for Standardization (CEN) was requested by the European Commission (EC) to draft European standards or European standardization deliverables to support the implementation of Regulation (EU) 2019/1009 of 5 June 2019 laying down rules on the making available on the market of EU fertilising products (“FPR” or “Fertilising Products Regulation”). This standardization request, presented as SR M/564 and M/564 Amd1, also contributes to the Communication on “Innovating for Sustainable Growth: A Bio economy for Europe”. Working Group 5 “Labelling and denominations”, was created to develop a work program as part of this standardization request.

Technical committee CEN/TC 455 “Plant Biostimulants” was established to carry out the work program that will prepare a series of standards. The interest in biostimulants has increased significantly in Europe as a valuable tool to use in agriculture. Standardization was identified as having an important role in order to promote the use of biostimulants. The work of CEN/TC 455 seeks to improve the reliability of the supply chain, thereby improving the confidence of farmers, industry, and consumers in biostimulants, and will promote and support commercialisation of the European biostimulant industry.

Biostimulants used in agriculture can be applied in multiple ways: on soil, on plants, as seed treatment, etc. A microbial plant biostimulant consists of a microorganism or a consortium of microorganisms, as referred to in Component Material Category 7 of Annex II of the EU Fertilising Products Regulation 2019/1009 [1].

The method is applicable to microbial plant biostimulants except those composed of aerobic bacterium to verify that the concentration of anaerobes does not exceed the respective limits described in the EU Fertilisers Regulation.

Table 1 summarizes many of the agro-ecological principles and the role played by biostimulants.

Table 1 — Agro-ecological principles and the role played by biostimulants

Increase biodiversity
By improving soil microorganism quality/quantity
Reinforce biological regulation and interactions
By reinforcing plant-microorganism interactions
— symbiotic exchanges i.e. <i>mycorrhize</i>
— symbiotic exchanges i.e. <i>rhizobiaciae/fava</i>
— secretions mimicking plant hormones (i.e. <i>trichoderma</i>)
By regulating plant physiological processes
— e.g. growth, metabolism, plant development...
Improve biogeochemical cycles
— improve absorption of nutritional elements
— improve bioavailability of nutritional elements in the soil
— stimulate degradation of organic matter

prEN 17719:2023 (E)

WARNING — Persons using this document should be familiar with normal laboratory practice. This document does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices and to ensure compliance with any national regulatory conditions.

IMPORTANT — It is absolutely essential that tests conducted in accordance with this document be carried out by suitably trained staff.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[oSIST prEN 17719:2023](https://standards.iteh.ai/catalog/standards/sist/905146b9-702f-4f6e-a2a4-f98567cd7a50/osist-pren-17719-2023)

<https://standards.iteh.ai/catalog/standards/sist/905146b9-702f-4f6e-a2a4-f98567cd7a50/osist-pren-17719-2023>

1 Scope

This document provides a horizontal method for enumeration of microorganisms that are able to grow and form colonies in a solid medium after anaerobic incubation at 30 °C.

The method is applicable to microbial plant biostimulants, except those composed of aerobic bacteria, for verifying that the concentration of anaerobes does not exceed the respective limits outlined in the EU Fertilizers Regulation [1].

This method does not apply to the microbiological monitoring of the environment in which microbial plant biostimulants are manufactured.

No information about potential human pathogens can be inferred from anaerobic plate counts.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN 17708:—¹, *Plant biostimulants — Preparation of sample for microbial analysis*

EN 17724:—², *Plant biostimulants — Terminology*

EN ISO 7218:2007³, *Microbiology of food and animal feeding stuffs — General requirements and guidance for microbiological examinations (ISO 7218:2007)*

EN ISO 11133:2014⁴, *Microbiology of food, animal feed and water — Preparation, production, storage and performance testing of culture media (ISO 11133:2014)*

3 Terms and definitions

For the purposes of this document, the terms and definitions given in EN 17724:—² and the following apply.

ISO and IEC maintain terminology databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <https://www.electropedia.org/>

3.1

microorganism

any microbiological entity, including lower fungi, bacteria and viruses, cellular or non-cellular, capable of replication or of transferring genetic material

[SOURCE: Regulation (EC) No 1107/2009, Article 3, point 15]

¹ Under preparation

² Under preparation

³ As amended by EN ISO 7218:2007/A1:2013

⁴ As amended by EN ISO 11133:2014/A1:2018 and EN ISO 11133:2014/A2:2020

prEN 17719:2023 (E)**3.2****facultative organism**

microorganism capable of both aerobic and anaerobic metabolism

[SOURCE: EN ISO 11139:2018, 3.114]

3.3**obligate anaerobe**

organism that only lives and grows in the absence of molecular oxygen

[SOURCE: EN ISO 11139:2018, 3.186]

3.4**viable count**

value established from enumeration of recoverable colony-forming units

[SOURCE: EN ISO 11139:2018, 3.316]

4 Principle**4.1 General**

Viable anaerobic bacteria are enumerated by the plate count technique under an anaerobic atmosphere [2]. Caution shall be exercised when applying the method since isolates may be pathogenic.

4.2 Brief description

A fixed amount of a dilution of the test sample is placed in an empty Petri dish and mixed in a specified melted agar culture medium to create a pour plate.

The plates are incubated at 30 °C ± 1 °C under anaerobic conditions for 48 h to 72 h.

The number of microorganisms per gram or millilitre of test sample is calculated as specified in Clause 9.

5 Culture media and reagents

For current laboratory practice, EN 17708:—¹ and EN ISO 11133:2014⁴ shall be used.

Composition of Anaerobic Agar (ANA) [3] and reagents and their preparation are described in Annex B.

6 Equipment and consumables

6.1 Special apparatus, anaerobic incubator, BBL GasPak or equivalent, equipped with GasPak hydrogen and CO₂ generator envelopes with an anaerobic indicator.

6.2 Apparatus for dry sterilization (oven) or wet sterilization (autoclave), according to EN ISO 7218:2007³ shall be used.

6.3 Drying cabinet or incubator, capable of operating at 30 °C to 35 °C ± 1 °C.

6.4 Water bath, capable of operating at 47 °C ± 2°C.

6.5 pH-meter, having an accuracy of calibration of ±0,1 pH unit at 25 °C.

6.6 Sterile graduated pipettes or automatic pipettes, of nominal capacities of 0,1 ml, 0,5 ml, 1 ml, and 10 ml.

6.7 Sterile Petri dishes, with a diameter of approximately 90 mm and (optional) large size (diameter approximately 140 mm).

6.8 Refrigerator, capable of operating at $5\text{ °C} \pm 3\text{ °C}$.

6.9 Peristaltic blender (Stomacher^{®5}) with 400 ml sterile bags.

6.10 Blender motor and jars or vortex.

7 Preparation of test sample

7.1 General

To ensure a truly representative analytical unit, agitate liquids or free flowing materials until the contents are homogeneous. If the sample unit is a solid, obtain the analytical unit by taking a portion from several locations within the sample unit. To reduce the workload, the analytical units may be combined for analysis. It is recommended that a composite contains no more than five analytical units.

General rules for the preparation of the initial suspension for microbiological examination are described in EN 17708:—¹.

A representative sample of the product should be prepared taking into consideration the different formulations of the plant biostimulants.

To minimize the effect of the sampling on the significance and the reliability of the analytical results, the number of replicates tested should be increased to 5.

7.2 Liquid – water-based formulations

Aseptically add 25 ml of the product (the analytical unit) in a 400 ml sterile Stomacher[®] bag (6.9) or in a blender jar (6.10) containing 225 ml of an appropriate sterile diluent (i.e. buffered peptone water) maintained at room temperature. Blend, stomach or vortex as required for thorough mixing.

7.3 Liquid – oil-based (emulsifiable concentrate - EC) formulations

Aseptically add 25 ml of the product (the analytical unit) in a 400 ml sterile Stomacher[®] bag (6.9) or in a blender jar (6.10) containing 225 ml of an appropriate sterile diluent (i.e. buffered peptone water) maintained at room temperature. Blend, stomach or vortex as required for thorough mixing.

7.4 Solid – wettable powder (WP) formulations

Aseptically add 25 g of the product (the analytical unit) in a 400 ml sterile Stomacher[®] bag (6.9) containing 225 ml of an appropriate sterile diluent (i.e. buffered peptone water) maintained at room temperature. Homogenize the mixture 2 min at higher speed with a Stomacher[®] (6.9).

7.5 Solid – water dispersible granules (WDG) formulations

Aseptically add 25 g of the product (the analytical unit) in a 400 ml sterile Stomacher[®] bag (6.9) containing 225 ml of an appropriate sterile diluent (i.e. buffered peptone water) maintained at room temperature. Homogenize the mixture 2 min at higher speed with a Stomacher[®] (6.9).

⁵ Stomacher[®] is an example of a suitable product available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement by CEN of this product.