INTERNATIONAL STANDARD

ISO 21317

First edition 2019-02

Traditional Chinese medicine — Lonicera japonica flower

Médecine traditionnelle chinoise — *Fleur de* Lonicera japonica

iTeh Standards (https://standards.iteh.ai) Document Preview

ISO 21317:2019



iTeh Standards (https://standards.iteh.ai) Document Preview

ISO 21317:2019

https://standards.iteh.ai/catalog/standards/iso/cdc85fa5-9673-4dea-87c6-dafa92e3cc5a/iso-21317-2019



COPYRIGHT PROTECTED DOCUMENT

© ISO 2019

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office CP 401 • Ch. de Blandonnet 8 CH-1214 Vernier, Geneva Phone: +41 22 749 01 11 Fax: +41 22 749 09 47 Email: copyright@iso.org Website: www.iso.org

Published in Switzerland

ii

Con	tents	Page
Forev	vord	iv
Intro	duction	v
1	Scope	1
2	Normative references	
	Terms and definitions	
3		
4	Descriptions	
5	Requirements	
	5.1 General characteristics	
	5.2 Macroscopic characteristics	
	5.3 Microscopic characteristics	
	5.4 Moisture	
	5.5 Total ash	
	5.6 Acid-insoluble ash	
	5.7 Dilute ethanol-soluble extract	
	5.8 Identification of chlorogenic acid and luteoloside	
	5.9 Chlorogenic acid content	
		0
6	Sampling iTeh Standards	6
7	Test methods	7
	7.1 Macroscopic identification	
	7.2 Microscopic identification	7
	7.3 Determination of moisture content	7
	7.4 Determination of total ash content	
	7.5 Determination of acid-insoluble ash content	7
	7.6 Determination of dilute ethanol-soluble extract content	7
	7.7 de Identification of chlorogenic acid and luteoloside and	<u></u>
	7.8 Determination of chlorogenic acid content	7
	7.9 Determination of luteoloside content	7
	7.10 Determination of heavy metals content	7
8	Test report	8
9	Packaging, storage and transportation	8
10	Marking and labelling	8
Anne	x A (informative) Identification of chlorogenic acid and luteoloside	9
Anne	x B (informative) Determination of chlorogenic acid content	11
Anne	x C (informative) Determination of luteoloside content	13
Anne	x D (informative) Reference values of national and regional limits of moisture, total ash, acid-insoluble ash, dilute ethanol-soluble extract, chlorogenic acid, luteoloside and heavy metals in <i>Lonicera japonica</i> flower	16
Diblia	ogranhy	18
	IVI ATITIV	124

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 249, *Traditional Chinese medicine*.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

ISO 21317:2019

Introduction

Lonicera japonica flower is the dried flower bud or flower starting to bloom of Lonicera japonica Thunb., which is recorded in the Chinese, South Korean, Japanese and United States Pharmacopeias. Lonicera japonica flower is internationally recognized as a traditional Chinese herbal medicine, and there is great demand for it in the international market. However, there are many problems seriously affecting the international trade of Lonicera japonica flower, including the following.

- 1) Quality requirements for *Lonicera japonica* flower are different among different countries and regions.
- 2) *Lonicera japonica* flower is often substituted with fake and inferior versions.
- 3) Different collecting times, processing methods, packaging, transportation and storage conditions often result in different qualities of *Lonicera japonica* flower.

Therefore, the establishment of an international standard for *Lonicera japonica* flower is necessary to guarantee the quality, safety and consistency of this valuable herbal medicine. This document includes sections on morphology evaluation, physicochemical indexes and heavy metals content.

As national implementation may differ, National Standards Bodies are invited to modify the values given in 5.4, 5.5, 5.6, 5.7, 5.9 and 5.10 in their national standards. Examples of national and regional values are given in Annex D.

iTeh Standards (https://standards.iteh.ai) Document Preview

ISO 21317:2019

iTeh Standards (https://standards.iteh.ai) Document Preview

ISO 21317:2019

Traditional Chinese medicine — Lonicera japonica flower

1 Scope

This document specifies minimum requirements and test methods for *Lonicera japonica* flower, which is derived from the plant *Lonicera japonica* Thunb. It is applicable to *Lonicera japonica* flower that is sold and used as traditional Chinese medicine.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 1575, Tea — Determination of total ash

ISO 1577, Tea — Determination of acid-insoluble ash

ISO 1666, Starch — Determination of moisture content — Oven-drying method

ISO 18664, Traditional Chinese Medicine — Determination of heavy metals in herbal medicines used in Traditional Chinese Medicine

World Health Organization. 2011, Quality control methods for herbal materials, General advice on sampling

CODEX STAN 1: 1985, Codex general standard for the labeling of prepackaged foods

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at https://www.iso.org/obp
- IEC Electropedia: available at http://www.electropedia.org/

3.1

Lonicera japonica flower

dried flower bud or flower starting to bloom of *Lonicera japonica* Thunb.

3.2

chlorogenic acid content

mass fraction of chlorogenic acid in the sample determined in accordance with Annex B

3.3

luteoloside content

mass fraction of luteoloside in the sample determined in accordance with Annex C

3.4

dilute ethanol-soluble extract

mass fraction of extract obtained from the sample using the method specified in 7.6

ISO 21317:2019(E)

3.5 batch


samples collected from the same particular place at the same time

4 Descriptions

In this document, *Lonicera japonica* flower is the dried flower bud or flower starting to bloom of *Lonicera japonica* Thunb. It is externally yellowish-white, greenish-white to yellowish-green, gradually darkening over time to a golden colour (see <u>Figure 1</u>).

iTeh Standards (https://standards.iteh.ai) Document Preview

ISO 21317:2019



Key

- 1 Lonicera japonica flower in different flowering stages
- 2 Lonicera japonica flower selected for medicinal use
- 3 length
- 4 diameter
- 5 calyx

- 6 bract (leaf-like)
- 7 calyx tube
- 8 calyx lobes
- 9 pistil
- 10 stamen

Figure 1 — The structure of Lonicera japonica flower

5 Requirements

5.1 General characteristics

The following requirements shall be met before sampling.

- a) Lonicera japonica flower shall be the dried flower bud or flower starting to bloom.
- b) *Lonicera japonica* flower shall be clean and free from foreign matter except for a few leaf-like bracts.
- c) The presence of living insects, mould and external contaminants which are visible to the naked eye shall not be permitted.

5.2 Macroscopic characteristics

Flower buds are clavate, tapered downwards, slightly curved; 1,5 cm to 3,5 cm long, about 3 mm in diameter in the upper part and 1,5 mm in diameter in the lower part, with a densely pubescent surface; externally yellowish-white or greenish-white, gradually darkening over time to yellowish-brown. The foliaceous bracts are occasionally visible. The calyx is green, pubescent, five-lobed at the apex and about 2 mm long.

5.3 Microscopic characteristics

Lonicera japonica flower pedicels are covered with numerous glandular hairs and non-glandular hairs. The heads of glandular hairs are multicellular, turbinate, subround or slightly oblate, usually 30 μ m to 70 μ m in diameter, exceptionally up to 110 μ m. The stalks of glandular hairs are unicellular or multicellular with up to five cells, usually 20 μ m to 70 μ m long, exceptionally up to 700 μ m. There are two types of non-glandular hairs: (i) thick walls, unicellular, 45 μ m to 900 μ m long, 15 μ m to 40 μ m in diameter, with fine verrucae on the surface, some have a corneous spiral; (ii) thin walls, slender, curved or shrinkage, with fine verrucae on the surface (see Figure 2).

ISO 21317:2019