



SLOVENSKI STANDARD
oSIST prEN ISO 11267:2022
01-september-2022

Kakovost tal - Zaviranje razmnoževanja vrste Folsomia candida iz rodu skakačev (Collembola) zaradi onesnaževal v tleh (ISO/DIS 11267:2022)

Soil quality - Inhibition of reproduction of Collembola (Folsomia candida) by soil contaminants (ISO/DIS 11267:2022)

Bodenbeschaffenheit - Hemmung der Reproduktion von Collembolen (Folsomia candida) durch Verunreinigungen (ISO/DIS 11267:2022)

Qualité du sol - Inhibition de la reproduction de Collembola (Folsomia candida) par des contaminants du sol (ISO/DIS 11267:2022)

Ta slovenski standard je istoveten z: prEN ISO 11267

ICS:

13.080.30 Biološke lastnosti tal Biological properties of soils

oSIST prEN ISO 11267:2022 **en,fr,de**

DRAFT INTERNATIONAL STANDARD

ISO/DIS 11267

ISO/TC 190/SC 4

Secretariat: AFNOR

Voting begins on:
2022-08-11Voting terminates on:
2022-11-03

Soil quality — Inhibition of reproduction of *Collembola* (*Folsomia candida*) by soil contaminants

Qualité du sol — Inhibition de la reproduction de Collembola (Folsomia candida) par des contaminants du sol

ICS: 13.080.30

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[oSIST prEN ISO 11267:2022](https://standards.iteh.ai/catalog/standards/sist/516137d2-d825-48ed-ac02-fec39be24aa6/osist-pren-iso-11267-2022)

<https://standards.iteh.ai/catalog/standards/sist/516137d2-d825-48ed-ac02-fec39be24aa6/osist-pren-iso-11267-2022>

This document is circulated as received from the committee secretariat.

THIS DOCUMENT IS A DRAFT CIRCULATED FOR COMMENT AND APPROVAL. IT IS THEREFORE SUBJECT TO CHANGE AND MAY NOT BE REFERRED TO AS AN INTERNATIONAL STANDARD UNTIL PUBLISHED AS SUCH.

IN ADDITION TO THEIR EVALUATION AS BEING ACCEPTABLE FOR INDUSTRIAL, TECHNOLOGICAL, COMMERCIAL AND USER PURPOSES, DRAFT INTERNATIONAL STANDARDS MAY ON OCCASION HAVE TO BE CONSIDERED IN THE LIGHT OF THEIR POTENTIAL TO BECOME STANDARDS TO WHICH REFERENCE MAY BE MADE IN NATIONAL REGULATIONS.

RECIPIENTS OF THIS DRAFT ARE INVITED TO SUBMIT, WITH THEIR COMMENTS, NOTIFICATION OF ANY RELEVANT PATENT RIGHTS OF WHICH THEY ARE AWARE AND TO PROVIDE SUPPORTING DOCUMENTATION.

ISO/CEN PARALLEL PROCESSING



Reference number
ISO/DIS 11267:2022(E)

© ISO 2022

iTeh STANDARD PREVIEW (standards.iteh.ai)

[oSIST prEN ISO 11267:2022](https://standards.iteh.ai/catalog/standards/sist/516137d2-d825-48ed-ac02-fec39be24aa6/osist-pren-iso-11267-2022)

<https://standards.iteh.ai/catalog/standards/sist/516137d2-d825-48ed-ac02-fec39be24aa6/osist-pren-iso-11267-2022>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2022

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

Contents		Page
Foreword		iv
Introduction		v
1 Scope		1
2 Normative references		1
3 Terms and definitions		1
4 Principle		3
5 Reagents and material		4
6 Apparatus		6
7 Procedure		6
7.1 Experimental design.....		6
7.1.1 General.....		6
7.1.2 Range-finding test.....		6
7.1.3 Definitive test.....		7
7.2 Preparation of test mixture.....		7
7.2.1 Testing contaminated soil.....		7
7.2.2 Testing substances added to the test substrate.....		8
7.2.3 Preparation of control container.....		9
7.3 Addition of the biological material.....		9
7.4 Test conditions and measurements.....		9
7.5 Determination of surviving Collembola.....		9
8 Calculation and expression of results		9
8.1 Calculation.....		9
8.2 Expression of results.....		10
9 Validity of the test		10
10 Statistical analysis		10
10.1 General.....		10
10.2 Single-concentration tests.....		10
10.3 Multi-concentration tests.....		11
10.3.1 Range-finding test.....		11
10.3.2 Definitive test.....		11
11 Test report		12
Annex A (informative) Techniques for rearing and breeding of Collembola		13
Annex B (informative) Determination of water-holding capacity		15
Annex C (informative) Guidance on adjustment of pH of artificial soil		16
Annex D (informative) Extraction and counting of Collembola		17
Annex E (informative) Specific information of alternative Collembolan species		18
Bibliography		30

ISO/DIS 11267:2022(E)

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the WTO principles in the Technical Barriers to Trade (TBT) see the following URL: [Foreword - Supplementary information](#).

This document was prepared by Technical Committee ISO/TC 190, *Soil quality*, Subcommittee SC 4, *Biological characterization*.

This third edition cancels and replaces the second edition (ISO 11267:2014), which has been technically revised. The main change to the previous edition is as follows:

- Addition of an annex of specific information when using alternative *Collembola* species for reproduction test.

Introduction

Ecotoxicological test systems are applied to obtain information about the effects of contaminants in soil and are proposed to complement conventional chemical analysis (see [2] and [4]). Reference [2] includes a list and short characterization of recommended and standardized test systems and [4] gives guidance on the choice and evaluation of the bioassays. Aquatic test systems with soil eluate are applied to obtain information about the fraction of contaminants potentially reaching the groundwater by the water path (retention function of soils), whereas terrestrial test systems are used to assess the habitat function of soils.

Soil-dwelling Collembola are ecologically relevant species for ecotoxicological testing. Springtails are prey animals for a variety of endogeic and epigeic invertebrates and they contribute to decomposition processes in soils. In acidic soils they may be the most important soil invertebrates besides enchytraeids with respect to that function, since earthworms are typically absent^[19]. Additionally, Collembola represent arthropod species with a different route and a different rate of exposure compared to earthworms^[1] and enchytraeids^[3]. Various species were used in bioassays of which four species were used most commonly, *Folsomia candida* Willem, *Folsomia fimetaria* L., *Onychiurus armatus*, and *Orchesella cincta*^[20]. Numerous soil toxicity tests supported by Environment Canada (EC) resulted in the development and standardization of a biological test method for determining the lethal and sublethal toxicity of samples of contaminated soil to Collembola^[10]. The method prepared by EC includes four species, *Orthonychiurus folsomi*, *Proisotoma minuta*, *F. candida*, and *F. fimetaria*. As standardized test systems using Collembola as indicator organisms for the habitat function of soil, another two methods exist. One is designed for assessing the effects of substances on the reproductive output of the Collembola, *F. fimetaria* and *F. candida* in soil,^{[19],[21]} and the other method described here, focuses on testing contaminated soil. Optionally the method can be used for testing substances added to standard soils (e.g. artificial soil) for their sublethal hazard potential to Collembola.

This International Standard describes a method that is based on the determination of sublethal effects of contaminated soils to adult Collembola of the species *Folsomia candida* Willem. The species is distributed worldwide. It plays a similar ecological role to *F. fimetaria* ^{[10],[19]} *F. candida* reproduces parthenogenetically and is an easily accessible species as it is commercially available and easy to culture. *F. candida* is considered to be a representative of soil arthropods and Collembola in particular. Background information on the ecology of springtails and their use in ecotoxicological testing is available.^[22]

Distinct Collembolan species inhabit various ecological niches at different soil depths and in different soil types across the globe. Although considered a surrogate species and therefore frequently used in ecotoxicological reproduction tests, *F. candida* is not common in most natural soils (OECD 2016). Furthermore, species specific morphological adaptations can influence exposure and toxic effects of chemicals on organisms (Chelinho et al., 2014). Thus, the use of a variety of Collembolan species representing different morphological adaptations may be advantageous to obtain a broad spectrum of sensitivities for this group. Therefore, other species like *F. fimetaria* (euedaphic, distributed worldwide and found in agricultural soils (OECD 2016)), *Onychiurus yodai* (an euedaphic Asian species (Janssens 2020)), *Proisotoma minuta* (hemiedaphic, distributed worldwide and inhabiting agricultural soils (Buch et al. 2016, Janssens 2020)), *Protaphorura fimata* (euedaphic, occurring through mild temperate to cold zones (Holmstrup & Slotsbo, 2018; Janssens, 2020)), and *Sinella curviseta* (epedaphic, distributed from North America to Europe, Southeast Asia and Japan (Salmon 1964)) were added as potential alternative test species ([Annex E](#)). These species have been used as ecotoxicological test species before, but available testing experience might be limited.

Soil quality — Inhibition of reproduction of *Collembola* (*Folsomia candida*) by soil contaminants

1 Scope

This International Standard specifies one of the methods for evaluating the habitat function of soils and determining effects of soil contaminants and substances on the reproduction of *Folsomia candida* Willem by dermal and alimentary uptake. This chronic test is applicable to soils and soil materials of unknown quality, e.g. from contaminated sites, amended soils, soils after remediation, industrial, agricultural or other sites of concern and waste materials.

Effects of substances are assessed using a standard soil, preferably a defined artificial soil substrate. For contaminated soils, the effects are determined in the soil to be tested and in a control soil. According to the objective of the study, the control and dilution substrate (dilution series of contaminated soil) are either an uncontaminated soil comparable to the soil to be tested (reference soil) or a standard soil (e.g. artificial soil).

This International Standard provides information on how to use this method for testing substances under temperate conditions.

The method is not applicable to volatile substances, i.e. substances for which H (Henry's constant) or the air/water partition coefficient is greater than 1, or for which the vapour pressure exceeds 0.013 3 Pa at 25 °C.

NOTE The stability of the test substance cannot be ensured over the test period. No provision is made in the test method for monitoring the persistence of the substance under test.

2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 10694, *Soil quality — Determination of organic and total carbon after dry combustion (elementary analysis)*

ISO 10390, *Soil, treated biowaste and sludge – Determination of pH*

ISO 11260, *Soil quality — Determination of effective cation exchange capacity and base saturation level using barium chloride solution*

ISO 11277, *Soil quality — Determination of particle size distribution in mineral soil material — Method by sieving and sedimentation*

ISO 11465, *Soil quality — Determination of dry matter and water content on a mass basis — Gravimetric method*

ISO 18400-206, *Soil quality — Sampling — Part 206: Collection, handling and storage of soil under aerobic conditions for the assessment of microbiological processes, biomass and diversity in the laboratory*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO/DIS 11267:2022(E)

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <https://www.electropedia.org/>

3.1

contaminant

substance or agent present in the soil as a result of human activity

3.2

EC_x

effect concentration for x % effect

concentration (mass fraction) of a test sample or a test substance that causes x % of an effect on a given end-point within a given exposure period when compared with a control

EXAMPLE An EC₅₀ is a concentration estimated to cause an effect on a test end-point in 50 % of an exposed population over a defined exposure period.

Note 1 to entry: The EC_x is expressed as a percentage of soil to be tested (dry mass) per soil mixture (dry mass). When substances are tested, the EC_x is expressed as mass of the test substance per dry mass of soil in milligrams per kilogram.

3.3

ER_x

effect rate for x % effect

rate of a contaminated soil that causes x % of an effect on a given end-point within a given exposure period when compared with a control

3.4

limit test

single concentration test consisting of at least four replicates each, the soil to be tested without any dilution or the highest concentration of test substance mixed into the control soil and the control

3.5

LOEC

lowest observed effect concentration

lowest test substance concentration that has a statistically significant effect ($p < 0,05$) when compared with the control

Note 1 to entry: In this test, the LOEC is expressed as a mass of test substance per dry mass of the soil to be tested. All test concentrations above the LOEC should usually show an effect that is statistically different from the control.

3.6

LOER

lowest observed effect rate

lowest rate of a contaminated soil tested in a control soil that has a statistically significant effect ($p < 0,05$) when compared with the control

3.7

NOEC

no observed effect concentration

highest test substance concentration immediately below the LOEC at which no statistically significant effect is observed when compared to the control

Note 1 to entry: In this test, the concentration corresponding to the NOEC has no statistically significant effect ($p < 0,05$) within a given exposure period when compared with the control.

3.8**NOER****no observed effect rate**

highest rate of a contaminated soil to be tested immediately below the LOER at which no statistically significant effect is observed when compared to the control

3.9**reference soil**

uncontaminated soil with comparable pedological properties (nutrient concentrations, pH, organic carbon content and texture) to the soil being studied

3.10**standard soil**

field-collected soil or artificial soil whose main properties (pH, texture, organic matter content) are within a known range

EXAMPLE Euro soils, artificial soil, LUFA standard soil.

Note 1 to entry: The properties of standard soils can differ from the soil to be tested.

3.11**control soil**

reference or standard soil used as a control and as a medium for preparing dilution series with soils to be tested or a reference substance, which fulfils the validity criteria

Note 1 to entry: In the case of natural soil, it is advisable to demonstrate its suitability for a test and for achieving the test validity criteria before using the soil in a definitive test.

3.12**test mixture**

mixture of contaminated soil or the test substance (e.g. chemical, biosolid, waste) with control soil

3.13**test mixture ratio**

ratio between the soil to be tested and the control soil in a test mixture

4 Principle

The effects on reproductive output of 10 d to 12 d old *Collembola* (*F. candida*) exposed to the soil to be tested are compared to those observed in a control soil. If appropriate, effects based on exposure to a test mixture of contaminated soil and control soil or a range of concentrations of a test substance mixed into control soil are determined. Test mixtures are prepared at the start of the test and are not renewed within the test period.

The *Collembola* are incubated until offspring (F_1) emerge from eggs laid by mature adults, and the number of offspring is determined. Usually offspring emerge within 28 d in control experiments. The results obtained from the tests are compared with a control or, if appropriate, are used to determine the concentrations which cause no effects on mortality and reproductive output (NOER/NOEC) and the concentration resulting in x % reduction of juveniles hatched from eggs compared to the control (ER_x/EC_x, 28 d) respectively.

If testing a concentration series, all test dilutions/concentrations above the LOER/LOEC have a harmful effect equal to or greater than that observed at the LOER/LOEC. Where there is no prior knowledge of the concentration of the soil to be tested or the test substance likely to have an effect, then it is useful to conduct the test in two steps:

- An acute toxicity test (range-finding test) is carried out, to give an indication of the effect dilution/concentration, and the dilution/concentration giving no mortality (NOER/NOEC). Dilutions/concentrations to be used in the definitive test can then be selected;

ISO/DIS 11267:2022(E)

- The definitive test on reproductive output to determine sublethal effects of (dilutions of) contaminated soil or the concentration of a substance which, when evenly mixed into the standard soil, causes no significant effects on numbers of offspring hatched from eggs compared with the control (NOER/NOEC), and the lowest concentration causing effects (LOER/LOEC).

NOTE The use of a reference soil is an essential requirement to demonstrate the present status of the test population, and to avoid misinterpretation of results.

5 Reagents and material

5.1 Biological material, in this test, 10 d to 12 d old juvenile springtails of the species *Folsomia candida* Willem are used (see [A.1](#) for details on synchronization of breeding).

5.2 Test mixture, which may consist of field-collected soil, waste material or control soil amended by the test substance.

5.2.1 Field-collected soil or waste

The sample(s) can be field-collected soil from an industrial, agricultural or other site of concern, or waste materials (e.g. dredged material, municipal sludge from a wastewater treatment plant, composted material, or manure) under consideration for possible land disposal.

The field-collected soils used in the test shall be passed through a sieve of 4 mm square mesh to remove coarse fragments and thoroughly mixed. If necessary, soil may be air-dried without heating before sieving. Storage of soil to be tested should be as short as possible. The soil shall be stored in accordance with ISO 18400-206 using containers that minimize losses of soil contaminants by volatilization and sorption to the container walls. If soils or test mixtures have been stored, they should be mixed a second time immediately before use. Soil pH should not be corrected as it can influence bioavailability of soil contaminants.

For interpretation of test results, the following characteristics shall be determined for each soil sampled from a field site:

- pH in accordance with ISO 10390,
- texture (sand, loam, silt) in accordance with ISO 11277,
- water content in accordance with ISO 11465,
- water-holding capacity according to [Annex B](#),
- cationic exchange capacity in accordance with ISO 11260,
- organic carbon in accordance with ISO 10694,
- percentage of material (mineral and organic) removed by the 4 mm sieve.

NOTE It is important to measure the water holding capacity of all mixtures used in the test.

5.2.2 Control soil, either a) reference soil ([3.9](#)) or b) standard soil ([3.10](#)) that allows the presence of Collembola. Control soil and soil used for dilution shall not differ in one test (either a) or b)).

- If reference soils from uncontaminated areas near a contaminated site are available, they should be treated and characterized like the soils to be tested. If a toxic contamination or unusual soil properties cannot be ruled out, standard control soils should be preferred.
- For testing the effects of substances mixed into soil, standard soils (e.g. artificial soil, LUFA) shall be used as test substrate. The properties of the field-collected standard soil shall be reported.

The substrate called artificial soil can be used as a standard soil and has the following composition:

Percentage expressed on dry mass basis

- Sphagnum peat finely ground [a particle size of (2 ± 1) mm is acceptable] and with no visible plant remains 10 %
- Kaolinite clay containing not less than 30 % kaolinite 20 %
- Industrial quartz sand (dominant fine sand with more than 50 % of particle size 0,05 mm to 0,2 mm) 69 %

Approximately 0,3 % to 1,0 % calcium carbonate (CaCO_3 , pulverized, analytical grade) are necessary to get a pH of $6,0 \pm 0,5$.

NOTE 1 Taking the properties of highly non-polar ($\log K_{ow} > 2$) or ionizing substances into account, 5 % of peat have proven to be sufficient for maintaining the desired structure of the artificial soil.

NOTE 2 It has been demonstrated that *F. candida* can comply with the validity criteria even on reproductive output when tested in field soils with lower organic carbon content (e.g. 2,7 %), and there is experience that this can be achieved in artificial soil with 5 % peat. Therefore, it is not necessary, before using such a soil in a definitive test, to demonstrate the suitability of the artificial soil for allowing the test to comply with the validity criteria unless the peat contents is lower than specified above.

Prepare the artificial soil at least three days prior to the start of the test, by mixing the dry constituents listed above thoroughly in a large-scale laboratory mixer. A portion of the deionized water required is added during mixing. Allowance should be made for any water that is used for introducing the test substance into the soil. The amount of calcium carbonate required can vary, depending on properties of the individual batch of sphagnum peat and should be determined by measuring sub-samples immediately before the test. Store the mixed artificial soil at room temperature for at least two days to equilibrate acidity. To determine pH and the maximum water holding capacity, the dry artificial soil is pre-moistened one or two days before starting the test by adding deionized water to obtain approximately half of the required final water content of 40 % to 60 % of the maximum water holding capacity. <https://standards.iteh.ai/catalog/standards/sist/516137d2-d825-48ed-ac02-fec39be24aa6/osist-pr-en-iso-11267-2022>

The total water-holding capacity is determined according to [Annex B](#), the pH is determined according to ISO 10390.

5.3 Food

A sufficient amount, e.g. 2 mg to 10 mg, of granulated dried baker's yeast, commercially available for household use, is added to each container as a suitable food source, at the beginning of the test and after about two weeks.

5.4 Reference substance

To ensure the quality of the test system, tests should be performed regularly (once or twice a year) with a reference substance.

Boric acid and the plant protection product Betosip¹⁾ (a.i. 157 g/l phenmedipham) have been tested in a ring test, and are recommended as reference substances.

WARNING — When handling these substances, appropriate precautions should be taken to avoid ingestion or skin contact.

1) Betosip is an example of a suitable product available commercially. This information is given for the convenience of users of this document and does not constitute an endorsement by ISO of the product named. Equivalent products may be used if they can be shown to lead to the same results.