



SLOVENSKI STANDARD
kSIST-TS FprCEN/TS 17709:2021
01-november-2021

[Not translated]

Plant biostimulants - Determination of Azotobacter spp.

Biostimulanzien für die pflanzliche Anwendung - Bestimmung von Azotobacter spp.

Biostimulants des végétaux - Détermination d'Azotobacter spp.

Ta slovenski standard je istoveten z: FprCEN/TS 17709

[kSIST-TS FprCEN/TS 17709:2021](https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-2b563054ce83/ksist-ts-fprcen-ts-17709-2021)

<https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-2b563054ce83/ksist-ts-fprcen-ts-17709-2021>

ICS:

65.080 Gnojila Fertilizers

kSIST-TS FprCEN/TS 17709:2021 en,fr,de

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[kSIST-TS FprCEN/TS 17709:2021](https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-2b563054ce83/ksist-ts-fprcen-ts-17709-2021)

<https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-2b563054ce83/ksist-ts-fprcen-ts-17709-2021>

TECHNICAL SPECIFICATION
SPÉCIFICATION TECHNIQUE
TECHNISCHE SPEZIFIKATION

FINAL DRAFT
FprCEN/TS 17709

September 2021

ICS 65.080

English Version

Plant biostimulants - Determination of *Azotobacter* spp.

Biostimulanzien für die pflanzliche Anwendung -
Bestimmung von *Azotobacter* spp.

This draft Technical Specification is submitted to CEN members for Vote. It has been drawn up by the Technical Committee CEN/TC 455.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Republic of North Macedonia, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.

Recipients of this draft are invited to submit, with their comments, notification of any relevant patent rights of which they are aware and to provide supporting documentation.

Warning : This document is not a Technical Specification. It is distributed for review and comments. It is subject to change without notice and shall not be referred to as a Technical Specification.

[kSIST-TS FprCEN/TS 17709:2021](https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-2b563054ce83/ksist-ts-fprcen-ts-17709-2021)

<https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-2b563054ce83/ksist-ts-fprcen-ts-17709-2021>



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Rue de la Science 23, B-1040 Brussels

Contents	Page
European foreword.....	3
Introduction	4
1 Scope	6
2 Normative references	6
3 Terms and definitions	6
4 Enumeration of <i>Azotobacter spp</i>	6
4.1 General	6
4.2 Sample preparation.....	6
4.2.1 General	6
4.2.2 Liquid (based water) formulations.....	6
4.2.3 Liquid - based oil, emulsifiable concentrate (EC) formulations.....	6
4.2.4 Solid - Wettable Powder (WP) formulations.....	7
4.2.5 Solid - Water dispersible granules (WDG) formulations	7
4.2.6 Solid - Pellets, granules, microgranules - slow release - formulations	7
4.2.7 Solid - substrate	7
4.3 Serial dilution	7
4.4 Plate counts of <i>Azotobacter sp</i> in sterile diluent.....	8
4.5 Spread-plate counting with ASHBY SUCROSE AGAR [6]	8
4.6 Calculation	9
5 Species Determination of <i>Azotobacter sp.</i> via genetic analysis	9
5.1 Preparation of the sample for the genomic DNA extraction.....	9
5.1.1 Isolation and preparation of the microorganism	9
5.1.2 Sample concentration	9
5.1.3 DNA extraction and storage.....	9
5.1.4 Partial PCR Amplification of the 16S rRNA Genes	10
Annex A (informative) Formula of culture media.....	11
A.1 Ashby's sucrose Agar medium.....	11
A.1.1 Ingredients	11
A.1.2 Method to prepare YT media	11
A.2 Procedure for preparing nutrient broth medium	11
A.3 Procedure for preparing 0,1 M Phosphate Buffer Saline (PBS).....	12
Bibliography.....	13

European foreword

This document (FprCEN/TS 17709:2021) has been prepared by Technical Committee CEN/TC 455 “Plant biostimulants”, the secretariat of which is held by AFNOR.

This document is currently submitted to the Vote on TS.

This document has been prepared under a mandate given to CEN by the European Commission and the European Free Trade Association.

iTeh STANDARD PREVIEW (standards.iteh.ai)

[kSIST-TS FprCEN/TS 17709:2021](https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-2b563054ce83/ksist-ts-fprcen-ts-17709-2021)

<https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-2b563054ce83/ksist-ts-fprcen-ts-17709-2021>

Introduction

This document was prepared by the experts of CEN/TC 455 ‘Plant Biostimulants’. The European Committee for Standardization (CEN) was requested by the European Commission (EC) to draft European standards or European standardization deliverables to support the implementation of Regulation (EU) 2019/1009 of 5 June 2019 laying down rules on the making available on the market of EU fertilising products (“FPR” or “Fertilising Products Regulation”). This request, presented as SR M/564, also contributes to the Communication on “Innovating for Sustainable Growth: A Bio economy for Europe”. The Working Group 5 “Labelling and denominations”, was created to develop a work program as part of this Request. The technical committee CEN/TC 455 ‘Plant Biostimulants’ was established to carry out the work program that will prepare a series of standards. The interest in biostimulants has increased significantly in Europe as a valuable tool to use in agriculture. Standardization was identified as having an important role in order to promote the use of biostimulants. The work of CEN/TC 455 seeks to improve the reliability of the supply chain, thereby improving the confidence of farmers, industry, and consumers in biostimulants, and will promote and support commercialisation of the European biostimulant industry.

Biostimulants used in agriculture can be applied in multiple ways: on soil, on plant, as seed treatment, etc. A microbial plant biostimulant consists of a microorganism or a consortium of microorganisms, as referred to in Component Material Category 7 of Annex II of the EU Fertilizing Products Regulation.

This document is applicable to all biostimulants in agriculture based on live microorganisms belonging to the genera *Azotobacter*.

The Table 1 below summarizes many of the agro-ecological principles and the role played by biostimulants.

Table 1 — Agro-ecological principles and the role played by biostimulants

Increase biodiversity
By improving soil microorganism quality/quantity
Reinforce biological regulation and interactions
By reinforcing plant-microorganism interactions
- symbiotic exchanges i.e. <i>mycorrhize</i>
- symbiotic exchanges i.e. <i>rhizobiaciae/fava</i>
- secretions mimicking plant hormones (i.e. <i>trichoderma</i>)
By regulating plant physiological processes
- for ex growth, metabolism, plant development...
Improve biogeochemical cycles
- improve absorption of nutritional elements
- improve bioavailability of nutritional elements in the soil
- stimulate degradation of organic matter

WARNING — Persons using this document should be familiar with normal laboratory practice. This document does not purport to address all of the safety problems, if any, associated with its use. It is the responsibility of the user to establish appropriate safety and health practices and to ensure compliance with any national regulatory conditions.

IMPORTANT — It is absolutely essential that tests conducted in accordance with this document be carried out by suitably trained staff.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[kSIST-TS FprCEN/TS 17709:2021
https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-2b563054ce83/ksist-ts-fprcen-ts-17709-2021](https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-2b563054ce83/ksist-ts-fprcen-ts-17709-2021)

FprCEN/TS 17709:2021 (E)**1 Scope**

This document was developed to provide the methodology for the enumeration and determination of *Azotobacter sp.* in plant biostimulant products in accordance with the Regulation (EU) 2019/1009 of the European Parliament and of the Council [1].

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

FprCEN/TS 17702-1, *Plant biostimulants — Sampling and sample preparation — Part 1: Sampling*

FprCEN/TS 17724, *Plant biostimulants — Terminology*

3 Terms and definitions

For the purposes of this document, the terms and definitions given in FprCEN/TS 17724 apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <https://www.electropedia.org/>
- ISO Online browsing platform: available at <https://www.iso.org/obp>

4 Enumeration of *Azotobacter spp***4.1 General**

[kSIST-TS FprCEN/TS 17709:2021
https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-2b563054ce83/ksist-ts-fprcen-ts-17709-2021](https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-2b563054ce83/ksist-ts-fprcen-ts-17709-2021)

This procedure is meant to determine the number of colony-forming units (CFU) of the above mentioned bacteria«Pathogen», per gram, per millilitre, per square centimetre, or per sampling device. The method, in order to be fast, cheap, repeatable, is based on serial dilutions and plating.

4.2 Sample preparation**4.2.1 General**

A representative sample of the product to be analysed according to FprCEN/TS 17702-1 shall be prepared according to following procedure which takes into consideration the different formulations of biostimulants based products.

4.2.2 Liquid (based water) formulations

Dispense 25 ml of sample (or more for low concentrated products) in 225 ml of sterile Phosphate Buffer Solution (PBS) maintained at room temperature, in a flask and shake for 10 min or more until the distribution is optimal, with a magnetic stirrer at half speed [6].

4.2.3 Liquid - based oil, emulsifiable concentrate (EC) formulations

Dispense 25 ml of sample (or more for low concentrated products) in 225 ml of sterile Phosphate Buffer Solution (PBS) maintained at room temperature, in a flask and shake for 10 min or more until the distribution is optimal, with a magnetic stirrer at half speed [6].

4.2.4 Solid - Wettable Powder (WP) formulations

Dispense 25 g of sample (or more for low concentrated products) in 225 ml of sterile Phosphate Buffer Solution (PBS) maintained at room temperature, in a flask and shake for 20 min or more until the distribution is optimal, with a magnetic stirrer at half speed [6].

4.2.5 Solid - Water dispersible granules (WDG) formulations

Dispense 25 g (or more for low concentrated products) of sample in 225 g of sterile Phosphate Buffer Solution (PBS) maintained at room temperature, in a flask and shake for 40 min or more until the distribution is optimal, with a magnetic stirrer at half speed. If required help the dispersion of the formulations with other apparatus such as a stomacher after having sieved (100 mesh sieve) the particles and resuspend them in the same suspension [6].

4.2.6 Solid - Pellets, granules, microgranules - slow release - formulations

Dispense 25 g (or more for low concentrated products) g of sample in 225 g of sterile Phosphate Buffer Solution (PBS) maintained at room temperature, in a sterile bag and disperse them using a magnetic stirrer for 40 min at half speed and then sieve in a 100 mesh sieve and if material remain in the sieve repeat the process for a maximum of three times. Put attention to all the buffer used to make the exact final calculation [6].

4.2.7 Solid - substrate


Dispense 25 g (or more for low concentrated products) g of sample in 225 g of sterile Phosphate Buffer Solution (PBS) maintained at room temperature, in a flask and shake for 20 min or more until the distribution is optimal, with a magnetic stirrer at half speed [6].

4.3 Serial dilution

The principle in counting bacteria by dilution is to serially dilute them to reduce the bacterial density to the level where individual cells can be differentiated. This may be, for example, as live cells under the microscope, as colonies that grow on plates from single cells, or estimated in the plant-infection technique (with the principle that a single cell can multiply to initiate an infection). Serial dilution can be applied to all kind of formulations. A 10-fold serial dilution is most often used (Figure 1) but if the number of *Azotobacter sp.* cells is expected to be low then a lower number of dilutions can be adopted.

A sample of the product is shaken in a bulk diluent (PBS) which represents the first level of dilution. This is then serially diluted with a sample at each level of dilution directly plated.

FprCEN/TS 17709:2021 (E)

**Key**

- A product suspended in sterile diluent
 B prepare dilution series at 10^{-8}
 C 3 replicates of Petri dishes with media

Figure 1 — Scheme of a serial dilution series

kSIST-TS FprCEN/TS 17709:2021

<https://standards.iteh.ai/catalog/standards/sist/cafa9812-d69e-4a6a-bb5e-26561054cc65/kSIST-TS-FprCEN-TS-17709-2021>

4.4 Plate counts of *Azotobacter sp* in sterile diluent

The counting of microorganisms on plates, following dilution, is also called direct counting. Counts only plates where there are between 30 to 300 colonies.

4.5 Spread-plate counting with ASHBY SUCROSE AGAR [6]

The steps of the procedure are the following:

1. inoculate 0,1 ml of the serial dilutions desired (e.g. 10^{-5} , 10^{-6} and 10^{-7}) on the surface of the culture medium in Petri dishes (Figure 1);
2. spread the 0,1 ml aliquot over the culture medium with a sterilized L-shaped glass spreader (or equivalent, e.g. a Drigalski loop);
3. there should be at least three separate replicate plates for each dilution;
4. after inoculation and absorption of the inoculum into the agar, the plates are placed in an incubator at approximately 30 °C, inverted and allowed to grow for a period of 4 days;
5. count the number of colonies on plates where colonies are well separated. If colony numbers are low, variation between plates and errors may be large. If colony numbers are too high, overcrowding may result in an underestimation of numbers. Many texts recommend counting between 30 CFU (Colony Forming Units) and 300 CFU per plate to give statistical robustness.