

SLOVENSKI STANDARD oSIST prEN ISO 18314-4:2021

01-marec-2021

Analizna kolorometrija - 4. del: Metamerični indeks parov vzorcev pri spremembi vrste svetila (ISO 18314-4:2020)

Analytical colorimetry - Part 4: Metamerism index for pairs of samples for change of illuminant (ISO 18314-4:2020)

Analytische Farbmessung - Teil 4: Metamerie-Index von Probenpaaren bei Lichtartwechsel (ISO 18314-4:2020) NDARD PREVIEW

Analyse colorimétrique - Partie 4. Indice de metamérisme de paires d'échantillon pour changement d'illuminant (ISO 18314-4:2020)

OSIST prEN ISO 18314-4:2021

https://standards.iteh.ai/catalog/standards/sist/48bb74e6-8910-43e6-80b5-

Ta slovenski standard je istoveten⁸2:5/osist-prEN ISO 18314-4

ICS:

17.180.20 Barve in merjenje svetlobe Colours and measurement of

ight

87.060.10 Pigmenti in polnila Pigments and extenders

oSIST prEN ISO 18314-4:2021 en,fr,de

iTeh STANDARD PREVIEW (standards.iteh.ai)

oSIST prEN ISO 18314-4:2021 https://standards.iteh.ai/catalog/standards/sist/48bb74e6-8910-43e6-80b5-2be2705a8c25/osist-pren-iso-18314-4-2021

INTERNATIONAL STANDARD

ISO 18314-4

First edition 2020-08

Analytical colorimetry —

Part 4:

Metamerism index for pairs of samples for change of illuminant

Analyse colorimétrique —

iTeh STPartie 4: Indice de metamérisme de paires d'échantillon pour changement d'illuminant (standards.iteh.ai)

oSIST prEN ISO 18314-4:2021 https://standards.iteh.ai/catalog/standards/sist/48bb74e6-8910-43e6-80b5-2be2705a8c25/osist-pren-iso-18314-4-2021



iTeh STANDARD PREVIEW (standards.iteh.ai)

oSIST prEN ISO 18314-4:2021 https://standards.iteh.ai/catalog/standards/sist/48bb74e6-8910-43e6-80b5-2be2705a8c25/osist-pren-iso-18314-4-2021



COPYRIGHT PROTECTED DOCUMENT

© ISO 2020

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office CP 401 • Ch. de Blandonnet 8 CH-1214 Vernier, Geneva Phone: +41 22 749 01 11 Email: copyright@iso.org Website: www.iso.org

Published in Switzerland

Cor	Contents					
Fore	word		iv			
Intro	oductio	n	v			
1	Scop	ne	1			
2	Nor	1				
3	Terr	1				
4	Sym	2				
5	Reference illuminant					
6	Test illuminant					
7	CIELAB coordinates L*, a*, b*					
8	Meta	amerism index for change in illuminant General calculation methods	4			
	8.1	General calculation methods	4			
	8.2	Basic calculation of the metamerism index from colour differences	4			
	8.3	Correction methods				
		8.3.1 Additive correction	5			
		8.3.2 Multiplicative correction	5			
		8.3.3 Spectral correction	6			
	8.4	Test report	9			
Anno	ex A (ir	Test report. formative) Calculation examples RD PREVIEW	10			
Bibli	iograp	ny (standards.iteh.ai)	23			

<u>oSIST prEN ISO 18314-4:2021</u> https://standards.iteh.ai/catalog/standards/sist/48bb74e6-8910-43e6-80b5-2be2705a8c25/osist-pren-iso-18314-4-2021

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see www.iso.org/iso/foreword.html. (Standards.iteh.ai)

This document was prepared by Technical Committee ISO/TC 256, *Pigments, dyestuff and extenders*.

A list of all parts in the ISO 18314 series can be found on the ISO website.0-43e6-80b5-

2be2705a8c25/osist-pren-iso-18314-4-2021

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

Introduction

For the phenomenon of metamerism of pairs of samples, three different kinds are distinguished:

- a) Illuminant metamerism occurs if both of the object colours of a pair of samples are perceived as being the same only under a specific illuminant (e.g. under illuminant D65), while they differ under a different illuminant (e.g. illuminant A).
- b) Observer metamerism occurs if the object colours of a pair of samples are perceived as being the same by one observer, while a different observer perceives a colour difference under the same illuminant and the same reference conditions.
- c) Field-size metamerism occurs if both of the object colours of a pair of samples are perceived as being the same on the retina for a size of an observation field (e.g. defined by the 2° standard observer), while they differ for a different observation field on the retina (e.g. 10°).
 - NOTE 2 The reason for field-size metamerism is based on the existent colour matching functions of an observer during an observation situation. The colour matching functions change with the size of the observation field on the retina. Such change of the observation field can also occur if, for example, the pair of samples is examined from different distances.

iTeh STANDARD PREVIEW (standards.iteh.ai)

<u>oSIST prEN ISO 18314-4:2021</u> https://standards.iteh.ai/catalog/standards/sist/48bb74e6-8910-43e6-80b5-2be2705a8c25/osist-pren-iso-18314-4-2021

iTeh STANDARD PREVIEW (standards.iteh.ai)

oSIST prEN ISO 18314-4:2021 https://standards.iteh.ai/catalog/standards/sist/48bb74e6-8910-43e6-80b5-2be2705a8c25/osist-pren-iso-18314-4-2021

Analytical colorimetry —

Part 4:

Metamerism index for pairs of samples for change of illuminant

1 Scope

This document specifies a formalism for the calculation of the illuminant metamerism of solid surface colours. It cannot be applied to colours of effect coatings without metrical adaptation.

This document only covers the phenomenon of metamerism for change of illuminant, which has the greatest meaning in practical application. In the case of chromaticity coordinates of a pair of samples under reference conditions that do not exactly match, recommendations are given on which correction measures are to be taken. Regarding the reproduction of colours, the metamerism index is used as a measure of quality in order to specify tolerances for colour differences between a colour sample and a colour match under different illumination conditions.

The quantification of the illuminant metamerism of pairs of samples is formally performed by a colour difference assessment, for which tolerances that are common for the evaluation of residual colour differences can be used. (standards.iteh.ai)

NOTE In the colorimetric literature and textbooks, the term geometric metamerism is sometimes used for the case that two colours appear to be the same under a specific geometry for visual assessment and selected standard observer and standard illuminant pair, but is perceived as two different colours at changed observation geometry. The term geometric metamerism is different to metamerism described in this document.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO/CIE 11664-1, Colorimetry — Part 1: CIE standard colorimetric observers

ISO/CIE 11664-2:—¹⁾, Colorimetry — Part 2: CIE standard illuminants

ISO/CIE 11664-4, Colorimetry — Part 4: CIE 1976 L*a*b* colour space

CIE 015, Colorimetry

CIE S 017, International Lighting Vocabulary

3 Terms and definitions

For the purposes of this document, the terms and definitions given in CIE S 017 and the following apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at https://www.iso.org/obp
- IEC Electropedia: available at http://www.electropedia.org/

¹⁾ Under preparation. Stage at the time of preparation: ISO/CIE DIS 11664-2:2020.

3.1

metamerism

property of spectrally different colour stimuli that have the same tristimulus values in a specified colorimetric system

[SOURCE: CIE 017:2016, 17-23-006]

3.2

paramerism

characteristic of a pair of samples with spectral colour stimulus functions which have different fundamental colour stimulus functions as well as different residuals or metameric black values within the visible spectral range

Note 1 to entry: Parameric objects are characterized by the fact that they reflect colour stimuli of different spectral power distribution functions under a specified standard illuminant, which cause approximately the same colour perception under the selected observation conditions.

3.3

colour difference

 ΔE^*

difference between two colour stimuli, defined as a distance between the points representing them in a specified colour space

[SOURCE: CIE 017:2016, 17-22-041, modified — symbol ΔE^* was amended, "Euclidean" and Note 1 to entry have been deleted.

3.4

reference illuminant (standards.iteh.ai) illuminant with which other illuminants are compared

[SOURCE: CIE S 017:2016, 17-22-109 17] <u>oSIST prEN ISO 18314-4:2021</u>

https://standards.iteh.ai/catalog/standards/sist/48bb74e6-8910-43e6-80b5-

iTeh STANDARD PREVIEW

3.5 2be2705a8c25/osist-pren-iso-18314-4-2021

test illuminant

illuminant, for which the colour difference (3.3) between the two samples to be tested is assessed

3.6

metamerism-index for change in illuminant

 M_{t}

colour difference ΔE^* (3.3) between the two samples under test illuminant (3.5) if $\Delta E^* = 0$ is observed under the reference illuminant (3.4)

3.7

correction method

algorithm for theoretically eliminating a *colour difference* (3.3) of the pair of samples under the *reference* illuminant (3.4)

Symbols and abbreviated terms

For the application of this document, the symbols given in <u>Table 1</u> apply.

Table 1 — Symbols

Symbol	Identification				
X, Y, Z	Standard tristimulus values of a measured object colour				
X_n , Y_n , Z_n	Standard tristimulus values of the used illuminant				
\overline{X} , \overline{Y} , \overline{Z}	Colour-matching functions				

Table 1 (continued)

Symbol	Identification				
L^* , a^* , b^*	Basic coordinates of the CIELAB system				
ΔL^* , Δa^* , Δb^*	Differences between basic coordinates of the CIELAB system				
M_{t}	Metamerism index for change in illuminant				
\vec{N} , $\vec{N}_{\rm f}$, $\vec{N}_{\rm r}$	Vector of the radiometric function of a sample with associated fundamental colour stimulus (f) and metameric black (r)				
λ	Wavelength				
S	Relative spectral distribution function of an illuminant				
\vec{W}	Vector of the standard tristimulus values				
w	Integration weights for the calculation of the standard tristimulus values				
A	Matrix of the integration weights <i>w</i> for the calculation of the standard tristimulus values				
R	Projection matrix				
I	Identity matrix				
Index spl	Sample				
Index std	Standard				
Index t	Test illuminant				
Index corr	Corrected value				
Index f	Fundamental colour stimulus A R D PREVIEW				
Index r	Metameric black values (residuals)				
Index ref	Reference illuminant and ards.iteh.ai)				
Index T	Transposed matrix				

https://standards.iteh.ai/catalog/standards/sist/48bb74e6-8910-43e6-80b5-

5 Reference illuminant be2705a8c25/osist-pren-iso-18314-4-2021

The standard illuminant D65 is chosen as reference illuminant in accordance with ISO/CIE 11664-2. Other reference illuminants required in special cases shall be particularly specified.

6 Test illuminant

The selection of the test illuminant depends on the application. If the test illuminants are not particularly specified, standard illuminant A in accordance with ISO 11664-2 and/or illuminants of the fluorescent lamp type, such as FL11 in accordance with CIE 015, shall preferably be selected. The test illuminant used shall be indicated as an index to M, e.g. M_A or $M_{\rm FL,11}$.

When calculating the standard tristimulus values *X*, *Y*, *Z* under the selected test illuminants, the basic raster of wavelengths given in ISO 11664-2 or CIE 015 for A and D65, and in CIE 015 for FL11 and FL2 shall be complied with. In cases of missing measuring values of the standard or sample for these wavelengths, these values shall be interpolated and/or extrapolated.

7 CIELAB coordinates L^* , a^* , b^*

The metamerism index $M_{\rm t}$ is based on the CIELAB coordinates L^* , a^* , b^* of samples 1 and 2 which are to be compared. L^* , a^* , b^* is calculated in accordance with ISO/CIE 11664-4 from the standard tristimulus values X, Y, Z of the sample for the CIE 1964 10° standard observer in accordance with ISO/CIE 11664-1 for the reference illuminant and the selected test illuminant. If calculating L^* , a^* , b^* under the test illuminant, the respective standard tristimulus values $X_{\rm n}$, $Y_{\rm n}$, $Z_{\rm n}$ of the entirely matt white surface shall be used (see CIE 015). For the standard illuminants A and D65 or for the illuminant

recommendation FL11, the standard tristimulus values X_n , Y_n , Z_n of the entirely matt white surface apply in accordance with Table 2.

Table 2 specifies standard tristimulus values for the frequently used standard illuminants D65 and A as well as illuminant FL11 and both of the standard observers in accordance with CIE 015.

	2° standard observer			10° standard observer				
Standard tristimulus values	Illuminant							
	D65	A	FL11	D65	A	FL11		
X _n	95,04	109,85	100,96	94,81	111,14	103,86		
Y _n	100,00	100,00	100,00	100,00	100,00	100,00		
$Z_{ m n}$	108,88	35,58	64,35	107,32	35,20	65,61		

Table 2 — Standard tristimulus values

For fluorescent samples, the illuminant used for measurement shall be adjusted as close as possible to that illuminant for which the standard tristimulus values are to be determined.

In contrast to non-fluorescent samples, the calculation of metamerism indices for fluorescent samples is erroneous if the samples are measured only under one illuminant.

Metamerism index for change in illuminant PREVIEW (standards.iteh.ai)

8.1 General calculation methods

Three different correction methods for calculating a metamerism index in the case of paramerism have been proposed in References [6] to [13]. All methods assume that, for practical cases, there might be already a small difference between the colours of the sample and the standard even under the reference illuminant from the very beginning, due to problems of fabrication. In the case of two methods, called the additive and the multiplicative correction, these inherent colour differences often merge with the difference introduced by the change of the illuminant. The third method, the spectral correction, works more fundamentally by the separation of inherent colour differences under the reference illuminant from those introduced by the change of the illuminant.

NOTE Annex A includes calculation examples.

Basic calculation of the metamerism index from colour differences 8.2

The common formula for a metamerism index at change in illuminant, expressed in CIELAB coordinates for the test illuminant (t), is given by Formula (1):

$$M_{t} = \sqrt{\left(\Delta L_{t}^{*}\right)^{2} + \left(\Delta a_{t}^{*}\right)^{2} + \left(\Delta b_{t}^{*}\right)^{2}} \tag{1}$$

where

is the test colour; t

$$\Delta L_{\rm t}^* = L_{\rm spl,corr,t}^* - L_{\rm std,t}^*;$$

$$\begin{split} \Delta a_{\mathsf{t}}^* &= a_{\mathsf{spl,corr},\mathsf{t}}^* - a_{\mathsf{std,t}}^*; \\ \Delta b_{\mathsf{t}}^* &= b_{\mathsf{spl,corr},\mathsf{t}}^* - b_{\mathsf{std,t}}^*. \end{split}$$

$$\Delta b_{\rm t}^* = b_{\rm spl,corr,t}^* - b_{\rm std,t}^*$$