

SLOVENSKI STANDARD kSIST-TS FprCEN/TS 17786-2:2021

01-december-2021

[Not translated]

Inorganic micronutrient fertilizers - Determination of the chelated micronutrient content and the chelated fraction of micronutrients - Part 2: Determination of EDTA, DTPA, HEEDTA, IDHA or EDDS

Anorganische Spurennährstoffdüngemittel - Bestimmung des Gehaltes an chelatisierten Spurennährstoffen und des chelatisierten Anteils an Spurennährstoffen - Teil 2: Bestimmung von EDTA, DTPA, HEEDTA, IDHA oder EDDS

(standards.iteh.ai)

<u>kSIST-TS FprCEN/TS 17786-2:2021</u> https://standards.iteh.ai/catalog/standards/sist/0dc6d8c6-0f76-4506-9fa6-

Ta slovenski standard je istoveten z: FprcEN/TS 17786-2

ICS:

65.080 Gnojila Fertilizers

kSIST-TS FprCEN/TS 17786-2:2021 en,fr,de

kSIST-TS FprCEN/TS 17786-2:2021

iTeh STANDARD PREVIEW (standards.iteh.ai)

kSIST-TS FprCEN/TS 17786-2:2021 https://standards.iteh.ai/catalog/standards/sist/0dc6d8c6-0f76-4506-9fa6-59efe9614788/ksist-ts-fprcen-ts-17786-2-2021

TECHNICAL SPECIFICATION SPÉCIFICATION TECHNIQUE TECHNISCHE SPEZIFIKATION

FINAL DRAFT FprCEN/TS 17786-2

October 2021

ICS 65.080

English Version

Inorganic micronutrient fertilizers - Determination of the chelated micronutrient content and the chelated fraction of micronutrients - Part 2: Determination of EDTA, DTPA, HEEDTA, IDHA or EDDS

Anorganische Spurennährstoffdüngemittel -Bestimmung des Gehaltes an chelatisierten Spurennährstoffen und des chelatisierten Anteils an Spurennährstoffen - Teil 2: Bestimmung von EDTA, DTPA, HEEDTA, IDHA oder EDDS

This draft Technical Specification is submitted to CEN members for Vote. It has been drawn up by the Technical Committee CEN/TC 260.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Republic of North Macedonia, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.

https://standards.itch.ai/catalog/standards/sist/0dc6d8c6-0f76-4506-9fa6-

Recipients of this draft are invited to submit, with their comments; notification of any relevant patent rights of which they are aware and to provide supporting documentation.

Warning: This document is not a Technical Specification. It is distributed for review and comments. It is subject to change without notice and shall not be referred to as a Technical Specification.



EUROPEAN COMMITTEE FOR STANDARDIZATION COMITÉ EUROPÉEN DE NORMALISATION EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Rue de la Science 23, B-1040 Brussels

European foreword	3
Introduction	4
1 Scope	5
2 Normative references	5
3 Terms and definitions	5
4 Principle	6
5 Interferences	6
6 Reagents	6
7 Apparatus	
8 Sampling and sample preparation	7
9 Procedure	7
 9.2 Water-soluble micronutrient content 9.3 Chelating agent content 	
10 Expression of results(standards.iteh.ai)	8
10.1 Water-soluble micronutrient content	8
10.2 Chelating agent contentksist-ts-FprGEN/TS-17786-22021	8
10.3 Chelated fraction (ChF) dards, itch ai/ontolog/standards/sist/0do6d8c6-0f76-4506-0fa6	
11 Test report	9
Annex A (informative) Complete names of chelating agents	10
Bibliography	11

European foreword

This document (FprCEN/TS 17786-2:2021) has been prepared by Technical Committee CEN/TC 260 "Fertilizers and liming materials", the secretariat of which is held by DIN.

This document is currently submitted to the Vote on TS.

This document has been prepared under a Standardization Request given to CEN by the European Commission and the European Free Trade Association.

iTeh STANDARD PREVIEW (standards.iteh.ai)

<u>kSIST-TS FprCEN/TS 17786-2:2021</u> https://standards.iteh.ai/catalog/standards/sist/0dc6d8c6-0f76-4506-9fa6-59efe9614788/ksist-ts-fprcen-ts-17786-2-2021

Introduction

Micronutrients are considered to be, in plant nutrition, a number of elements known to be needed in small amounts for proper plant growth and development. The most common are Iron (Fe), Manganese (Mn), Molybdenum (Mo), Copper (Cu), Zinc (Zn) and Boron (B).

If an inorganic micronutrient fertilizer contains a substance, or one of the substances in the mixture, which is intended to enhance the long term availability to plants of micronutrients in the EU fertilizing product, that substance is either a chelating agent or a complexing agent.

The chelating agents are divided into two groups1:

- Group 1: EDTA, DTPA, HEEDTA, IDHA and [S,S]-EDDS;
- Group 2: Chelating agents present in UVCB (unknown or variable composition, complex reaction products and biological materials) chelates including [o,o] EDDHA, [o,p] EDDHA, [o,o] EDDHMA, HBED and EDDHSA.

This document defines the test method to be used in order to measure the compliance with the chelated fraction of micronutrients in product function category (PFC) 1(C) (II) (classified according to Regulation (EU) 2019/1009 [7]) as inorganic micronutrient fertilizer containing one or more chelating agents of Group 1.

iTeh STANDARD PREVIEW (standards.iteh.ai)

<u>kSIST-TS FprCEN/TS 17786-2:2021</u> https://standards.iteh.ai/catalog/standards/sist/0dc6d8c6-0f76-4506-9fa6-59efe9614788/ksist-ts-fprcen-ts-17786-2-2021

_

¹ Abbreviated terms are described in Annex A.

1 Scope

This document specifies a method for the determination of the chelated fraction of micronutrients for fertilizers when one or many micronutrients are chelated by EDTA, DTPA, HEEDTA, IDHA or [S,S]-EDDS in fertilizers.

This method is used for inorganic micronutrient fertilizers when micronutrients are chelated only by EDTA, DTPA, HEEDTA, IDHA or [S,S]-EDDS or for mixtures in which EDTA, DTPA, HEEDTA, IDHA or [S,S]-EDDS is one of the chelating agents.

The method is applicable to all inorganic micronutrient fertilizers containing EDTA, DTPA, HEEDTA, IDHA or [S,S]-EDDS as chelating agent for contents > 0,1% (g/100 g).

The method is based on ICP or AAS measurement of the concentration of micronutrients according to EN 16963 or EN 16965 after water extraction according to EN 16962 and LC measurement of the chelating agents according to EN 15950, EN 13368-1 and EN 13368-3.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

iTeh STANDARD PREVIEW

EN 12944-1, Fertilizers and liming materials and soil improvers - Vocabulary - Part 1: General terms

EN 12944-2, Fertilizers and liming materials and soil improvers - Vocabulary - Part 2: Terms relating to fertilizers

https://standards.iteh.ai/catalog/standards/sist/0dc6d8c6-0f76-4506-9fa6-

EN 13368-1, Fertilizers - Determination of chelating agents in fertilizers by chromatography - Part 1: Determination of EDTA, HEEDTA and DTPA by ion chromatography

EN 13368-3, Fertilizers - Determination of chelating agents in fertilizers by chromatography - Part 3: Determination of [S,S]-EDDS by ion pair chromatography

EN 15950, Fertilizers - Determination of N-(1,2-dicarboxyethyl)-D,L-aspartic acid (Iminodisuccinic acid, IDHA) using high-performance liquid chromatography (HPLC)

EN 16962, Fertilizers - Extraction of water soluble micro-nutrients in fertilizers and removal of organic compounds from fertilizer extracts

EN 16963, Fertilizers - Determination of boron, cobalt, copper, iron, manganese, molybdenum and zinc using ICP-AES

EN 16965, Fertilizers - Determination of cobalt, copper, iron, manganese and zinc using flame atomic absorption spectrometry (FAAS)

3 Terms and definitions

For the purposes of this document, the terms and definitions given in EN 12944-1 and EN 12944-2 and the following apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

— ISO Online browsing platform: available at https://www.iso.org/obp

IEC Electropedia: available at https://www.electropedia.org/

3.1

chelated fraction

ChF

chelating agent content divided by the water-soluble micronutrient content

Note 1 to entry: The chelated fraction (ChF) is expressed as a percentage.

4 Principle

The principle of the method is to determine the content of micronutrients and the content of one or more chelating agents declared in the fertilizer followed by calculating the chelated fraction.

The content of micronutrients is determined using ICP-AES (inductive coupled plasma atomic emission spectrometry) or FAAS (flame atomic absorption spectrometry) methods using EN 16963 or EN 16965 standards after previous extraction using the EN 16962 method.

The content of the declared chelating agent is determined using one of the following methods:

- for EDTA, DTPA and HEEDTA, EN 13368-1 shall be used;
- for [S,S]-EDDS, EN 13368-3 shall be used;
- for IDHA, EN 15950 shall be used TANDARD PREVIEW

The chelated fraction is calculated by comparing the water soluble micronutrient content with the chelating agent content.

5 Interferences

kSIST-TS FprCEN/TS 17786-2:2021

https://standards.iteh.ai/catalog/standards/sist/0dc6d8c6-0f76-4506-9fa6-

Other chelating agents such as DTPA, [o,o] EDDHA or [o,p] EDDHA do not interfere the determination of IDHA. EDTA may interfere with the determination of IDHA with some equipment, especially with certain columns in the HPLC-equipment.

In case of [S,S]-EDDS no interferences have been detected. Metal chelates with [o,o] EDDHA, [o,o] EDDHMA, HBED, EDDHSA, EDTA, DTPA, HEEDTA, IDHA, [o,p] EDDHA, lignosulfonates and heptagluconates as well as the chelating agents do not interfere since after Cu derivatization they are separate from Cu-[S,S]-EDDS. Since retention times depend on the column type, interferences should be checked if a mixture with other chelating or complexing agents is suspected.

6 Reagents

6.1 Water

All water used should conform to EN ISO 3696 and be degassed.

6.2 Reagents of recognized analytical grade.

Those included in the following standards shall be used:

- EN 16962, when extracting boron, cobalt, copper, iron, manganese, molybdenum and zinc;
- EN 16963, when determining the content of boron, cobalt, copper, iron, manganese, molybdenum and zinc by ICP-AES;

- EN 16965, when determining the content of cobalt, copper, iron, manganese and zinc by FAAS;
- EN 13368-1, when determining the content of EDTA, HEEDTA and DTPA;
- EN 13368-3, when determining the content of [S,S]-EDDS;
- EN 15950, when determining the content of IDHA.

7 Apparatus

The usual laboratory apparatus and, in particular, the following:

- **7.1 Rotary shaker**, capable of operation at a rotational speed of about 35 min⁻¹ to 45 min⁻¹.
- **7.2 Inductive coupled plasma-atomic emission spectrometer**, with axial or radial viewing of the plasma and with suitable background correction.
- **7.3 Atomic absorption spectrometer**, equipped with hollow cathode lamp or other suitable light emission source, background correction and burner.
- **7.4 Chromatograph,** equipped with pump, injection valve and injection loop. The columns included in the following standards shall we used: DARD PREVIEW
- EN 13368-1 when determining the content of EDTA, HEEDTA and DTPA;
- EN 13368-3 when determining the content of [S,S]-EDDS;
- EN 15950 when determining the content of IDHA.
- **7.5 Membrane filters,** micro membrane filters, resistant to aqueous solutions, with porosity of respectively $0.45 \mu m$ and $0.2 \mu m$.

8 Sampling and sample preparation

Sampling and sample preparation are not part of the method specified in this document. A recommended sampling method is given in EN 1482-1 and a recommended sample preparation method is given in EN 1482-2.

For the size reduction of samples with a high amount of chelating agents, it is not recommended to use a high speed laboratory mill. It is more convenient to grind the sample to a particle size less than 1 mm with a mortar. Special care shall be taken with NPK samples due to their high hygroscopicity.

9 Procedure

9.1 General

For the determination of the micronutrient content and the content of the chelating agents, proceed in accordance with the procedures included in the standards mentioned below.

9.2 Water-soluble micronutrient content

Perform the extraction of the water-soluble micronutrients according to EN 16962 and determine the content of the water soluble micronutrients according to EN 16963 or EN 16965.

9.3 Chelating agent content

Determine the content of the chelating agent(s) by one or more of the following standards: EN 13368-1; EN 13368-3; EN 15950. Perform the analysis according to the procedures included in these standards.

NOTE To determine the total content of chelating agents, the adequate standard to determine the chelating agents can be based on, yet not limited to, the declared chelating agents.

10 Expression of results

10.1 Water-soluble micronutrient content

The water-soluble micronutrient concentration obtained from EN 16963 or EN 16965 presented as w_x in percent, should be calculated in millimoles per 100 grams by Formula (1):

$$c_{\rm x} = \frac{w_{\rm x}}{M_{\rm x}} \times 1000 \tag{1}$$

where

 $c_{\rm x}$ is the concentration of water-soluble micronutrient, in mmol per 100 g;

 $w_{\rm x}$ is the mass fraction of water-soluble micronutrient, in g per 100 g;

 $M_{\rm x}$ is the molar mass of the water-soluble micronutrient, in g per mol.

In the case of a mixture of micronutrients calculate the sum of all the micronutrients expressed in millimoles per 100 grams. https://standards.iteh.ai/catalog/standards/sist/0dc6d8c6-0f76-4506-9fa6-

10.2 Chelating agent content 59efe9614788/ksist-ts-fprcen-ts-17786-2-2021

The mass fraction of the chelating agents EDTA, HEEDTA and DTPA obtained from EN 13368-1 and [S,S]-EDDS obtained from EN 13368-3 is presented as $w_{\rm L}$ in percent, expressed as free acid. Convert the result to millimoles per 100 grams as given by Formula (2):

$$c_{\rm LCh} = \frac{w_{\rm L}}{M_{\rm I}} \times 1\,000\tag{2}$$

where

 c_{LCh} is the content of the chelating agent, expressed as free acid, in mmol per 100 g;

 $w_{\rm L}$ is the mass fraction of the chelating agent, expressed as free acid, in g per 100 g;

 $\rm M_L$ is the molar mass, in g/mol, of the chelating agent, expressed as free acid. NOTE: for EDTA: 292 g/mol, HEEDTA: 278 g/mol, DTPA: 393 g/mol [S,S]-EDDS: 292,2 g/mol and for IDHA 249,1 g/mol.

The chelating agent content ($C_{L,total}$) in mmol/100 g is the sum of the content of EDTA, HEEDTA, DTPA, [S,S]-EDDS and IDHA, expressed as free acid, in millimoles per 100 grams.