
Živalske in rastlinske maščobe ter olja - Določevanje maščobnokislinsko vezanih kloropropanediolov (MCPD) in glicidola z GC/MS - 3. del: Metoda z uporabo kislinske transesterifikacije in meritev 2-MCPD, 3-MCPD in glicidola (ISO 18363-3:2017)

Animal and vegetable fats and oils - Determination of fatty-acid-bound chloropropanediols (MCPDs) and glycidol by GC/MS - Part 3: Method using acid transesterification and measurement for 2-MCPD, 3-MCPD and glycidol (ISO 18363-3:2017)

Tierische und pflanzliche Fette und Öle - Bestimmung von fettsäuregebundenen Chlorpropanediol (MCPD) und Glycidol mittels GC/MS - Teil 3: Verfahren mittels Säureumesterung und Messung für 2 MCPD, 3 MCPD und Glycidol (ISO 18363 3:2017)

Corps gras d'origines animale et végétale - Détermination des esters de chloropropanediols (MCPD) et d'acides gras et des esters de glycidol et d'acides gras par CPG/SM - Partie 3: Méthode par transestérification acide et mesure du 2-MCPD, du 3-MCPD et du glycidol (ISO 18363-3:2017)

Ta slovenski standard je istoveten z: prEN ISO 18363-3

ICS:

67.200.10	Rastlinske in živalske maščobe in olja	Animal and vegetable fats and oils
-----------	--	------------------------------------

oSIST prEN ISO 18363-3:2021

en,fr,de

INTERNATIONAL
STANDARD

ISO
18363-3

First edition
2017-12

**Animal and vegetable fats and oils —
Determination of fatty-acid-bound
chloropropanediols (MCPDs) and
glycidol by GC/MS —**

Part 3:

**Method using acid transesterification
and measurement for 2-MCPD,
3-MCPD and glycidol**

*Corps gras d'origines animale et végétale — Détermination des
esters de chloropropanediols (MCPD) et d'acides gras et des esters de
glycidol et d'acides gras par CPG/SM —*

*Partie 3: Méthode par transestérification acide et mesure pour le
2-MCPD, le 3-MCPD et le glycidol*

<https://standards.iteh.ai/catalog/standards/sist/756ad273-ceec-4700-900f-34c29cb95655/sist-en-iso-18363-3-2022>



Reference number
ISO 18363-3:2017(E)

© ISO 2017

iTeh Standards
(<https://standards.iteh.ai>)
Document Preview

[SIST EN ISO 18363-3:2022](https://standards.iteh.ai/catalog/standards/sist/756ad273-ceec-4700-900f-34c29cb95655/sist-en-iso-18363-3-2022)

<https://standards.iteh.ai/catalog/standards/sist/756ad273-ceec-4700-900f-34c29cb95655/sist-en-iso-18363-3-2022>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2017, Published in Switzerland

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
Ch. de Blandonnet 8 • CP 401
CH-1214 Vernier, Geneva, Switzerland
Tel. +41 22 749 01 11
Fax +41 22 749 09 47
copyright@iso.org
www.iso.org

Contents

Page

Foreword	iv
Introduction	v
1 Scope	1
2 Normative references	1
3 Terms and definitions	1
4 Principle	2
5 Reagents	2
5.1 Standard and reference compounds.....	2
5.2 Standard solutions.....	3
5.2.1 General.....	3
5.2.2 Stock solutions (1 mg/ml).....	3
5.2.3 Working solutions.....	3
5.3 Other reagents.....	4
5.4 Reagent solutions.....	4
6 Apparatus	5
7 Sample	5
7.1 Sampling.....	5
7.2 Preparation of the test sample.....	5
8 Procedure	6
8.1 Test sample preparation.....	6
8.2 Preparation of the calibration curve.....	7
8.3 Gas chromatography/mass spectrometry references.....	7
9 Expression of results	8
9.1 Quantification of 3-MCPD esters.....	8
9.2 Quantification of 2-MCPD esters.....	9
9.3 Quantification of glycidyl esters.....	9
10 Precision	10
10.1 General.....	10
10.2 Repeatability.....	10
10.3 Between-day reproducibility.....	10
11 Test report	11
Annex A (informative) Construction of the calibration curves	12
Annex B (informative) Results of the interlaboratory test	16
Bibliography	18

ISO 18363-3:2017(E)

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see the following URL: www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 11, *Animal and vegetable fats and oils*.

A list of all parts in the ISO 18363 series can be found on the ISO website.

[SIST EN ISO 18363-3:2022](https://standards.iteh.ai/catalog/standards/sist/756ad273-ceec-4700-900f-34c29cb95655/sist-en-iso-18363-3-2022)

<https://standards.iteh.ai/catalog/standards/sist/756ad273-ceec-4700-900f-34c29cb95655/sist-en-iso-18363-3-2022>

Introduction

The ISO 18363 series is a family of International Standards which can be used for the determination of ester-bound MCPD and glycidol. This introduction describes the methods specified in the three documents currently published or proposed so that the analyst can decide which methods are suitable for application. The detailed application of each method is contained within the scope of the individual method.

ISO 18363-1 is a differential method equivalent to the DGF standard C-VI 18 (10) and identical to AOCS Official Method Cd 29c-13. Briefly, it is based on a fast alkaline catalysed release of 3-MCPD and glycidol from the ester derivatives. Glycidol is subsequently converted into induced 3-MCPD. It consists of two parts. The first part (A) allows the determination of the sum of ester-bound 3-MCPD and ester-bound glycidol, whereas the second part (B) determines ester-bound 3-MCPD only. Both assays are based on the release of the target analytes 3-MCPD and glycidol from the ester-bound form by an alkaline catalysed alcoholysis carried out at room temperature. In part A, an acidified sodium chloride solution is used to stop the reaction and subsequently convert the glycidol into induced 3-MCPD. Thus, 3-MCPD and glycidol become indistinguishable in part A. In part B, the reaction stop is achieved by the addition of an acidified chloride-free salt solution which also prevents the conversion of glycidol into induced MCPD. Thereby, part B allows the determination of the genuine 3-MCPD content. Finally, the glycidol content of the sample is proportional to the difference of both assays (A – B) and can be calculated when the transformation ratio from glycidol to 3-MCPD has been determined. ISO 18363-1 is applicable to the fast determination of ester-bound 3-MCPD and glycidol in refined and non-refined vegetable oils and fats. ISO 18363-1 can also apply to animal fats and used frying oils and fats, but a validation study has to be undertaken before the analysis of these matrices. Any free analytes within the sample would be included in the results, but the document does not allow the distinction between free and bound analytes. However, as of publication, research has not shown any evidence of a free analyte content as high as the esterified analyte content in refined vegetable oils and fats. In principle, ISO 18363-1 can also be modified in such a way that the determination of 2-MCPD is feasible, but again, a validation study has to be undertaken before the analysis of this analyte.

ISO 18363-2 (proposed) will represent AOCS Official Method Cd 29b-13. Briefly, it will be based on a slow alkaline release of MCPD and glycidol from the ester derivatives. Glycidol is subsequently converted into 3-MBPD. ISO 18363-2 will consist of two sample preparations that differ in the use of internal standards. Both preparations will be used for the determination of ester-bound 2-MCPD and 3-MCPD. In part A, a preliminary result for ester-bound glycidol will be determined. Because the 3-MCPD present in the sample will be converted to some minor extent into induced glycidol by the sample preparation, part B will serve to quantify this amount of induced glycidol that is subsequently subtracted from the preliminary glycidol result of part A. By the use of isotopically labelled free MCPD isomers in assay A and isotopically labelled ester-bound 2-MCPD and 3-MCPD in part B, the efficiency of ester cleavage will be able to be monitored. Both assays A and B will be based on the release of the target analytes 2-MCPD, 3-MCPD, and glycidol from the ester-bound form by a slow alkaline catalysed alcoholysis in the cold. In both sample preparations, the reaction will be stopped by the addition of an acidified concentrated sodium bromide solution so as to convert the unstable and volatile glycidol into 3-MBPD which shows comparable properties to 3-MCPD with regard to its stability and chromatographic performance. Moreover, the major excess of bromide ions will prevent the undesired formation of 3-MCPD from glycidol in the case of samples which contain naturally occurring amounts of chloride. ISO 18363-2 will be applicable to the determination of ester-bound 3-MCPD, 2-MCPD, and glycidol in refined and unrefined vegetable oils and fats. It will also apply to animal fats and used frying oils and fats, but a validation study will have to be undertaken before the analysis of these matrices. Any free analytes within the sample would be included in the results, but the document will not allow the distinction between free and bound analytes. However, as of publication of this document, research has not shown any evidence of a free analyte content as high as the esterified analyte content in vegetable oils and fats.

This document represents AOCS Official Method Cd 29a-13. Briefly, it is based on the conversion of glycidyl esters into 3-MBPD esters and a slow acid catalysed release of MCPD and MBPD from the ester derivatives. This document is based on a single sample preparation in which glycidyl esters are converted into MBPD monoesters, and subsequently, the free analytes 2-MCPD, 3-MCPD, and 3-MBPD are released by a slow acid catalysed alcoholysis. The 3-MBPD represents the genuine content of bound glycidol. This document can be applied for the determination of ester-bound 2-MCPD, 3-MCPD, and

ISO 18363-3:2017(E)

glycidol in refined and non-refined vegetable oils and fats. It can also apply to animal fats and used frying oils and fats, but a validation study has to be undertaken before the analysis of these matrices. The method is suited for the analysis of bound (esterified) analytes, but if required, this document can be also performed without the initial conversion of glycidyl esters. In such a setup, both free and bound 2-MCPD and 3-MCPD forms would be included in the results and the amount of free analytes can be calculated as a difference between two determinations performed in both setups. However, as of publication of this document, research has not shown any evidence of a free analyte content as high as the esterified analyte content in vegetable oils and fats.

iTeh Standards
(<https://standards.iteh.ai>)
Document Preview

[SIST EN ISO 18363-3:2022](https://standards.iteh.ai/catalog/standards/sist/756ad273-ceec-4700-900f-34c29cb95655/sist-en-iso-18363-3-2022)

<https://standards.iteh.ai/catalog/standards/sist/756ad273-ceec-4700-900f-34c29cb95655/sist-en-iso-18363-3-2022>

Animal and vegetable fats and oils — Determination of fatty-acid-bound chloropropanediols (MCPDs) and glycidol by GC/MS —

Part 3:

Method using acid transesterification and measurement for 2-MCPD, 3-MCPD and glycidol

1 Scope

This document specifies a procedure for the simultaneous determination of 2-MCPD esters (bound 2-MCPD), 3-MCPD esters (bound 3-MCPD) and glycidyl esters (bound glycidol) in a single assay, based on acid catalysed ester cleavage and derivatization of cleaved (free) analytes with phenylboronic acid (PBA) prior to GC/MS analysis.

This document is applicable to solid and liquid fats and oils. For all three analytes the limit of quantification (LOQ) is 0,1 mg/kg and the limit of detection (LOD) is 0,03 mg/kg.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 3696, *Water for analytical laboratory use — Specification and test methods*

3 Terms and definitions

[SIST EN ISO 18363-3:2022](https://standards.iteh.ai/catalog/standards/sist/756ad273-ceec-4700-900f-34c29cb95655/sist-en-iso-18363-3-2022)

<https://standards.iteh.ai/catalog/standards/sist/756ad273-ceec-4700-900f-34c29cb95655/sist-en-iso-18363-3-2022>

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <http://www.electropedia.org/>
- ISO Online browsing platform: available at <https://www.iso.org/obp>

3.1

bound 2-MCPD

amount of 2-MCPD cleaved from its esterified (bound) forms by acid catalysed transesterification according to the reference method

Note 1 to entry: The content of 2-MCPD is calculated and reported as a mass fraction, in milligrams per kilogram (mg/kg).

3.2

bound 3-MCPD

amount of 3-MCPD cleaved from its esterified (bound) forms by acid catalysed transesterification according to the reference method

Note 1 to entry: The content of 3-MCPD is calculated and reported as a mass fraction, in milligrams per kilogram (mg/kg).

ISO 18363-3:2017(E)

3.3

bound glycidol

amount of glycidol cleaved from its esterified (bound) forms by acid catalysed transesterification according to the reference method

Note 1 to entry: The content of glycidol is calculated and reported as a mass fraction, in milligrams per kilogram (mg/kg).

4 Principle

The oil/fat sample is dissolved in tetrahydrofuran, and the internal standards (pentadeuterated 3-MCPD diester and pentadeuterated glycidyl ester) are added. During the first step of sample preparation, glycidyl esters are converted into 3-MBPD monoesters by the addition of an acidified solution of sodium bromide. Upon completion of the reaction, the organic phase, containing 2- and 3-MCPD esters and 3-MBPD esters, is separated and evaporated to dryness. In the second step the residue is dissolved in tetrahydrofuran and the acid transesterification is initiated by the addition of an acid alcoholic solution. After 16 h incubation at 40 °C, the sample mixture is neutralized and the fatty acid methyl esters generated during the transesterification are removed. Finally, the purified sample [containing cleaved (free) analytes] is derivatized with phenylboronic acid prior to GC/MS analysis.

The quantification of 2- and 3-MCPD esters (expressed as bound 2- and 3-MCPD) is based on the 2-MCPD/3-MCPD-d5 and 3-MCPD/3-MCPD-d5 signal ratio, respectively. The quantification of glycidyl esters (expressed as bound glycidol) is based on the 3-MBPD/3-MBPD-d5 signal ratio.

This method allows the simultaneous quantification of all three analytes in a single assay.

5 Reagents

WARNING — This document requires handling of hazardous substances. Technical, organizational, and personal safety measures shall be followed.

Unless otherwise stated, analytically pure reagents shall be used. Water shall comply with grade 3 of ISO 3696.

5.1 Standard and reference compounds

5.1.1 1,2-Dipalmitoyl-3-chloropropanediol (PP-3-MCPD), purity ≥ 95 % (e.g. from a supplier or synthesized from 3-MCPD and palmitoyl chloride as described by Reference [1]).

NOTE 1,2-Dipalmitoyl-3-chloropropanediol can be substituted by 1,2-dioleoyl-3-chloropropanediol or other fatty acid diesters of 3-MCPD with similar chain length (C16-C18 are preferred as they are the most abundant in the majority of oils/fats).

5.1.2 1,3-Dipalmitoyl-2-chloropropanediol (PP-2-MCPD), purity ≥ 95 % (e.g. synthesized from 2-MCPD and palmitoyl chloride as described by Reference [1]).

NOTE In analogy with the recommendations given for PP-3-MCPD, 1,3-dipalmitoyl-2-chloropropanediol can be substituted by other fatty acid diesters of 2-MCPD with similar chain length (C16-C18 are preferred as they are the most abundant in the majority of oils/fats).

5.1.3 Pentadeuterated 1,2-dipalmitoyl-3-chloropropanediol (PP-3-MCPD-d5), purity ≥ 95 %.

NOTE The same consideration applied to 1,2-dipalmitoyl-3-chloropropanediol is valid also for its pentadeuterated analogue, see the Note in 5.1.1.

5.1.4 Glycidyl palmitate (Gly-P), purity ≥ 98 %.