# INTERNATIONAL STANDARD

ISO 19698

First edition 2020-09

# Sludge recovery, recycling, treatment and disposal — Beneficial use of biosolids — Land application

Valorisation, recyclage, traitement et élimination des boues — Utilisation bénéfique des boues d'épuration — Épandage

# iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 19698:2020 https://standards.iteh.ai/catalog/standards/sist/af6c286e-db36-4c8b-aa18-211d71b79920/iso-19698-2020



# iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 19698:2020 https://standards.iteh.ai/catalog/standards/sist/af6c286e-db36-4c8b-aa18-211d71b79920/iso-19698-2020



# **COPYRIGHT PROTECTED DOCUMENT**

© ISO 2020

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office CP 401 • Ch. de Blandonnet 8 CH-1214 Vernier, Geneva Phone: +41 22 749 01 11 Email: copyright@iso.org Website: www.iso.org

Published in Switzerland

Co	Contents Pa				
Fore	eword		vi		
Intr	oductio	n	vii		
1	Scone	e	1		
2	•	native references			
3		s and definitions			
4	Bene	fits of biosolids land application	4		
5		ents in biosolids			
	5.1	General			
	5.2	Nitrogen			
		5.2.2 Nitrogen content and availability in biosolids			
		5.2.3 Determination of biosolids application rate according to nitrogen content			
	5.3	Phosphorus			
	5.4	Potassium	6		
	5.5	Calcium			
	5.6	Sulfur	6		
6	Nuisa	ance and risks associated with biosolids land application	7		
	6.1	General Odours iTeh STANDARD PREVIEW	7		
	6.2	Odours 1	7		
	6.3 6.4	Vector attraction Pathogens (standards.iteh.ai)	/		
	0.4	6.4.1 General	7 7		
		6.4.2 Viruses ISO 19698-2020			
		6.4.3 http://Bacterials.iteh.ai/eatalog/standards/sist/af6c286c-db36-4c8b-aa18-			
		6.4.4 Protozoa211d71b79920/jso=19698=2020	8		
		6.4.5 Helminths			
	6.5	Trace elements			
	6.6	Organic compounds			
		6.6.1 General 6.6.2 Source control of potential contaminants			
	6.7	Environmental considerations			
	0.7	6.7.1 General			
		6.7.2 Climate and season	10		
		6.7.3 Topography			
		6.7.4 Protection of water sources			
	6.0	6.7.5 Identification of sensitive uses and associated setbacks			
	6.8	Biosolids treatment			
		6.8.2 Foreign matter			
		6.8.3 Odour reduction			
		6.8.4 Vector attraction reduction			
		6.8.5 Pathogen reduction			
	6.9	Biosolids quality criteria — Groups of biosolids			
		6.9.1 General			
		6.9.2 Group 1 biosolids			
	6.10	Potential uses			
	6.11	Biosolids and soil monitoring			
7	Ringe	olids application programme development and management			
,	7.1	General			
	7.2	Community consultation			
		7.2.1 Involving the community	16		

# ISO 19698:2020(E)

		7.2.2 Consultation principles	
	7.3	Programme development	18
		7.3.1 General	
		7.3.2 Programme design and decision considerations	
	7.4	Programme management	
		7.4.1 General	
		7.4.2 Programme management considerations	
		7.4.3 Biosolids and soil monitoring	
		7.4.4 Continuous improvement	
		7.4.5 Voluntary agreement and quality assurance	Z4
8	Obje	ctives of agricultural land application	26
	8.1	General	
	8.2	Agronomic considerations	
		8.2.1 Nitrogen management — General	
		8.2.2 Nitrogen in biosolids	
	8.3	Phosphorus management	
		8.3.1 General	
	8.4	8.3.2 Determination of biosolids application rate according to phosphorus content Annual biosolids application rate	
9	Stora	ge, staging, fencing and signage	21
,	9.1	General	
	9.2	Signage	
10	Haula	age and field deliveries STANDARD PREVIEW	33
11	Appli	ication techniques	33
12	Speci	fic pathogen mitigation measures for agricultural land application	34
13			
13	13.1	rd keeping ISO 19698:2020	34 2 <i>1</i>
	13.2	General https://standards.iteh.a/catalog/standards/sist/af6c286e-db36-4c8b-aa18- Spreading records 211d71b79920/iso-19698-2020	3±
	13.3	Field inspection 211d71b79920/80-19698-2020	35
1 4		-	
14	14.1	ctives of land reclamation	
	14.1	Site management	
	14.2	14.2.1 General	
		14.2.2 Soil/substrate properties	
	14.3	Environmental considerations	
	11.0	14.3.1 Climate and season	
		14.3.2 Topography	
		14.3.3 Protection of water sources	
		14.3.4 Identification of sensitive uses and associated buffer zones	
	14.4	Operations management	39
		14.4.1 Site preparation	
		14.4.2 Biosolids batch management and transport	
		14.4.3 Substrate sampling	41
		14.4.4 Substrate analysis	
	14.5	Biosolids applications equipment and considerations	
	14.6	Determination of biosolids application rate	
		14.6.1 General	
		14.6.2 Biosolids application rates based on agronomic nitrogen application rate	
		14.6.3 Biosolids application rates based on maximum nutrient loads	
		14.6.4 Biosolids application rates based on target carbon to nitrogen ratio (C:N)	
		14.6.5 Biosolids application rates based on target organic matter	
	147	14.6.6 Biosolids application rates based on target pH adjustment	
	14.7 14.8	Revegetation Environmental post application monitoring	
	14.0	14.8.1 General	
		Thor denotal	тЭ

		14.8.2 Soil monitoring	
		14.8.3 Water monitoring	
	14.9	14.8.4 Foliage monitoring Quality assurance	
4 =			
15		nce and risk management for biosolids use for land reclamation	
16		lids treatment	
	16.1	Biosolids quality criteria	
	16.2 16.3	Pathogens	
	16.4	Odour	
	16.5	Fencing and signage	
	16.6	Record keeping	48
Annex	A (info	ormative) <b>Soil carbon dynamics</b>	49
Annex	<b>B</b> (info	ormative) Benefits of biosolids land application	51
Annex		ormative) Average concentrations of organic matter and plant macronutrients solids	54
Annex		ormative) Comparative pathogen and indicator limits for Group 1 biosolids	
	•	ormative) Source of trace elements in wastewater and biosolids and examples	
7 1111102		ce elements standards in biosolids and in soil after biosolids application	56
Annex	<b>F</b> (info	ormative) Standards for maximum concentration of organic compounds in	
	biosol	lids iTeh STANDARD PREVIEW	62
Annex	<b>G</b> (info	ormative) Organic compounds	63
Annex	<b>H</b> (info	ormative) <b>Organic compounds</b> ( <b>Standards.iteh.ai</b> ) ormative) <b>Setbacks (buffer zones) in different regions</b>	64
Annex	I (info	rmative) Biosolids quality based on treatment method	66
Annex	J (info	rmative) <b>Determination of plant available phosphate</b>	67
Annex	<b>K</b> (info	ormative) <b>Determination of maximum biosolids application rates based on</b>	
		elements	68
Annex	<b>L</b> (info	ormative) Examples of source control/pollution prevention tools and	
		nting industrial discharges	73
Annex	<b>M</b> (inf	ormative) Biosolids sampling frequencies	74
Annex	<b>N</b> (info	ormative) Soil and biosolids tests and methodologies	75
Annex	<b>o</b> (info	ormative) Community consultative workshop	82
Annex	<b>P</b> (info	ormative) Benefits and value of biosolids use in rehabilitation	84
		ormative) Examples of restricted activities and withholding periods for	
		2 biosolids	85
Annex	<b>R</b> (info	ormative) <b>Revegetation</b>	86
Biblio	graphy	7	87

# **Foreword**

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see <a href="www.iso.org/directives">www.iso.org/directives</a>).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see <a href="https://www.iso.org/patents">www.iso.org/patents</a>).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see <a href="https://www.iso.org/iso/foreword.html">www.iso.org/iso/foreword.html</a>. (standards.iteh.ai)

This document was prepared by Technical Committee ISO/TC 275, *Sludge recovery, recycling, treatment* ISO 19698:2020
https://standards.iteh.ai/catalog/standards/sist/af6c286e-db36-4c8b-aa18-

Any feedback or questions on this document should be directed to the user's national standards body. A

Any feedback or questions on this document should be directed to the user's national standards complete listing of these bodies can be found at www.iso.org/members.html.

# Introduction

Biosolids are organic-based materials from industrial or municipal sludge and municipal biosolids derived products, in the form of solids, semi-solids, semi-liquids, and liquids which have been treated to meet applicable standards, guidelines or requirements including the reduction of pathogens, vector attraction and contaminant criteria.

The land application of biosolids, which is considered an integrated approach to sustainable management of this resource, can be beneficial in many ways such as:

- increasing soil organic matter;
- increasing biological activity in soil;
- decreasing soil bulk density and improving soil porosity;
- improving water infiltration rate, water holding capacity and erosion prevention;
- improving soil aggregate stability;
- increasing cation exchange capacity, which can result in a lower frequency of fertilizer application;
- increasing soil pH;
- providing additional nutrients to the soil for plant growth;
- recovering phosphorus from the urban and industrial environment;
- providing potential for carbon sequestration in soil; and in providing potential for carbon sequestration in soil; and in providing potential for carbon sequestration in soil; and in providing potential for carbon sequestration in soil; and in providing potential for carbon sequestration in soil; and in providing potential for carbon sequestration in soil; and in providing potential for carbon sequestration in soil; and in providing potential for carbon sequestration in soil; and in providing potential for carbon sequestration in soil; and in providing potential for carbon sequestration in soil; and in providing potential for carbon sequestration in soil; and in providing potential for carbon sequestration in the sequestration in soil; and in providing potential for carbon sequestration in the sequestrat
- decreasing the use of mineral fertilizers and related greenhouse gas (GHG) emissions related to production and application of mineral fertilizers and related greenhouse gas (GHG) emissions related to production and application of mineral fertilizers.

This document does not prioritize, or suggest a hierarchy amongst various beneficial use options, but aims to identify and address the different criteria that could be considered to develop a sustainable and environmentally successful land application programme. These criteria include the nature of the treatment process, the selection of an appropriate application site, the method of application, the rate of application and the establishment of protective barriers or setbacks to environmentally and socially sensitive areas such as surface water and residences.

Control of non-beneficial substances, odour and potential risk to human, animal and environmental health are important parts of any beneficial use strategy. These can be managed by employing tools such as point source control, appropriate treatment methods and land-use restrictions.

Application of this document presupposes awareness of applicable legal requirements.

# iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO 19698:2020 https://standards.iteh.ai/catalog/standards/sist/af6c286e-db36-4c8b-aa18-211d71b79920/iso-19698-2020

# Sludge recovery, recycling, treatment and disposal — Beneficial use of biosolids — Land application

# 1 Scope

This document provides guidance on the conditions of beneficial use of biosolids produced from industrial and municipal sludge and municipal biosolids derived products (e.g. composts, growing media) in the production of food and feed crops, energy crops, forestry crops and for the remediation of disturbed sites.

This document applies to biosolids for land application and includes biosolids from wastewater treatment (municipal, industrial and private onsite systems).

This document does not apply to hazardous sludge that originates from wastewater which, due to its nature, physical, chemical or infectious properties, is potentially hazardous to human health and/or the environment during use, handling, storage or transportation and which requires special disposal techniques to eliminate or reduce the hazard.

This document includes:

- general guidelines for the land application of biosolids and biosolids derived products;
- specific guidelines for the land application of biosolids and biosolids derived products for food and feed crop production and for non-food and non-feed crop production (e.g. horticulture, fibre for bio-mass, silviculture, etc.); and

ISO 19698:2020

— specific guidelines: for the land application of biosolids and biosolids derived products for other beneficial uses (e.g. land reclamation of rehabilitation) 20

### 2 Normative references

There are no normative references in this document.

# 3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <a href="https://www.iso.org/obp">https://www.iso.org/obp</a>
- IEC Electropedia: available at <a href="http://www.electropedia.org/">http://www.electropedia.org/</a>

# 3.1

# alkaline stabilized biosolids

biosolids or non-toxic sludge which has undergone alkaline treatment to meet specific requirements for reduction of pathogens and vector attraction

# 3.2

# alkaline treatment

process where biosolids or non-toxic sludge is mixed with alkaline additives to enhance wastewater solids stabilization by increasing the pH of the biosolids up to 12 or higher for a minimum amount of time

# ISO 19698:2020(E)

## 3.3

### batch

definite quantity of material manufactured or produced under conditions which are presumed to be uniform

# 3.4

### biocenosis

community of biologically integrated and interdependent plants and animals

### 3.5

### biosolids

organic-based materials from industrial or municipal wastewater sludge and their derived products, in the form of solids, semi-solids, semi-liquids (pasty), and liquids which have been treated to meet specific standards, guidelines or requirements including the reduction of pathogens, vector attraction and contaminant criteria

### 3.6

# cation exchange capacity

measure of the soil's ability to hold positively charged ions (cations)

### 3.7

### compost

organic soil improver obtained by decomposition of a mixture consisting principally of various plant residues, occasionally with organic materials of animal origin, and having a limited mineral content

# [SOURCE: ISO 8157:2015, 2.2.8.6]eh STANDARD PREVIEW

# 3.8

# composting

(standards.iteh.ai)

natural aerobic biological process, carried out under controlled conditions, which converts organic material into a stable humus–like product ISO 19698:2020 https://standards.iteh.ai/catalog/standards/sist/af6c286e-db36-4c8b-aa18-

Note 1 to entry: During the composting process, Various microofganisms, including bacteria and fungi, break down organic material into simpler substances.

### 3.9

# contaminant

biological, chemical, physical, or radiological substance released to the environment from anthropogenic sources which, in sufficient concentration, can adversely affect living organisms through air, water, soil, and/or food

# 3.10

# dewatered biosolids

biosolids that have undergone a reduction of the water content to produce paste-like biosolids or solid biosolids by the use of one or several technologies, usually by natural or mechanical means

Note 1 to entry: These treatments lead to the production of biosolids whose mechanical characteristics allow a storage in heap on a minimum height of 1 m. As an indication, the dryness obtained is generally within a range between 15 % and 40 % (wet mass).

# 3.11

# foreign matter

material of anthropogenic origin as opposed to natural objects such as sand, stones and wood fibres

EXAMPLE Plastics, glass, metal, small/large or sharp debris.

# 3.12

### industrial sludge

mixture of water and solids separated from various types of industrial wastewater (e.g. food processing plants) as a result of natural or artificial processes and meet specific standards, guidelines or requirements including the reduction of pathogens, vector attraction and contaminant criteria

## 3.13

# mineralization

final stage of the biodegradation of organic matter or organic substances into carbon dioxide, water and the hydrides, oxides and other mineral salts

[SOURCE: ISO 11074:2015, 3.3.19]

### 3.14

# mesophilic anaerobic digestion

biological conversion of organic matter to biogas and residual solids at temperatures between 20 °C and about 40 °C, typically 37 °C with a mean residence time of 15 to 30 days

### 3.15

### micronutrient

element, such as boron, manganese, iron, zinc, copper, molybdenum, cobalt, and/or chlorine, which are essential, in relatively small quantities, for plant growth

[SOURCE: ISO 8157:2015, 2.1.3.3, modified — Note 1 to entry has been deleted.]

### 3.16

# municipal biosolids

biosolids produced from municipal sludge which has been treated to meet jurisdictional standards, guidelines or requirements including the reduction of pathogens and vector attraction

### 3.17

# municipal sludge

mixture of water and non-stabilized solids separated from various types of municipal wastewater as a result of natural or artificial processes ndards.iteh.ai)

### 3.18

# organic matter

# ISO 19698:2020

matter consisting of plant and or animal organic materials, and the conversion products of those 211d71b79920/iso-19698-2020 materials

[SOURCE: ISO 11074:2015, 2.1.8]

# 3.19

# organic compound

any of a large class of chemical compounds in which one or more atoms of carbon are covalently linked to atoms of other elements, most commonly hydrogen, oxygen or nitrogen

### 3.20

### plant nutrient

chemical element, which is essential for plant growth

[SOURCE: ISO 8157:2015, 2.1.2]

# 3.21

# sample

part of a defined bulk product taken for the purpose of characterization

[SOURCE: ISO 14488:2007, 3.8]

# 3.22

# setback

## buffer

determined distance, sometimes based on risk assessment, that provides protection to environmentally sensitive features such as humans and water

# ISO 19698:2020(E)

## 3.23

## sludge

mixture of water and solids originating from various types of wastewater during natural and artificial treatment

# 3.24

# sodium adsorption ratio

measure of the amount of sodium (Na) relative to calcium (Ca) and magnesium (Mg) in the water extract from saturated soil paste

### 3.25

# soil quality

all current positive or negative properties with regard to soil utilization and soil functions

[SOURCE: ISO 11074:2015, 2.1.15]

### 3.26

### stakeholder

person or organization that can affect, be affected by or perceive themselves to be affected by a decision or activity

[SOURCE: ISO 28007-1:2015, 3.6, modified — Note 1 to entry has been deleted.]

### 3.27

# thermophilic anaerobic digestion

biological conversion of organic matter to biogas and residual solids, that takes place between 49  $^{\circ}\text{C}$  and 57  $^{\circ}\text{C}$ 

# 3.28

# (standards.iteh.ai)

# total organic carbon

TOC

ISO 19698:2020

amount of carbon found in an organic compound by standards/sist/af6c286e-db36-4c8b-aa18-211d71b79920/iso-19698-2020

# 3.29

### trace element

element present in very low concentrations

# 3.30

### vector

living organism capable of transmitting a pathogen from one organism to another either mechanically (by simply transporting the pathogen) or biologically by playing a specific role in the life cycle of the pathogen

[SOURCE: EPA/625/R-92/013 Revised July 2003: Control of Pathogens and Vector Attraction in Sewage Sludge]

# 3.31

# vector attraction reduction

treatment processes that stabilize and reduce the odours and other aspects of biosolids that attract flies, rodents and other potential vectors

# 4 Benefits of biosolids land application

The land application of biosolids can be beneficial in many ways including:

- adding organic matter to the soil which can have a positive impact on soil biological activity, soil
  porosity, soil bulk density, soil water infiltration rate, aggregate stability and cation exchange
  capacity;
- increased soil organic matter can also reduce soil erosion;

- potential to increase carbon sequestration in the soil while reducing chemical fertilizer input and greenhouse gas emissions from their production (see <u>Annex A</u>); and
- responsible utilization of essential macro and micro nutrients that could otherwise be wasted.

Alkaline stabilized biosolids, in the context of the many positive soil effects, can also be an economical and effective soil amendment to increase the pH of acidic soils.

The use of biosolids should go through a global assessment taking into account the various factors, in order to ensure the positive effect on crops and soils.

See Annex B for further information regarding benefits of biosolids.

# 5 Nutrients in biosolids

## 5.1 General

Biosolids contain organic matter and plant nutrients. See <u>Annex C</u> for average concentrations of organic matter and plant macronutrients in biosolids.

# 5.2 Nitrogen

# 5.2.1 General

Growing plants require a continuous source of nitrogen (N), which is an essential component of proteins that build cell material and plant tissue. It is also necessary for other plant functions.

Nitrogen applied to soils in mineral fertilizers, biosolids or other organic amendments is subjected to biochemical and physical processes which form the nitrogen cycle. Inorganic forms of nitrogen are the available forms to plants. All forms of nitrogen can be present in the soil at any point of time, mainly because nitrogen readily shifts from one form to another:020

# 5.2.2 Nitrogen content and availability in biosolids

Nitrogen content and availability in biosolids can vary greatly depending on the source of the wastewater and the treatment process. Biosolids produced from some industrial sludges can have a high content of nitrogen while it is the opposite for others such as paper mill biosolids. Forms of nitrogen that can be present in biosolids include organic nitrogen (i.e. nitrogen bound in organic molecules such as proteins), nitrate  $(NO_3^-)$ , nitrite  $(NO_2^-)$ , ammonia  $(NH_3)$  and ammonium  $(NH_4^+)$ .

As plants can only assimilate mineral nitrogen, a primary factor in determining the nutrient value of biosolids should be the mineralization rate of its organic nitrogen. The nitrogen mineralization rate is dependant in part on the sludge treatment process. Other factors that can influence the availability of the nitrogen are intrinsic to the land application sites such as:

- temperature (air and soil);
- moisture;
- soil porosity;
- pH and texture;
- microbial activity; and
- method of application to the land.

Two main mechanisms for nitrogen loss should also be considered:

volatilisation of the ammonia; and

nitrate leaching.

# 5.2.3 Determination of biosolids application rate according to nitrogen content

The rate for biosolids application should be determined to provide the amount of nitrogen needed by the crop vegetation, or in reclamation scenarios, by the biosolids application management plan, to attain a desired result. For information regarding application rates for reclamation projects see 14.6. Whatever the project target is, special care should be applied to minimize or manage the amount of nitrogen that could be leached below the root zone of the crop, or through the soil to the ground water. Specifics related to nitrogen management on agricultural and non-agricultural lands are further explored in 8.2.2 and 14.6.2.

Biosolids application rates should be adjusted to avoid excess trace elements or phosphorus loading. In such case, if insufficient nitrogen is applied in the biosolids, inorganic fertilizer nitrogen can be applied. The addition of biosolids in a fertility plan adds multiple beneficial aspects.

# 5.3 Phosphorus

Phosphorus (P) is a macronutrient that is present in many organic materials including biosolids, and is important for healthy plant development. Specifically, phosphorus is needed for plant growth, rigidity of cell walls, and for the development of the root system. Phosphorus is of particular value as it is a limited natural resource.

Phosphorus is present in biosolids in organic and inorganic (phosphate) forms. Organic phosphorus undergoes mineralization through a (bio)degradation process in the soil before plant assimilation. Inorganic phosphorus is often predominant in biosolids [6].

The solubility and availability of phosphorus in biosolids is also dependant on soil pH. Plant available phosphorus is the phosphorus that is in the soil solution or is weakly adsorbed by soil particles and organic matter. Biological wastewater treatment does not change the availability of phosphorus. Nitrogen and phosphorus are found in similar concentrations in biosolids but often crop requirements are significantly lower for phosphorus. Subsequently application rates are based on the most restrictive element: nitrogen or phosphorus.

# 5.4 Potassium

Potassium (K) is soluble in wastewater and remains in the liquid stream. Therefore, potassium is not typically found in biosolids in high concentrations and the ratios K:N or K:P are much lower than with livestock manures, thus additional potassium can be required through the addition of inorganic fertilizers for soil health and plant growth.

# 5.5 Calcium

Most biosolids contain concentrations of calcium, approximately 2,1 % – 3,92 % (wet mass), similar to the content in animal manures. When lime is added to biosolids during an alkaline stabilization process, (e.g. 30 % lime addition by dry mass to the biosolids) the calcium content is increased. Applying biosolids at agronomic rates can supply a sufficient amount of calcium to correct deficiencies or the application of alkaline stabilized biosolids may be used to increase the soil pH where necessary or maintain the soil pH within a range that is optimum for plant growth.

# 5.6 Sulfur

Most biosolids contain sulfur, approximately 0,01 % to 2,42 % of  $SO_3$  (wet mass) which promotes plant growth, development and seed formation. In biosolids, sulfur exists in available and slow–release forms resulting from the oxidation of sulphides and decomposition of organic matter respectively. When biosolids are applied at agronomic rates the sulfur demand of the crops can also be met.

# 6 Nuisance and risks associated with biosolids land application

## 6.1 General

The beneficial use of biosolids can pose potential nuisance and risks which should be considered to help determine appropriate mitigation measures for public health and environmental protection.

Many of the potential environmental and health risks associated with the land application of biosolids are related to storage prior to spreading. These potential risks include:

- flystrike<sup>1)</sup> in stockpiles < 8 weeks aged;</li>
- ground and surface water contamination;
- leaching of contaminants during heavy rainfall events and floods;
- ammonia volatilisation: and
- public and occupational health risks from uncontrolled access and/or inappropriate storage. transport or handling.

The following factors can influence the severity of these risks:

- quantity stored;
- biosolids quality;

Teh STANDARD PREVIEW

- length of time biosolids will be stored;
- (standards.iteh.ai)

   time of the year (that is, effects of moisture, temperature and wind gradients); and
- storage design and location.

ISO 19698:2020 https://standards.iteh.ai/catalog/standards/sist/af6c286e-db36-4c8b-aa18-

211d71b79920/iso-19698-2020

### 6.2 Odours

Biosolids are an abundant source of food for microorganisms due to the fact that they contain proteins, amino-acids and carbohydrates. The degradation of these energy sources leads to the emission of odorous compounds in organic and inorganic forms such as hydrogen sulfide, mercaptan, ammonia, amines and organic fatty acids. These compounds can be released from biosolids during treatment, storage, transport and spreading, causing trouble with the neighbourhood.

# 6.3 Vector attraction

Vectors which include insects, rodents and birds can transmit pathogens to human and other hosts, physically through contact, and biologically by playing a specific role in the life cycle of the pathogen. Pathogens in biosolids may pose a risk when they are brought in contact with humans and other susceptible hosts such as plants or animals. Vectors can be attracted to a land application site by odour and the potential for food sources.

# 6.4 Pathogens

### General 6.4.1

Biosolids can contain microorganisms, some of which could be pathogens. The species and concentrations of organisms that are present in biosolids are dependent on the wastewater treatment

<sup>1)</sup> Flystrike (Myiasis) is often caused by the green bottle fly (Lucilia sericata) and related fly species laying eggs on rabbits, sheep and other animals and is used in Australia as a measure for vector attraction reduction efficiency.