

ISO/TC 34/SC 9

Secretariat: AFNOR

Voting begins on:
2020-12-29

Voting terminates on:
2021-02-23

Microbiology of the food chain — Methods for the detection of Anisakidae L3 larvae in fish and fishery products —

Part 2: Artificial digestion method

iTeh STANDARD PREVIEW
(standards.iteh.ai)

*Microbiologie de la chaîne alimentaire — Méthodes de recherche des
larves L3 d'Anisakidae dans les poissons et produits de la pêche —*

Partie 2: Méthode de digestion artificielle

<https://standards.iteh.ai/catalog/standards/sist/3019ffb6-f45b-4fd4-a7e9-50690f3a99ec/iso-fdis-23036-2>

RECIPIENTS OF THIS DRAFT ARE INVITED TO SUBMIT, WITH THEIR COMMENTS, NOTIFICATION OF ANY RELEVANT PATENT RIGHTS OF WHICH THEY ARE AWARE AND TO PROVIDE SUPPORTING DOCUMENTATION.

IN ADDITION TO THEIR EVALUATION AS BEING ACCEPTABLE FOR INDUSTRIAL, TECHNOLOGICAL, COMMERCIAL AND USER PURPOSES, DRAFT INTERNATIONAL STANDARDS MAY ON OCCASION HAVE TO BE CONSIDERED IN THE LIGHT OF THEIR POTENTIAL TO BECOME STANDARDS TO WHICH REFERENCE MAY BE MADE IN NATIONAL REGULATIONS.

ISO/CEN PARALLEL PROCESSING



Reference number
ISO/FDIS 23036-2:2020(E)

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO/FDIS 23036-2

<https://standards.iteh.ai/catalog/standards/sist/3019ffb6-f45b-4fd4-a7e9-50690f3a99ec/iso-fdis-23036-2>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2020

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

Contents

Page

Foreword	iv
Introduction	v
1 Scope	1
2 Normative references	1
3 Terms and definitions	1
4 Principle	2
4.1 General.....	2
4.2 Sample size.....	2
4.3 Sample preparation.....	2
4.4 Digestion of the sample.....	2
4.5 Filtration of the digest fluid.....	2
4.6 Verification of findings.....	2
5 Reagents	3
6 Equipment and consumables	3
7 Sampling	4
8 Procedure	4
8.1 Preparation of the sample.....	4
8.2 Preparation of the digest fluid.....	4
8.3 Digestion of the sample in the glass beaker.....	5
8.4 Filtration of the digest fluid.....	5
8.5 Microscopic examination.....	5
9 Expression of the results	5
10 Performance characteristics of the method	6
11 Test report	6
12 Quality assurance	6
Annex A (informative) Sample collection	7
Annex B (informative) Example of a laboratory worksheet for recording data when testing fish fillets with the artificial digestion method	8
Bibliography	9

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 34, *Food products*, Subcommittee SC 9, *Microbiology*, in collaboration with the European Committee for Standardization (CEN) Technical Committee CEN/TC 275, *Food analysis – Horizontal methods*, in accordance with the Agreement on technical cooperation between ISO and CEN (Vienna Agreement).

A list of all parts in the ISO 23036 series can be found on the ISO website.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

Introduction

Nematodes of the Anisakidae family have a complex life cycle involving a high number of hosts. Adult stages of Anisakidae reside in the stomach of marine mammals, where they are embedded in the mucosa. Unembryonated eggs produced by adult females are released with the faeces of marine mammals and become embryonated in seawater, where first-stage larvae (L1) develop in the eggs. The larvae moult to become free-swimming second-stage larvae (L2) and, if ingested by crustaceans, mature into third-stage larvae (L3). This stage is infective to fish and squid, and larvae are transferred between fishes through predation, maintaining the L3 stage. Some larvae migrate from the abdominal cavity into muscle tissues. Humans are incidental hosts and can be infected after ingesting raw or undercooked infected fish or cephalopods containing viable L3.

Nematodes of the family Anisakidae are the causative agents of human anisakidosis, a disease that is not only a public health hazard affecting humans, but also represents an economic problem in fishery and food safety (the term “anisakiasis”, designating the disease caused by members of the genus *Anisakis*, is also sometimes used). Worldwide, marine and wild anadromous fishes are intermediate hosts of Anisakidae, whereas marine mammals are the definitive hosts.

Visual inspection procedures for the detection of Anisakidae larvae in fish are employed to minimize the risk that contaminated fish will reach the consumer,^{[1],[2]} thus preventing human anisakidosis.

The UV-press and the artificial digestion of the fish muscular tissue are the methods specifically designed to detect nematode larvae in fish and to evaluate the infestation level of a batch, and have been validated and tested in multicentre collaborative studies^[3] (see [Clause 11](#)).

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[ISO/FDIS 23036-2](#)

<https://standards.iteh.ai/catalog/standards/sist/3019ffb6-f45b-4fd4-a7e9-50690f3a99ec/iso-fdis-23036-2>

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO/FDIS 23036-2

<https://standards.iteh.ai/catalog/standards/sist/3019ffb6-f45b-4fd4-a7e9-50690f3a99ec/iso-fdis-23036-2>

Microbiology of the food chain — Methods for the detection of Anisakidae L3 larvae in fish and fishery products —

Part 2: Artificial digestion method

1 Scope

This document specifies a method for the detection of Anisakidae L3 larvae commonly found in marine and anadromous fishes. The method is applicable to fresh fish and/or frozen fish, as well as lightly processed fish products, such as marinated, salted or smoked. It is also suitable for visceral organs as a confirmatory method for a visual inspection scheme.

The artificial digestion method^{[4]–[6]} is applicable to quantifying parasitic infections by estimating the number of parasites in the fish musculature and, when applied to fresh fish or lightly processed fish products (never frozen before processing), determining the viability of Anisakidae L3, which can be present.

This method does not apply to determining the species or genotype of detected parasites. Final identification is made by morphological and/or molecular methods.

2 Normative references

ISO/FDIS 23036-2

<https://standards.iteh.ai/catalog/standards/sist/3019ffb6-f45b-4fd4-a7e9-50690f3a99ec/iso-fdis-23036-2>

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

ISO 7218, *Microbiology of food and animal feeding stuffs — General requirements and guidance for microbiological examinations*

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <http://www.electropedia.org/>

3.1

Anisakidae L3 larvae

third-stage larvae (L3) belonging to the Anisakidae family, in particular to *Anisakis*, *Contracaecum* and *Pseudoterranova* genera

Note 1 to entry: For practical purposes, the genus *Hysterothylacium* belonging to Raphidascaerididae family and already classified as Anisakidae, can be also included.

3.2 artificial digestion method

method to detect Anisakidae larvae in fish muscle tissue by an enzymatic (pepsin-HCl) digestion step to release larvae from muscle tissues followed by a filtration step and detection of larvae by microscopy

Note 1 to entry: If applied to fresh fish, it allows checking of the viability of larvae.

4 Principle

4.1 General

The artificial digestion method relies on the enzymatic degradation of muscle fibres in a fluid composed of pepsin and hydrochloric acid followed by filtration and washing steps.

The procedure allows a differentiation between dead and viable anisakid larvae if the temperature of the digestion solution does not exceed 37 °C (with the exception of *Hysterothylacium* sp. larvae, which are killed at 37 °C), and assuming the fish was never frozen.

There are no internal quality controls that can be used while performing the method.

NOTE Alternative methods can be used for analysis, provided their equivalence with the methods described in this document are demonstrated.

4.2 Sample size

For inspection of fishery products for public health purposes, the sample size shall be risk-based, as determined by the competent authority.

4.3 Sample preparation

To increase the surface area for enzymatic degradation, samples are gently eased apart taking care not to disrupt larvae by checking for them. Alternatively, a smasher/stomacher, that facilitates the digestion but does not damage the nematode larvae, can be used.

A blending or grinding procedure should be avoided as this can damage or disrupt larvae.

4.4 Digestion of the sample

Viable Anisakidae larvae are resistant to the pepsin-HCl digest fluid and therefore can be recovered free from muscle tissues.

To facilitate an efficient and rapid digestion, a maximum ratio of 1:20, meat to digest fluid, and a temperature of 37 °C ± 2 °C shall be maintained throughout the process. The time required for digestion shall be 15 min to 30 min. For muscle samples that are less digestible, the digestion time should be increased but, unless otherwise validated for a particular sample matrix, the time shall not exceed 45 min.

4.5 Filtration of the digest fluid

Following digestion, the digest fluid shall be filtered through a sieve with specific mesh (6.9). Retained larvae shall be rinsed with tap water.

4.6 Verification of findings

If positive or doubtful findings occur, confirmation and identification at the species level should be performed by a qualified reference laboratory, by means of morphological and/or molecular methods.

5 Reagents

5.1 **Tap water.**

5.2 **Hydrochloric acid, 25 %.**

5.3 **Pepsin**, powder or granular: 1:10 000 NF, 1:12 500 BP, 2 000 FIP; liquid: 660 U/ml.

Pepsin is commercially available in powder, granular and liquid forms. The activity of the pepsin shall be certified. The pepsin shall be stored according to the manufacturer's recommendation.

The use of a liquid pepsin formulation can be advantageous as it could reduce the risk of occupational hazard, such as allergic reaction in laboratory staff.

NOTE The activity of pepsin powder is expressed per gram, either in "NF" (US National Formulary), "BP" (British Pharmacopoeia) or "FIP" (Fédération Internationale de Pharmacie). The activity of liquid pepsin is expressed in European Pharmacopoeia units per millilitre with a minimum of 660 U/ml. Other pepsin activities can be used, provided the final activity in the digest fluid is equivalent to the activity of 10 g of 1:10 000 NF.

5.4 **90 % ethanol.**

5.5 **Acidic solution, 1 % acetic acid.**

5.6 **Tap water**

iTeh STANDARD PREVIEW
(standards.iteh.ai)

6 Equipment and consumables

Usual microbiological laboratory equipment (in accordance with ISO 7218) and, in particular, the following.

6.1 **Labelled collection trays or plastic bags**, for samples.

6.2 **Knives, scissors and forceps**, for cutting samples.

6.3 **Calibrated scale**, for weighing samples and/or pepsin, accuracy $\pm 0,1$ g.

6.4 **Magnetic stirrer**, with an adjustable heating plate or magnetic stirrer put in an incubator.

6.5 **Thermometer**, accurate to at least $\pm 0,5$ °C, minimum range of temperatures from 20 °C to 70 °C.

6.6 **Polytetrafluoroethylene (PTFE)-coated triangular stir bar**, 5 cm in length, minimum.

6.7 **Glass beakers**, appropriate capacity.

6.8 **Aluminium foil or lids**, to cover the top of the glass beaker.

6.9 **Sieve**, made of brass or stainless steel, mesh size around 180 microns to 500 microns, with diameter of approximately 10 cm or larger.

6.10 **Petri dishes**, approximately 90 mm in diameter, for larval counting.

6.11 **Small vials**, for the collection of recovered larvae.