

FINAL
DRAFT

INTERNATIONAL
STANDARD

ISO/FDIS
23190

ISO/TC 249

Secretariat: SAC

Voting begins on:
2021-03-02

Voting terminates on:
2021-04-27

Traditional Chinese medicine — Determination of aristolochic acids in natural products by high-performance liquid chromatography (HPLC)

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[ISO/FDIS 23190](https://standards.iteh.ai/catalog/standards/sist/ba7e745e-d951-40d3-a331-c5d0d6da76d2/iso-fdis-23190)

<https://standards.iteh.ai/catalog/standards/sist/ba7e745e-d951-40d3-a331-c5d0d6da76d2/iso-fdis-23190>

RECIPIENTS OF THIS DRAFT ARE INVITED TO SUBMIT, WITH THEIR COMMENTS, NOTIFICATION OF ANY RELEVANT PATENT RIGHTS OF WHICH THEY ARE AWARE AND TO PROVIDE SUPPORTING DOCUMENTATION.

IN ADDITION TO THEIR EVALUATION AS BEING ACCEPTABLE FOR INDUSTRIAL, TECHNOLOGICAL, COMMERCIAL AND USER PURPOSES, DRAFT INTERNATIONAL STANDARDS MAY ON OCCASION HAVE TO BE CONSIDERED IN THE LIGHT OF THEIR POTENTIAL TO BECOME STANDARDS TO WHICH REFERENCE MAY BE MADE IN NATIONAL REGULATIONS.



Reference number
ISO/FDIS 23190:2021(E)

© ISO 2021

iTeh STANDARD PREVIEW (standards.iteh.ai)

ISO/FDIS 23190

<https://standards.iteh.ai/catalog/standards/sist/ba7e745e-d951-40d3-a331-c5d0d6da76d2/iso-fdis-23190>



COPYRIGHT PROTECTED DOCUMENT

© ISO 2021

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office
CP 401 • Ch. de Blandonnet 8
CH-1214 Vernier, Geneva
Phone: +41 22 749 01 11
Email: copyright@iso.org
Website: www.iso.org

Published in Switzerland

Contents

	Page
Foreword.....	iv
Introduction.....	v
1 Scope.....	1
2 Normative references.....	1
3 Terms and definitions.....	1
4 Abbreviated terms.....	1
5 Principle.....	2
6 Reagents.....	2
7 Apparatus.....	3
8 Test method.....	3
8.1 General principle.....	3
8.2 Caution and safety considerations.....	3
8.3 Stock solution of standard.....	3
8.4 HPLC condition.....	3
8.4.1 Chromatographic system and system suitability.....	3
8.4.2 Standard solution.....	4
8.4.3 Test solution I.....	4
8.4.4 Test solution II.....	4
8.4.5 Procedure.....	5
8.5 Method validation.....	5
8.6 Evaluation of qualitative results.....	5
9 Sampling and preservation of samples.....	5
9.1 Sampling.....	5
10 Test report.....	5
Annex A (informative) Typical HPLC conditions.....	7
Annex B (informative) Qualitative determination of aristolochic acid I by LC-MS.....	9
Bibliography.....	11

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular, the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation of the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT), see www.iso.org/iso/foreword.html.

This document was prepared by Technical Committee ISO/TC 249, *Traditional Chinese medicine*.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at www.iso.org/members.html.

Introduction

Aristolochic acids, a class of chemical compounds with renal toxicity, carcinogenic and mutagenic toxicity, are widely distributed in over 350 species of plant from around the world, many of which have been used as natural products to treat gout, arthritis, rheumatism and acute inflammation of the skin; some species from North America have been used to treat snake bites. Clinical practice and research have confirmed that long-term use of natural products containing aristolochic acids can cause chronic renal failure and renal tubules, and natural products containing aristolochic acids have been prohibited and restricted to use in clinics in many countries. Aristolochic acid toxicity is of great concern worldwide.

Safety and efficacy are basic requirements for the use of natural medicines. Although many natural products containing aristolochic acids have been strictly controlled in clinics, some are still used as raw herbal materials or to produce manufactured products such as asarum, Kaempfer dutchmanspipe root, Herba Aristolochiae mollissimae, German birthwort, American snakeroot and Indian Aristolochia tagala. In addition, some prohibited plant medicines are easily confused or misused during manufacturing, which can cause large safety concerns in the application of natural products.

This document is beneficial for effectively supervising and reducing the toxic side effects of natural-medicine-derived products and ensuring their safety and efficacy in clinical use.

The high-performance liquid chromatography (HPLC) method is applied in organizations in such places as Europe, China, the United States of America, Japan and the Republic of Korea for the determination of aristolochic acid I, both qualitatively and quantitatively. The HPLC method is recommended internationally for the qualitative determination of aristolochic acid I in natural products.

iTeh STANDARD PREVIEW
(standards.iteh.ai)

[ISO/FDIS 23190](https://standards.iteh.ai/catalog/standards/sist/ba7e745e-d951-40d3-a331-c5d0d6da76d2/iso-fdis-23190)

<https://standards.iteh.ai/catalog/standards/sist/ba7e745e-d951-40d3-a331-c5d0d6da76d2/iso-fdis-23190>

iTeh STANDARD PREVIEW
(standards.iteh.ai)

ISO/FDIS 23190

<https://standards.iteh.ai/catalog/standards/sist/ba7e745e-d951-40d3-a331-c5d0d6da76d2/iso-fdis-23190>

Traditional Chinese medicine — Determination of aristolochic acids in natural products by high-performance liquid chromatography (HPLC)

1 Scope

This document specifies a method for using high-performance liquid chromatography (HPLC) to determine the presence of aristolochic acid I in natural products.

2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

World Health Organization, Quality control methods for herbal materials. World Health Organization, 2011

3 Terms and definitions

For the purposes of this document, the following terms and definitions apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- ISO Online browsing platform: available at <https://www.iso.org/obp>
- IEC Electropedia: available at <http://www.electropedia.org/>

3.1

raw herbal material

medicinal part of medicinal plants after *preliminary processing* (3.2)

3.2

preliminary processing

physical technique of converting medicinal part of medicinal plants into *raw herbal material* (3.1)

3.3

extract

preparation containing the active ingredient of a substance from *raw herbal material* (3.1) in concentrated form based on the standardized production process and meeting certain quality standards

4 Abbreviated terms

DAD	diode array detector
HPLC	high-performance liquid chromatography
UV	ultraviolet absorption
LC-MS	liquid chromatography-mass spectrometry

5 Principle

The HPLC method is employed to determine the presence of aristolochic acid I. The HPLC system consists of a quaternary pump, continuous vacuum degasser, thermostated auto-sampler and column compartment coupled to a variable wavelength diode array detector (DAD).

The aristolochic acid I determinable by this method is shown in [Table 1](#). The chemical structure of aristolochic acid I is given in [Figure 1](#).

Table 1 — Aristolochic acid I determinable by this method

Analyte	Molecular formula	CA determinable S ^a No.	Molar mass
Aristolochic acid I	C ₁₇ H ₁₁ NO ₇	313-67-7	341,27




Figure 1 — Chemical structure of aristolochic acid I

6 Reagents

All reagents shall be of recognized chromatographic or analytical grade. The purity of the reagents used shall be checked by running a blank determination. The chromatogram obtained from the solvents shall have a baseline without noticeable peak that would interfere with those from the targeted aristolochic acid I for detection.

6.1 Aristolochic acid. The purity of aristolochic acid I should be more than 95 %.

6.2 Acetonitrile. The molecular formula of acetonitrile is CH₃CN, with a HPLC grade for HPLC and a LC-MS grade for LC –MS.

6.3 Phosphoric acid. The molecular formula of phosphoric acid is H₃PO₄, with a HPLC grade for HPLC.

6.4 Formic acid. The molecular formula of formic acid is HCOOH, with a LC-MS grade for LC –MS.

6.5 Water. The molecular formula of water is H₂O with a resistivity of 18,2 MΩ.

6.6 Methanol. The molecular formula of methanol is CH₃OH with a purity of analytical grade.

7 Apparatus

Use the usual laboratory apparatus and, in particular, the following.

- 7.1 **HPLC**, consisting of a high-pressure infusion pump, sampler and chromatographic column, DAD, integrator or data processing system.
- 7.2 **Glassware**. All glassware shall be thoroughly cleaned before use.
- 7.3 **Grinder**, used to smash the raw herbal materials.
- 7.4 **Volumetric flasks**, with a capacity of 2,0 ml and 10,0 ml.
- 7.5 **Conical flask**, with a capacity of 100 ml and a glass stopper.

8 Test method

8.1 General principle

The HPLC-DAD method is applied to detect the presence of aristolochic acid I. The parameters can vary depending on the particular condition of different instruments. The procedures of the HPLC-DAD method can be modified if the accuracy and precision are validated, such as a new analysis method with the same or higher accuracy and precision.

8.2 Caution and safety considerations

Due to the toxicity of aristolochic acid I, necessary precautions shall be taken in all operations. The solutions and the apparatus used in the test should not pollute the environment.

8.3 Stock solution of standard

Dissolve 2 mg of aristolochic acid I in methanol or 80 % methanol and store at 4 °C.

8.4 HPLC condition

8.4.1 Chromatographic system and system suitability

a) Column


Stationary phase: octadecylsilyl bonded silica gel as chromatography column or equivalent.

b) Mobile phase

- Mobile phase A: acetonitrile.
- Mobile phase B: 0,05 % phosphoric acid solution

c) Wavelength of the detector: 254 nm.

d) The ultraviolet absorption (UV) spectrum of aristolochic acid I is shown in [Figure 2](#).

**Key**

X nm
Y mAU

iTeh STANDARD PREVIEW
(standards.iteh.ai)

Figure 2 — UV spectrum of aristolochic acid I

[ISO/FDIS 23190](#)

See [Annex A](#) for typical HPLC-DAD chromatographic conditions. [7e745e-d951-40d3-a331-c5d0d6da76d2/iso-fdis-23190](#)

8.4.2 Standard solution

Dilute the stock solution of aristolochic acid I with methanol or 80 % methanol to prepare the standard solution of aristolochic acid I (0,2 ug/ml).

8.4.3 Test solution I

Weigh 100 g of raw herbal materials to grind and pass it through an 80-mesh or finer sieve. Weigh accurately 0,5 g of the powdered raw herbal material in a conical flask with stopper. Add accurately 25 ml of 70 % methanol or 80 % methanol and weigh. Ultrasonicate (powder, 250 W; frequency, 40 kHz) for 40 minutes, cool, weigh again, make up the loss of weight with 70 % methanol or 80 % methanol, mix well and filter through a membrane filter (normal pore size 0,45 µm) using the successive filtrate as the test solution.

8.4.4 Test solution II

Grind approximately 10 g to 100 g of the extract, bolus, powder, pellet and tablet to fine powder, weigh an appropriate amount (equivalent to 0,5 g of aristolochic acid-containing raw herbal material) in a conical flask with stopper. Add accurately an appropriate amount of 70 % methanol or 80 % methanol with a ratio of material to liquid of 50:1. Ultrasonicate (powder, 250 W; frequency, 40 kHz) for 40 minutes, cool and then filter. Evaporate the filtrate to dryness, dissolve the residue in an appropriate amount of 70 % methanol or 80 % methanol to prepare a solution containing 20 mg aristolochic acid-containing raw herbal material per ml and filter through a membrane filter (normal pore size 0,45 µm) using the successive filtrate as the test solution.